

🌸 日本数学会

2018年度年会

英 文 サ マ リ 集

2018年3月

於 東京大学

2018 日本数学会 年会プログラム

期 日 2018年3月18日(日)～3月21日(水)

会 場 東京大学大学院数理科学研究科
〒153-8914 東京都目黒区駒場3-8-1

連絡先 東京大学大学院数理科学研究科
〒153-8914 東京都目黒区駒場3-8-1

E-mail tokyo18mar@mathsoc.jp

(会期中) Tel 090-1791-3483

一般社団法人 日本数学会
Tel 03-3835-3483

	第I会場 11号館1階 1101教室	第II会場 11号館1階 1102教室	第III会場 11号館2階 1106教室	第IV会場 11号館2階 1108教室	第V会場 13号館1階 1311教室	第VI会場 13号館1階 1313教室	第VII会場 13号館2階 1321教室	第VIII会場 13号館2階 1322教室	第IX会場 13号館3階 1331教室
18日 (日)	幾何学 9:15～11:35 14:15～16:35	トポロジー 9:40～12:00 15:30～18:00	代数学 9:00～11:45 14:20～17:20	函数解析学 14:15～16:15	数学基礎論 および歴史 9:00～11:30 14:10～16:50	応用数学 9:30～11:45 14:20～16:30	函数論 9:45～11:45	統計数学 9:15～12:00 14:15～15:00	函数方程式論 9:15～12:00 14:15～16:15
	企画特別講演 13:00～14:00								
	特別講演 16:45～17:45	特別講演 14:15～15:15		特別講演 16:30～17:30	特別講演 17:00～18:00	特別講演 16:45～17:45	特別講演 14:15～15:15	特別講演 15:15～16:15 16:30～17:30	特別講演 16:30～17:30
19日 (月)	幾何学 9:00～11:45	トポロジー 9:30～10:30 13:00～14:20	代数学 9:15～12:00	函数解析学 9:30～12:00	数学基礎論 および歴史 9:00～11:30	応用数学 9:30～11:45	函数論 10:00～11:40	統計数学 9:30～11:30	函数方程式論 9:15～12:00
	特別講演 13:15～14:15	特別講演 10:45～11:45	特別講演 13:15～14:15	特別講演 13:15～14:15	特別講演 13:15～14:15		特別講演 13:15～14:15		特別講演 13:15～14:15
	日本数学会賞授賞式(講堂 900番教室)……………(14:50～15:20)								
	総合講演() 日本数学会賞春季賞受賞者……………(15:30～16:30) 山口孝男(京大理)……………(16:45～17:45) 懇親会(駒場ファカルティハウス)……………(18:00～20:00)								
20日 (火)	幾何学 9:15～11:45 14:15～16:30	トポロジー 9:20～12:00 15:35～17:45	代数学 9:15～12:00	函数解析学 9:30～12:00 14:15～16:00	無限可積分系 10:00～11:30 14:00～15:30	応用数学 9:15～11:50 14:15～16:40	実函数論 9:00～12:00 14:15～16:15	統計数学 10:00～11:40 14:15～15:05	函数方程式論 9:15～12:00 14:15～16:15
	企画特別講演 13:00～14:00								
	特別講演 16:45～17:45	特別講演 14:20～15:20	特別講演 14:30～15:30 15:40～16:40 16:50～17:50	特別講演 16:15～17:15	特別講演 15:45～16:45	特別講演 16:50～17:50	特別講演 16:30～17:30	特別講演 15:20～16:20 16:40～17:40	特別講演 16:30～17:30
21日 (水)			代数学 9:15～12:00 14:20～17:00		無限可積分系 9:30～10:30	応用数学 9:15～11:55 14:15～16:40	実函数論 9:15～12:00 14:15～16:00	統計数学 10:00～12:00	函数方程式論 9:15～12:00 14:15～16:15
	企画特別講演 13:00～14:00								
					特別講演 10:45～11:45	特別講演 16:50～17:50	特別講演 16:15～17:15		特別講演 16:30～17:30

目次

総合講演	1
企画特別講演	2
数学基礎論および歴史	5
3月18日(日)	5
3月19日(月)	10
代数学	13
3月18日(日)	13
3月19日(月)	19
3月20日(火)	23
3月21日(水)	27
幾何学	33
3月18日(日)	33
3月19日(月)	36
3月20日(火)	38
函数論	43
3月18日(日)	43
3月19日(月)	45
函数方程式論	47
3月18日(日)	47
3月19日(月)	53
3月20日(火)	56
3月21日(水)	62
実函数論	68
3月20日(火)	68
3月21日(水)	71
函数解析学	76
3月18日(日)	76
3月19日(月)	78
3月20日(火)	81
統計数学	86
3月18日(日)	86
3月19日(月)	89
3月20日(火)	92
3月21日(水)	96

応用数学	98
3月18日(日)	98
3月19日(月)	102
3月20日(火)	105
3月21日(水)	110
トポロジー	116
3月18日(日)	116
3月19日(月)	120
3月20日(火)	122
無限可積分系	128
3月20日(火)	128
3月21日(水)	130

総 合 講 演

3月19日(月) 総合講演会場

日本数学会賞春季賞受賞者 (15:30~16:30)
Spring Prize Winner

山口孝男(京大理) 境界付きリーマン多様体の崩壊 (16:45~17:45)
Takao Yamaguchi (Kyoto Univ.) Collapsing Riemannian manifolds with boundary

概要 As the Gauss-Bonnet theorem shows, there are relations between curvature and topology of Riemannian manifolds, and studying such relations has been one of the main problems in Differential geometry. The theory of Gromov-Hausdorff convergence of Riemannian manifolds is known to be an effective method in this direction. In this theory, we consider a certain family of closed Riemannian manifolds with a lower curvature bound. The lower curvature bound ensures the precompactness of the family, and one can expect that some topological, geometric or analytic invariants should be bounded on that family. Moreover, it is natural to ask deeper information on manifolds themselves through the convergence. The theory of collapsing Riemannian manifolds answers this question and turns out to be quite effective in some cases. For example, the theory of collapsing Riemannian three-manifolds was used in Perelman's work on the geometrization conjecture to determine the topology of the collapsed part of a closed three-manifold under the Ricci flow. In this lecture, after a brief survey of the development of the collapsing theory for closed manifolds, I will mainly focus on recent development of the theory of collapsing Riemannian manifolds with boundary, based on joint works with Zhilang Zhang.

企 画 特 別 講 演

3月18日(日)

第VI会場

田村明久 (慶大理工) 離散凸解析と数理経済モデル (13:00~14:00)
Akihisa Tamura (Keio Univ.) Discrete convex analysis and mathematical economic models

概要 Discrete convex analysis is a unified framework of discrete optimization and two concepts, called L-convexity and M-convexity, play important roles in this framework. Structural results of discrete convex analysis include the conjugacy theorem between L-convexity and M-convexity, separation theorems for L-convex/L-concave functions and for M-convex/M-concave functions, and the Fenchel-type discrete duality theorem. Algorithmic aspects of L-convex and M-convex functions have also been discussed in discrete mathematics. In mathematical economics, discrete convex analysis has been applied to models with indivisible commodities. In this talk, I will cover these fundamental topics and recent developments.

第IX会場

八杉満利子 「竹内の証明論」について (13:00~14:00)
Mariko Yasugi On “Takeuti’s Proof Theory”

概要 Gaisi Takeuti has left us numerous distinguished works in various fields of the foundations of mathematics. Among them, “proof theory” can be proclaimed to have been the matter of his foremost concern. It was deeply related to his desire (dream) to understand the true nature of “sets”.

He formulated second order logic (and higher order ones), which is a formalized framework for mathematics, and posed a conjecture, known as “Takeuti’s fundamental conjecture”, which would provide many logical facts, including the consistency, of the formal systems. An attempt of (partially) proving the conjecture is called, as a catchword, “cut elimination” or “consistency proof”.

The fundamental conjecture claims that its proofs must be performed within “Hilbert’s Program”, that is, the metamathematical proofs of formal systems must abide by the “finitist viewpoint”. Formalism and finitist viewpoint are the important thoughts behind Takeuti’s mathematical achievements.

In order to prove the fundamental conjecture, Takeuti created and developed a theory of constructive well-ordered structures of notations, called “ordinal diagrams”. Applying this theory, he made a monumental advance of the consistency proof.

I will introduce Takeuti’s proof-theoretical results along with the background thoughts, which determined the direction of “Takeuti’s proof theory”.

3月20日(火)

第III会場

渡辺 純三 (東海大*) 次数付き0次元ゴレンスタイン環のレフシェッツ性 …… (13:00~14:00)
 Junzo Watanabe (Tokai Univ.*) The Lefschetz properties of graded Artinian Gorenstein algebras

概要 The strong Lefschetz property for Artinian Gorenstein algebras is a ring-theoretic abstraction of the Hard Lefschetz Theorem for compact Kähler manifolds.

Suppose that $A = \bigoplus_{i=0}^d A_i$ is a graded Artinian Gorenstein algebra. We say that A has the **strong Lefschetz property** if there exists a linear element $l \in A$ such that the multiplication map

$$\times l^{d-2i} : A_i \rightarrow A_{d-i}$$

is bijective for all $i = 0, 1, 2, \dots, [d/2]$.

This can be defined for graded vector spaces and basic properties can be derived as properties of an endomorphism of graded vector spaces. As a consequence, it can be proved that “almost all” Artinian Gorenstein algebras have the strong Lefschetz property. It gives us new problems to ask (1) what classes of Gorenstein algebras have the strong/weak Lefschetz property without exception and (2) what Gorenstein algebras fail to have the strong/weak Lefschetz property. I would like to speak about known results, methods to prove them, applications and problems of the Lefschetz properties of Artinian Gorenstein algebras.

第VI会場

特別招待講演 (日本応用数理学会)

岩田 覚 (東大情報理工) マトロイド・パリテイ …… (13:00~14:00)
 Satoru Iwata (Univ. of Tokyo) Matroid parity

概要 This talk provides an overview on matroid parity, which was introduced in the 70s as a common generalization of matching and matroid intersection. In particular, we present a combinatorial, deterministic, polynomial-time algorithm for the weighted linear matroid parity problem.

第IX会場

葉 廣和夫 (京大数理研) 3次元トポロジーにおける圏論的構造について …… (13:00~14:00)
 Kazuo Habiro (Kyoto Univ.) Category-theoretic structures in 3-dimensional topology

概要 We will review algebraic and category-theoretic structures in 3-dimensional topology. The most well-known structures of such kind are topological quantum field theories, which are vector-space-valued functors on the category of closed surfaces and 3-dimensional cobordisms. We will mainly focus on the category Cob of once-punctured surfaces and cobordisms between them, introduced by Crane and Yetter and by Kerler, and some of its subcategories. The category Cob has a structure of a braided monoidal category, and it is equipped with a Hopf algebra object. We will consider braided subcategories sLCob and LCob of Cob, which are called the category of special Lagrangian and Lagrangian cobordisms, respectively, where sLCob is also a subcategory of LCob. The category sLCob may be identified with the opposite of the category H of handlebodies and “disc-based” embeddings. We will also discuss functors on these categories, which may be regarded as functorial 3-manifold invariants.

3月21日(水)

第VI会場

谷崎俊之(阪市大理) 量子旗多様体と量子群の表現 (13:00~14:00)
Toshiyuki Tanisaki (Osaka City Univ.) Quantized flag manifolds and representations of quantum groups

概要 By the work of Brylinski-Kashiwara and Beilinson-Bernstein we can localize representations of complex simple Lie algebras on the flag manifolds. There is also a similar theory in positive characteristics due to Bezrukavnikov-Mirkovic-Rumynin.

In this talk I would like to talk about the corresponding results for quantized enveloping algebras using the quantized flag manifolds. The quantized flag manifold is not an algebraic variety in the ordinary sense; it is a non-commutative scheme equipped with non-commutative ring of functions. Nevertheless, we have the notion of D -modules on the quantized flag manifolds, by which we can localize representations of quantized enveloping algebras on the quantized flag manifolds.

第IX会場

田中直樹(静岡大理) 距離空間における作用素半群と微分方程式 (13:00~14:00)
Naoki Tanaka (Shizuoka Univ.) Semigroups of operators and mutational equations in metric spaces

概要 The nonlinear Hille-Yosida theorem in Hilbert spaces was established in 1967 by Kōmura. After his pioneering work, a generation theorem of semigroups of contractions in general Banach spaces was proved in 1971 by Crandall and Liggett. The notion of semigroups of Lipschitz operators in Banach spaces was introduced by Kobayashi in 1990's as a nonlinear analogue of strongly continuous semigroups of bounded linear operators. Metric-like functionals, not the metrics induced by norms, play an important role in characterizing such semigroups with continuous generators. What happens if we avoid using not only norms but also linear structures? I will talk about mutational equations described by 'transitions' and 'mutations', which are mathematical tools extending the concept of differential equations to the case of metric spaces. The mutational analysis was initiated by Aubin in 1990's to analyze varying shapes, and has been recently developed by Lorenz to give a unified way to various types of evolution equations. Their abstract results can be extended to apply to quasilinear evolution equations due to Kato.

数学基礎論および歴史

3月18日(日) 第V会場

9:00~11:30

- 1 田中昭太郎 * 分数関数をローラン展開と衰壊展開で表現し対比する 15
Shotaro Tanaka To express fractions into power series by Suida expansion

概要 The fraction $1/(z^2 + 1)$ has two singularities $z = i, -i$. Expand in $z = 2i$. (1) Devide the domain into $0 \leq |z - 2i| < 1$, $1 < |z - 2i| < 3$, $3 < |z - 2i|$. (2) Decomposite it into partials: $(i/2)\{1/(z + 1) - i/(z - i)\}$. Ex. $1/(z + 1) = 1/\{(z - 2i) + 3i\}$. Let us omit Laurent's expansion. Wada's theorem: $1/(\square - \Delta)^{p+1} = \Sigma(k = 1 \rightarrow \infty)d_p(k)\square^{-p-k}\Delta^{k-1}$, $0 < |\Delta/\square| < 1$, where integers $p \geq 0$; $k \geq 1$ and $d_p(k) \equiv (k + p - 1)!/p!(k - 1)!$, named Suida expansion. If $1 \div \{(z - 2i) - (-3i)\}^{0+1} = \Sigma(k = 1 \rightarrow \infty)d_0(k)(z - 2i)^{-0-k}(-3i)^{k-1}$, $0 < |-3i/(z - 2i)| < 1$. Then $\Sigma(k = 1 \rightarrow \infty)(-3i)^{k-1}(z - 2i)^{-k}$, $3 < |z - 2i|$. If $1/\{3i - (-(z - 2i))\}^{0+1} = \Sigma(k = 1 \rightarrow \infty)d_0(k)(3i)^{-0-k}(-(z - 2i))^{k-1}$, $0 < |-(z - 2i)/3i| < 1$. Then $\Sigma(k = 1 \rightarrow \infty)d_0(k)(-1)^{k-1}(1/3i)^k(z - 2i)^{k-1}$, $0 < |z - 2i| < 3$.

- 2 増田 茂 (京大数理研) Mathematical principles treated in mechanics by Poisson 15
Shigeru Masuda (Kyoto Univ.) Mathematical principles treated in mechanics by Poisson

概要 Maupertuis treats principle of minimum action and Clairaut discuss effort and principles of equilibrium on earth. Poisson issues Study of Mechanics in 1833, which consists the second book of three books, entitled A Study of Mathematical Physics. He discusses the mathematical principles from many side of mathematics. Our present to this session shows the points of mathematical scopes in mechanics by Poisson.

- 3 増田 茂 (京大数理研) Proof of rise of capillary surface by Poisson 15
Shigeru Masuda (Kyoto Univ.) Proof of rise of capillary surface by Poisson

概要 Providing capillary action in the equilibrium, Poisson assures that the rise of the surface of water is due to the abrupt variation of density in the neighborhood of the wall and of the surface. Poisson discusses this problem in 1831, in the rivalry to the paper/book of Laplace 1806-7 and Gauss 1831. We show Poisson's discussion.

- 4 中根美知代 20世紀初頭のハミルトン・ヤコビ理論と変換論 15
Michiyo Nakane An introduction of an idea of transformation to Hamilton-Jacobi theory in the beginning of twentieth century

概要 In the early twentieth century, mathematicians began to eagerly discuss transformations of variables that keep the canonical form of the differential equations, the so-called "contact transformation" or "canonical transformation". This paper examines origins of these two transformations. We also discuss a process how these two transformations were bound and how they were introduced to the Hamilton-Jacobi theory.

- 5 張替俊夫 『数』の斗食算題について 15
(大阪産大全学教育機構)

Toshio Harikae (Osaka Sangyo Univ.) On 'Toshoku' problems in "Shu"

概要 Two 'Toshoku (doushi)' problems contained in the book "Shu", and one of them had been left undeciphered. In this talk, we decipher the problem from the other deciphered problem, and discuss that the method used in Toshoku problems is 'shaoguang-shu.'

- 6 脇 克志 (山形大理) NMFによる和算図形問題類似評価 10
 土橋拓馬 (明大先端数理)
 阿原一志 (明大総合数理)
 Katsushi Waki (Yamagata Univ.) The similarity evaluation of geometric problems in WASAN by NMF
 Takuma Tsutiihashi (Meiji Univ.)
 Kazushi Ahara (Meiji Univ.)

概要 We try to evaluate the similarity of geometric problems in WASAN using NMF (Non-negative Matrix Factorization). In our study, we calculate characteristic vectors of geometric problems in “SANPOUTEN-SEISHINAN” written by “AIDA YASUAKI” by NMF. And we try to evaluate the similarity of geometric problems by these characteristic vectors.

- 7 森本光生 大成算経の日用術について 15
 (四日市大関孝和数学研・上智大*)
 Mitsuo Morimoto Daily Mathematics in the Taisei Sankei
 (Yokkaichi Univ./Sophia Univ.*)

概要 Volumes 8 and 9 of the Taisei Sankei (Great Accomplished Mathematical Treatise, 1710 or 11) are entitled Daily Mathematics and contain 220 problems of mathematics stemmed from daily lives. Because of their elementary character, they have been neglected by historians of Japanese mathematics; for example, in the Meijizen Hihon Sūgakushi (History of Mathematics before the Meiji Restoration), Fujiwara Matsusaburo paid almost no attention to them. Takebe Katahiro, one of three authors of the Taisei Sankei published the Sangaku Keimō Genkai Taisei (Great Colloquial Commentary on the Suanxue Qimeng) in 1690. The Suanxue Qimeng (Introduction to Mathematics, 1299), written by Zhu Shijie of the Yuan dynasty, was a collection of mathematical problems. We shall discuss the relation of the Daily Mathematics with the Suanxue Qimeng.

- 8 小川 束 (四日市大環境情報) 会田安明の数学思想 (その2) 15
 Tsukane Ogawa (Yokkaichi Univ.) Mathematical philosophy of Aida Yasuaki, part 2

概要 Aida Yasuaki (1747–1817) wrote the 6 volume *Sanpou Kokon Tsuuran* in 1797, which incisively criticized 19 already published books of mathematics. I have discussed the first volume of Aida’s work, which contained comments on 7 books from the *Sanpou Kongen Ki* (1666) to the *Katsuyou Sanpou* (1712). In this instance, I examined the second volume, which contained comments on 8 books from the *Kagaku Sanpou* (1715) to the *Meigen Sanpou* (1764). Though some comments were the same as those found in volume 1, some were new. I will comment on the history of difficult problems, and on the relationship between solutions using equations and solutions using the abacus.

- 9 土橋拓馬 (明大先端数理) 画像認識に基づく和算図形問題への自動タグ付け 15
 脇克志 (山形大理)
 阿原一志 (明大総合数理)
 Takuma Tsuchihashi (Meiji Univ.) Automatic tagging to geometric problems in Japanese mathematics
 Katsushi Waki (Yamagata Univ.) (wasan) based on image recognition
 Kazushi Ahara (Meiji Univ.)

概要 In recent years, construction of image databases from digitized historical documents of Japanese mathematics (wasan) has been progressing. Since previous studies, the authors have proposed ideas for automatic tagging in these image databases of wasan based on geometric elements (triangles, squares, circles and so on) and those relationship (tangency of circles, number of elements and so on). In this study, the authors implement the program that tags images of geometric problems in wasan automatically and verify the effectiveness of our proposal for geometric problems included in actual documents of wasan. As a result, it is found that automatic tagging succeeds for geometric problems of more than 80 percent in “Sanpo tensei-ho shinan” by Yasuaki Aida.

11:30~12:00 歴史部門懇談会

14:10~16:50

- 10 鈴木登志雄 (首都大東京理工) Non-depth-first search of an AND-OR tree 15
 Toshio Suzuki (Tokyo Metro. Univ.) Non-depth-first search of an AND-OR tree

概要 Liu and Tanaka (2007) asserted that among independent distributions on a uniform binary AND-OR tree, the minimum cost (achieved by an algorithm) is maximized only by an independent and identical distribution. In this decade, the assertion has been justified under a hypothesis that only depth-first algorithms are taken into consideration. The uniform binary tree case was shown by S. and Niida (2015). Balanced multi-branching tree case was shown (with a certain hypothesis) by Peng et al. (2017). We extend the results of S.-Niida and Peng et al. to the case where non-depth-first algorithms are taken into consideration.

- 11 田中一之 (東北大理) 重み付き AND-OR 木における固有分布の一意性について 15
 沖坂祥平 (東北大理)
 Kazuyuki Tanaka (Tohoku Univ.) On the uniqueness of the eigen-distribution for weighted AND-OR trees
 Shohei Okisaka (Tohoku Univ.)

概要 In 2007, Liu and Tanaka characterized the eigen-distributions that achieve the distributional complexity for AND-OR trees, and among others, they proved the uniqueness of eigen-distribution for a uniform binary tree. Later, Suzuki and Nakamura showed that the uniqueness fails if only directional algorithms are considered. In this talk, we introduce the weighted trees, namely, trees with weighted cost depending on the value of a leaf. Using such models, we prove that for balanced multi-branching trees, the uniqueness of eigen-distribution holds w.r.t. all deterministic algorithms, but fails w.r.t. only directional algorithms.

- 12 水澤勇気 (首都大東京理工) 偽ソロベイ還元性に関する幾つかの結果 15
 Yuki Mizusawa (Tokyo Metro. Univ.) Some results of pseudo Solovay reducibility

概要 Solovay reducibility is a well-known and important notion in theory of randomness. We defined pseudo Solovay reducibility to generalize Solovay reducibility. We have some results of pseudo Solovay reducibility.

- 13 横山 啓太 (北陸先端大情報) 指標関数, 強制法と証明の変換 15
Keita Yokoyama (JAIST) Indicators, forcing and proof-transformation

概要 The indicator argument is a model-theoretic framework to obtain independence and conservation results in the study of first-order arithmetic. In the talk, we will consider new formulation of indicator arguments with the idea of generic cuts and forcing. With this method, we will analyze the conservation results for bounding principle and Ramsey's theorem for pairs.

- 14 只木 孝太郎 (中部大工) 量子力学では確率1の事象は必ず起こる 15
Kohtaro Tadaki (Chubu Univ.) An event with probability one occurs certainly in quantum mechanics

概要 The notion of probability plays a crucial role in quantum mechanics. It appears as the Born rule. In this talk we reveal that every event with probability one occurs certainly in quantum measurement.

- 15 矢島 幸信 (神奈川大工) Undecidability of the existence of C^* -embedded but not C -embedded
平田 康史 (神奈川大工) subsets in a product of natural numbers 15
Yukinobu Yajima (Kanagawa Univ.) Undecidability of the existence of C^* -embedded but not C -embedded
Yasushi Hirata (Kanagawa Univ.) subsets in a product of natural numbers

概要 It is proved that every C^* -embedded subset in \mathbb{N}^{ω_1} is C -embedded in \mathbb{N}^{ω_1} under a certain set-theoretic assumption, where \mathbb{N}^{ω_1} denotes the product of ω_1 copies of natural numbers \mathbb{N} . As a consequence, it is independent of ZFC that there is a (closed) C^* -embedded subset in \mathbb{N}^{ω_1} .

- 16 薄葉 季路 (早大理工) Products of Lindelöf spaces 15
Toshimichi Usuba (Waseda Univ.) Products of Lindelöf spaces

概要 For a topological space X , the Lindelöf degree of X is the minimal cardinal κ such that every open cover of X has a subcover of size $\leq \kappa$. If S is the Sorgenfrey line, then its product $S \times S$ has the Lindelöf degree 2^ω . On the other hand, it is unknown whether there are Lindelöf spaces X and Y with $L(X \times Y) > 2^\omega$. In this talk, we prove that, in the Cohen forcing extension, the Lindelöf degree of the product of two regular Lindelöf spaces can be arbitrary large up to the least ω_1 -strongly compact cardinal. We also show that if there is no such Lindelöf spaces, then ω_2 is weakly compact in L .

- 17 依岡 輝幸 (静岡大理) Aronszajn 木が持つふたつの組合せ的性質の違い 15
Teruyuki Yorioka (Shizuoka Univ.) On two combinatorial properties which come from Aronszajn trees

概要 The forcing notion which adds an uncountable antichain through an Aronszajn tree has two similar combinatorial properties, which are called the rectangle refining property and the property R_{1, \aleph_1} . By a viewpoint of specialization of an Aronszajn tree, we can conclude that two properties are different.

- 18 D. A. Mejía (静岡大理) 強制法理論及び連続体上の組合せ論 15
Diego A. Mejía (Shizuoka Univ.) Forcing theory and combinatorics of the real line

概要 We present recent developments in forcing theory that have been motivated by problems on combinatorics of the real line. Concretely, the construction of tree-dimensional arrangements of forcing generic extensions (joint work with Fischer, Friedman and Montoya), and the incorporation of ultrafilter limits in two-dimensional arrangements of generic extensions (joint with Brendle and Cardona). These techniques work to construct models where the cardinals in Cichoń's diagram (classical diagram of cardinal numbers associated with combinatorial properties of the real line) can be divided into 7 different values, which is the maximum number known modulo ZFC alone.

- 19 酒井拓史 (神戸大システム情報) Embeddability of uncountable LO into models generated by uncountable indiscernible sequences 15
- Hiroshi Sakai (Kobe Univ.) Embeddability of uncountable LO into models generated by uncountable indiscernible sequences

概要 We discuss what kinds of uncountable linearly ordered sets are embeddable into models generated by uncountable indiscernible sequences.

17:00~18:00 特別講演

- 渕野昌 (神戸大システム情報) Set-theoretic reflection principles
- Sakaé Fuchino (Kobe Univ.) Set-theoretic reflection principles

概要 For a class \mathcal{C} of structures (with a fixed notion \mathcal{N} of substructure) and a property \mathcal{P} , the reflection cardinal of $(\mathcal{C}, \mathcal{P})$ is the minimal cardinal κ such that, for any $M \in \mathcal{C}$ of cardinality $> \kappa$, if M does not satisfy the property \mathcal{P} , then there are stationarily many substructures N of M of cardinality $< \kappa$. If κ is the reflection cardinal of $(\mathcal{C}, \mathcal{N})$, we shall write $\kappa = \text{Ref}(\mathcal{C}, \mathcal{P})$.

By choosing \mathcal{C} , \mathcal{N} and \mathcal{P} , we can represent many set-theoretic reflection statements. If, for example \mathcal{P} is simply a contradiction, and \mathcal{N} is the elementary submodel relation for some logic \mathcal{L} , then $\kappa = \text{Ref}(\mathcal{C}, \mathcal{P})$ is the strong form of Downward Löwenheim-Skolem Theorem down to $< \kappa$ for \mathcal{L} .

Of these reflection statements, the cases $\aleph_2 = \text{Ref}(\mathcal{C}, \mathcal{P})$ and $2^{\aleph_0} = \text{Ref}(\mathcal{C}, \mathcal{P})$ seems to be of special interest. The former may be interpreted as a pronouncement that the first uncountable cardinal \aleph_1 captures the situation $\neg\mathcal{P}$ good enough while the latter as the pronouncement that the continuum is large enough in connection with the property \mathcal{P} .

For a class \mathcal{C} of structures (with a fixed notion \mathcal{N} of substructure) and a property \mathcal{P} , the reflection cardinal of $(\mathcal{C}, \mathcal{P})$ is the minimal cardinal κ such that, for any $M \in \mathcal{C}$ of cardinality $> \kappa$, if M does not satisfy the property \mathcal{P} , then there are stationarily many substructures N of M of cardinality $< \kappa$. If κ is the reflection cardinal of $(\mathcal{C}, \mathcal{N})$, we shall write $\kappa = \text{Ref}(\mathcal{C}, \mathcal{P})$.

By choosing \mathcal{C} , \mathcal{N} and \mathcal{P} , we can represent many set-theoretic reflection statements. If, for example \mathcal{P} is simply a contradiction, and \mathcal{N} is the elementary submodel relation for some logic \mathcal{L} , then $\kappa = \text{Ref}(\mathcal{C}, \mathcal{P})$ is the strong form of Downward Löwenheim-Skolem Theorem down to $< \kappa$ for \mathcal{L} .

Of these reflection statements, the cases $\aleph_2 = \text{Ref}(\mathcal{C}, \mathcal{P})$ and $2^{\aleph_0} = \text{Ref}(\mathcal{C}, \mathcal{P})$ seems to be of special interest. The former may be interpreted as a pronouncement that the first uncountable cardinal \aleph_1 captures the situation $\neg\mathcal{P}$ good enough while the latter as the pronouncement that the continuum is large enough in connection with the property \mathcal{P} .

The stronger assertions among $\aleph_2 = \text{Ref}(\mathcal{C}, \mathcal{P})$ imply the Continuum Hypothesis while assertions of the form $2^{\aleph_0} = \text{Ref}(\mathcal{C}, \mathcal{P})$ tend to imply that the continuum is extremely large.

Most of the natural assertions of the form $\aleph_2 = \text{Ref}(\mathcal{C}, \mathcal{P})$ or $2^{\aleph_0} = \text{Ref}(\mathcal{C}, \mathcal{P})$ involves some kind of countability in the property \mathcal{P} . This is the case with the reflection assertion $\aleph_2 = \text{Ref}(\mathcal{C}, \mathcal{P})$ where \mathcal{C} is the class of all graphs with induced subgraphs as the notion of substructure and \mathcal{P} is the property “of countable coloring number”. It is shown that this assertion is equivalent to the Fodor-type Reflection Principle (FRP). We can also consider the reflection number for the property obtained from these properties of countable character by replacing the countability by of cardinality κ . Recently many interesting results about reflection statements in this vein are obtained.

In this talk we will give a survey on these reflection statements.

3月19日(月) 第V会場

9:00~11:30

- 20 大 藪 卓 b 数・数体系, 他 5 件 5
 Takashi Oyabu Number; number system, and other 5 talks

概要 (1) Axiomatic set theory: Construction of number system: Some problems are presented: Construction of mathematics::: From ZFC axioms:
 (2) Entropy concept is introduced::: Theory of H-theorems: $H=dAUlogU$:
 (3) Poincaré conjecture is discussed: $AUT(R)$: —Mathematics:
 (4) Our theories are in a sense: Analysis on manifold: i.e. global analysis:
 (5) Some mathematical principles are presented: They are in a sense decisive:
 (6) Phase transition: symmetry breaking: Some schemes are proposed:

- 21 伊 東 由 文 (徳 島 大*) 集合の概念の定義とその存在定理 15
 Yoshifumi Ito (Tokushima Univ.*) Definition of the concept of sets and its existence theorem

概要 In this paper, we give the definition of sets and prove its existence theorem by using the axiomatic method. The system of axioms I used here is the system of axioms ZFC and the axiom of ordinary numbers and the axiom of transfinite induction.

- 22 田 中 義 人 (九州産大経済) A predicate extension of the logic of provability 15
 Yoshihito Tanaka A predicate extension of the logic of provability
 (Kyushu Sangyo Univ.)

概要 In this talk, we introduce a Kripke complete predicate extension of the logic of provability, that is, the propositional modal logic defined by the Löb formula.
 The proof system for the logic of the talk is a modal extension of Gentzen-style sequent calculus for predicate logic. It has a standard derivation rule for necessitation, but does not include the Löb formula as an axiom schema. Instead, it has a non-compact inference rule. We show the logic is complete with respect to the class of Kripke frames of bounded length.

- 23 鈴 木 信 行 (静岡大理) 存在特性を持ち選言特性を持たない中間述語論理をたくさん作る 15
 Nobu-Yuki Suzuki (Shizuoka Univ.) Constructing uncountably many intermediate predicate logics having
 disjunction property but lacking existence property

概要 The disjunction and existence properties in intermediate predicate logics were revealed to be independent in our previous paper. In that paper, we constructed a continuum of intermediate predicate logics having existence property but lacking disjunction property. In this talk, we report the existence of a continuum of intermediate predicate logics having disjunction property but lacking existence property.

- 24 藤 田 憲 悦 (群馬大工) The Church–Rosser Theorem and quantitative analysis of witnesses . . . 15
 Ken-etsu Fujita (Gunma Univ.) The Church–Rosser Theorem and quantitative analysis of witnesses

概要 We analyze quantitative properties of witnesses of the Church–Rosser Theorem for beta-equality in terms of Takahashi translation and by using the notion of parallel reduction. We show that the proof method developed here can be applied to other reduction systems such as lambda-calculus with beta-eta-reduction, Girard’s system F, and Gödel’s system T as well.

- 25 関 隆 宏 (新潟大経営戦略本部) 縮減フレームで特徴づけられる適切様相論理 15
Takahiro Seki (Niigata Univ.) Some relevant modal logics characterized by reduced frames

概要 In Routley–Meyer semantics, relevant logics or relevant modal logics are characterized by unreduced frames in general. In this talk, we consider modal extensions of Slaney’s reduced modeling theorem for relevant logics without WI.

- 26 倉 橋 太 志 (木更津工高専) 理論の分解と証明可能性論理 15
Taishi Kurahashi Provability logics and decompositions of theories
(Nat. Inst. of Tech., Kisarazu Coll.)

概要 We proved that if L is one of the modal logics GL_α , D_β , S_β and GL_β^- where $\alpha \subseteq \omega$ is Σ_1 and $\beta \subseteq \omega$ is cofinite, then for any Σ_1 -definable consistent extension U of Peano Arithmetic PA, there exists a Σ_1 definition $\tau(v)$ of some extension of $I\Sigma_1$ such that the provability logic $PL_\tau(U)$ of $\tau(v)$ relative to U is exactly L . We proved this theorem by using Jeroslow’s method of decomposing theories.

- 27 板 井 昌 典 (東海大理) A model theoretic Rieffel’s theorem of quantum 2-tori 15
Masanori Itai (Tokai Univ.) A model theoretic Rieffel’s theorem of quantum 2-tori

概要 We associate quantum 2-tori T_θ with the structure over $\mathbb{C}_\theta = (\mathbb{C}, +, \cdot, y = x^\theta)$, where $\theta \in \mathbb{R} \setminus \mathbb{Q}$, and introduce the notion of geometric isomorphisms between such quantum 2-tori.

We show that this notion is closely connected with the fundamental notion of Morita equivalence of non-commutative geometry. Namely, we prove a model theoretic version of Rieffel’s theorem of quantum 2-tori.

- 28 岡 部 峻 典 (神戸大システム情報) ジェネリック構造上の自己同型群の幅について 15
Shunsuke Okabe (Kobe Univ.) On widths of automorphism groups on generic structures

概要 The automorphism group on the random graph is simple. Moreover, for all non identical element g of the automorphism group, every element can be denoted by a product of at most three conjugates of g or g^{-1} , by Truss in 2003. We call the number of the product *the width of the automorphism group*. It’s known that the similar fact holds in generic structures analogues of the random graph. That is, if the generic structure has FAP (Free Amalgamation Property) and its automorphism group is transitive, then the group is simple and the width is less than or equal to 32. In this talk, we proof we can improve the width to 12.

- 29 桔 梗 宏 孝 (神戸大システム情報) Hrushovski の ab initio 融合クラスについて 15
岡 部 峻 典 (神戸大システム情報)
Hirohata Kikyo (Kobe Univ.) On Hrushovski’s ab initio amalgamation class
Shunsuke Okabe (Kobe Univ.)

概要 Hrushovski defined an amalgamation class K_f by defining a concave increasing function f referring to an irrational number α with $1/2 < \alpha < 2/3$. His construction works for any real number α with $0 < \alpha < 1$. We obtained the following: (1) f is concave and strictly increasing. (2) K_f is an amalgamation class for any α . (3) f is unbounded if α is rational. (4) If $\alpha = m/d$ then $f(2x) \leq f(x) + 1/d$. Therefore, the generic model of K_f has a model complete theory. (5) There is α (e.g. $\alpha = 1/\sqrt{2}$) such that f is bounded. (6) If f is bounded then the generic model of K_f has no model complete theory. (7) We can give an α in a form of continued fractions where f is unbounded.

11:30~12:00 数学基礎論および歴史分科会総会

13:15~14:15 特別講演

松崎 拓也 (名古屋大 工) 計算機が大学入試数学問題を解く

Takuya Matsuzaki (Nagoya Univ.) A computer program that solves pre-university mathematical problems

概要 In a joint work with many people, we have developed a computer system that solves pre-university level math problems written in natural language. The system is comprised of two parts. One is a language processing pipeline, which translates a math problem into a logical formula. The other is a computer algebra system that derives an answer from the translated problem. In the talk, I will mainly talk about the former part. The main obstacle in the translation from a natural language into a logical language is the flexibility of the natural language, which enables us to convey complex meaning in a concise expression but makes the sentences highly ambiguous for a machine. I will explain how we combat with it using both logical and statistical means.

代 数 学

3月18日(日) 第Ⅲ会場

9:00~11:45

- 1 野村 泰敏 * アペリ型数列の半1次合同式 10
Yasutoshi Nomura Quasi-linear congruences of Apery-like numbers

概要 Based upon machine-experiment we assert that, if Apery-like numbers $X(p-r)$ with prime p is not ultimately constant then $X(p-r)$ is congruent to $(xp-em)/q \pmod{p}$, where $e=1$ or -1 and m is divisible by prime u with $X(u-r)=0 \pmod{p}$, and that, in case $X(p-r)$ has the ultimate constant “Aperi quotient” $W(p)$ satisfies similar congruence.

- 2 鈴木 雄太 (名大多元数理) 互いに素な友愛数について 10
Yuta Suzuki (Nagoya Univ.) On relatively prime amicable pairs

概要 An *amicable pair* is a pair of distinct positive integers each of which is the sum of the proper divisors of the other. Gmelin (1917) conjectured that there is no relatively prime amicable pairs. Artjuhov (1975) and Borho (1974) proved that for any fixed positive integer K , there are only finitely many relatively prime amicable pairs (M, N) with $\omega(MN) = K$, where $\omega(n)$ denotes the number of the distinct prime factors of n . Recently, Pollack (2015) obtained an upper bound $MN < (2K)^{2^{K^2}}$ for such amicable pairs. In this talk, we improve this upper bound to $MN < \frac{\pi^2}{6} 2^{4^K - 2 \cdot 2^K}$.

- 3 飯高 茂 (学習院大*) 究極の完全数と超完全数 10
Shigeru Itaka (Gakushuin Univ.*) Ultimate perfect numbers and hyper perfect numbers

概要 Given an integer m and an odd prime P , if the following equality $(P-1)\sigma(a) = Pa - m$ holds then a is said to be a hyper perfect number with base P and translation parameter m .

- 4 金子 元 (筑波大数理物質) 代数的数のベータ展開における digit 変化数について 10
川島 誠 (阪大理)
Hajime Kaneko (Univ. of Tsukuba) On the number of digit exchanges in the beta expansion of real numbers
Makoto Kawashima (Osaka Univ.)

概要 Many mathematicians have researched the uniformity of the digit expansion of real numbers. In this talk we consider the beta expansion of algebraic numbers, which is a generalization of base- b expansion for a fixed integer $b \geq 2$. For instance, Borel conjectured for each integral base- b that any algebraic irrational number has uniform digits in its base- b expansion. For the study of the uniformity, Bugeaud suggested to consider the number of digit exchanges in the beta expansion of algebraic numbers. In our main result, we considerably improve known results on the asymptotic behavior of the number of digit exchanges.

- 5 渋川 元樹 (阪大情報) Some arithmetic properties of the elliptic Dedekind sums 10
Genki Shibukawa (Osaka Univ.) Some arithmetic properties of the elliptic Dedekind sums

概要 We prove some arithmetic properties of the elliptic Dedekind sums introduced by Egami. Further, we also talk about some conjectures of the elliptic Dedekind sums.

- 6 飛車来人 (徳山工高専) L 関数を任意の精度かつリーマン・ジーゲル公式と同じ速さで計算できる計算法 10

Kurt Fischer (Tokuyama Coll. of Tech.) The Zetafast algorithm to compute L functions to arbitrary precision as fast as the Riemann–Siegel formula

概要 We express the Riemann zeta function $\zeta(s)$ of argument $s = \sigma + i\tau$ with imaginary part τ in terms of three absolutely convergent series. The resulting simple algorithm allows to compute, to arbitrary precision, $\zeta(s)$ and its derivatives using at most $C(\epsilon)|\tau|^{\frac{1}{2}+\epsilon}$ summands for any $\epsilon > 0$, with explicit error bounds. It can be regarded as a quantitative version of the approximate functional equation. The numerical implementation is straightforward. The approach works for any type of zeta function with a similar functional equation such as Dirichlet L -functions.

- 7 スリアジャヤアデイルマ (理化学研) An approximate functional equation for the fourth moment of the Riemann zeta function on the critical line 10

J. Steuding (Univ. of Würzburg)

Ade Irma Suriajaya (RIKEN) An approximate functional equation for the fourth moment of the Riemann zeta function on the critical line

概要 In 2015, Li and Radziwiłł proved an approximate functional equation of the second moment of the Riemann zeta function $\zeta(s)$ on vertical arithmetic progressions on $\text{Re}(s) = 1/2$. Using this formula, they could show that there is an at least $1/3$ proportion of points on arithmetic progressions on $\text{Re}(s) = 1/2$ such that $\zeta(s)$ does not vanish. We are interested in finding the proportion of points on the line such that two consecutive values of $\zeta(s)$ differ. For this purpose, we need an approximate functional equation for the fourth moment of $\zeta(s)$ of Li and Radziwiłł's form. In this talk, we introduce this approximate functional equation we obtained.

- 8 井上翔太 (名大多元数理) Extreme values of Dirichlet L -functions on horizontal line in critical strip 10

Shota Inoue (Nagoya Univ.) Extreme values of Dirichlet L -functions on horizontal line in critical strip

概要 We consider the existence of horizontal lines in critical strip on which a Dirichlet L -function takes its extreme values uniformly. This is an extension of the result for the Riemann zeta-function that was shown by Ramachandra and Sankaranarayanan in 1991. In addition, as an application of this theorem, we also obtain the estimate of the sum of derivatives of Dirichlet L -functions and certain Dedekind zeta-functions over the non-trivial zeros.

- 9 峰正博 (東工大理) The distribution of zeros of the Hurwitz zeta-function on the right side of the critical line 10

Masahiro Mine (Tokyo Tech) The distribution of zeros of the Hurwitz zeta-function on the right side of the critical line

概要 The study of zeros of zeta-functions is a classical topic in analytic number theory. The Riemann hypothesis assert that any nontrivial zeros of the Riemann zeta-function are located on the critical line. On the other hand, the Hurwitz zeta-function has zeros off the critical line in general. Then the upper and lower bounds for the number of such zeros has been considered. In this presentation, we obtain an asymptotic formula on the number of zeros of the Hurwitz zeta-function on the right side of the critical line, applying a certain density function related to the value-distribution of the Hurwitz zeta-function.

- 10 井川 祥彰 (山口大理)* On the number of k -free integers $\leq x$ which are coprime to m 10
 南出 真 (山口大理)
 古屋 淳 (浜松医大)
 谷川 好男

Tadaaki Igawa (Yamaguchi Univ.) On the number of k -free integers $\leq x$ which are coprime to m
 Makoto Minamide (Yamaguchi Univ.)
 Jun Furuya
 (Hamamatsu Univ. School of Medicine)
 Yoshio Tanigawa

概要 Let $k \geq 2$ and $m \geq 1$ be integers. Suryanarayana and Sitaramachandra Rao studied the number of k -free integers $n \leq x$ satisfying $(n, m) = 1$. We shall reconsider error terms in their formula.

- 11 D. Banerjee (IISER)* Bounds of double zeta-function 10
 南出 真 (山口大理)
 谷川 好男

Debika Banerjee (IISER) Bounds of double zeta-function
 Makoto Minamide (Yamaguchi Univ.)
 Yoshio Tanigawa

概要 Let $\zeta_2(s_1, s_2)$ be the double zeta-function of Euler–Zagier type $\zeta_2(s_1, s_2) = \sum_{m=1}^{\infty} \sum_{n=m+1}^{\infty} \frac{1}{m^{s_1} n^{s_2}}$. We shall give several bounds of an error of approximate formula of $\zeta_2(s_1, s_2)$.

- 12 鈴木 正俊 (東工大) ゼータ関数から生ずる積分作用素の族について 10
 Masatoshi Suzuki (Tokyo Tech) On the family of integral operators arising from zeta functions

概要 We consider a family of integral operators arising from zeta functions, and state an equivalence condition of the Riemann hypothesis in terms of operators.

- 13 小山 信也 (東洋大理工) Selberg ゼータ関数の Euler 積の収束性 10
 金子 生弥 (竹園東中)

Shin-ya Koyama (Toyo Univ.) Convergence of Euler products of Selberg zeta functions
 Ikuya Kaneko
 (Tsukuba Takezono Higashi Junior High School)

概要 For any cocompact subgroup of $PSL(2, \mathbb{R})$, and its finite dimensional unitary representation ρ not containing the trivial representation, the Selberg zeta function $Z(s, \rho)$ is defined by the Euler product, which is regular at $s = 1$. It is known that the Euler product is absolutely convergent in $\text{Re}(s) > 1$. In this talk we show convergence of the Euler product for $\text{Re}(s) \geq 3/4$, under assuming the analog of the Riemann hypothesis for the Selberg zeta function.

14:20~17:20

- 14 富安亮子(山形大理) Kaplansky 予想に現れる同じ \mathbb{Z} 上表現を持つ 3 変数 2 次形式の系列について 10

Ryoko Tomiyasu (Yamagata Univ.) On the infinite families of ternary quadratic forms with the same representations over \mathbb{Z} in Kaplansky conjecture

概要 Kaplansky conjectured that if two positive-definite real ternary quadratic forms have perfectly identical representations over \mathbb{Z} , they are equivalent over \mathbb{Z} or constant multiples of regular forms, or is included in either of two families parametrized by \mathbb{Q} . Firstly, the result of an exhaustive search for such pairs of integral quadratic forms is presented, in order to provide a concrete version of the Kaplansky conjecture. The obtained list contains a small number of non-regular forms that were confirmed to have the identical representations up to 3,000,000. Secondly, we prove that if two pairs of ternary quadratic forms have the identical simultaneous representations over \mathbb{Q} , their constant multiples are equivalent over \mathbb{Q} . This was motivated by the question why the other families were not detected in the search.

- 15 富安亮子(山形大理)^b 周期的点集合の平均テータ級数からの決定問題に現れる 2 次形式の問題について 10

Ryoko Tomiyasu (Yamagata Univ.) Problem on quadratic forms that is required to solve for determination of periodic point sets from their average theta series

概要 A problem that has been discussed in crystallography is introduced. When we consider a crystal structure as a periodic point set in R^n , this problem is equivalent to the determination of the periodic points set from its average theta series. We prove that the theta series can almost determine the smallest set that contains the difference set of the periodic point set and is invariant by the action of the automorphism group of the period lattice. This result explains how the problem is reduced to a problem about integral representations of quadratic forms.

- 16 小野雅隆(慶大理工) 2 色根付き木に付随する有限多重ゼータ値 10

Masataka Ono (Keio Univ.) Finite multiple zeta values associated with 2-colored rooted trees

概要 Finite multiple zeta values (FMZVs) were defined first by Kaneko and Zagier and studied by many mathematicians. In this talk, we introduce 2-colored rooted trees, which are some combinatorial objects, and define FMZVs associated with 2-colored rooted trees. We will show that they can be regarded as common generalizations of FMZVs and finite Mordell–Tornheim multiple zeta values defined first by Kamano. Moreover, we will explain that with a mild assumption, FMZVs associated with 2-colored rooted trees can be written as a sum of usual FMZVs. As a corollary, we will give another proof of the shuffle relation among FMZVs, which was first proved by Kaneko and Zagier.

- 17 小野雅隆(慶大理工) 2 色根付き木に付随する多重ゼータ関数 10

Masataka Ono (Keio Univ.) Multiple zeta functions associated with 2-colored rooted trees

概要 In my recent study, I introduced a combinatorial object called 2-colored rooted tree and finite multiple zeta value (FMZV) associated with it to generalize FMZV of Euler–Zagier type and Mordell–Tornheim type simultaneously. In this talk, we introduce multiple zeta function (MZF) associated with 2-colored rooted trees and give some analytical properties. In particular, we give a conjecture on the singularities of MZFs associated 2-colored rooted trees and a new example of MZF satisfying this conjectures.

- 18 町出 智也 多重ゼータスター値の対称和に関する恒等式について 10
 (国立情報学研・JST ERATO)
 Tomoya Machide On an identity involving symmetric sums of regularized multiple zeta-
 (Nat. Inst. of Information/JST ERATO) star values

概要 An identity involving symmetric sums of regularized multiple zeta-star values of harmonic type was proved by Hoffman. In this talk, we prove an identity of shuffle type. In the proof, we meet Bell polynomials appearing in the study of set partitions.

- 19 中筋 麻貴 (上智大理工) Casselman 基底に関する変換係数について 10
 D. Bump (Stanford Univ.)
 Maki Nakasuji (Sophia Univ.) The transition matrix of Casselman basis
 Daniel Bump (Stanford Univ.)

概要 In the spherical principal series representation of a p -adic group, we consider the space of Iwahori-fixed vectors, which has a natural basis and the so-called Casselman basis both indexed by the Weyl group. The latter is defined by using the intertwining integrals. We are interested in the transition matrix of these bases. In order to describe the matrix, we introduced a deformation of the Kazhdan–Lusztig R -polynomials, and proved certain functional equations and a duality formula.

- 20 大井 雅雄 (東大数理) Simple supercuspidal L -packets of quasi-split classical groups 10
 Masao Oi (Univ. of Tokyo) Simple supercuspidal L -packets of quasi-split classical groups

概要 We consider a quasi-split classical group G over a p -adic field F . By the local Langlands correspondence for G , which is recently established by Arthur, we have a natural partition of the set of irreducible smooth representations of $G(F)$ into finite sets which are parametrized by L -parameters. On the other hand, Gross and Reeder defined a some special class of supercuspidal representations which they call the simple supercuspidal representations. In this talk, I will explain a result on an explicit description of the local Langland correspondence for simple supercuspidal representations of quasi-split classical groups.

- 21 小塚 和人 (都城工高専)* p 進 Dedekind–Rademacher 和について 10
 Kazuhito Kozuka p -adic Dedekind–Rademacher sums
 (Miyakonjo Nat. Coll. of Tech.)

概要 In this talk, we consider p -adic functions interpolating Dedekind–Rademacher sums and their reciprocity formula. The results are natural generalizations of the ones due to Rosen and Snyder.

- 22 太田 和惟 (慶大理工) On the rank-part of the Mazur–Tate refined conjecture for modular
 forms 10
 Kazuto Ota (Keio Univ.) On the rank-part of the Mazur–Tate refined conjecture for modular
 forms

概要 The Mazur–Tate refined conjecture connects arithmetic invariants of modular forms with associated L -functions by using Mazur–Tate elements, which are certain elements of group rings of Galois groups and regarded as analogues of Stickelberger elements. In this talk, we discuss its rank-part, which compares the rank of Selmer groups with the order of zeros of Mazur–Tate elements, and our main result is as follows. Under some assumptions, we prove it for higher weight modular forms, generalizing our previous proof for elliptic curves.

- 23 山田一紀 (慶大理工) Comparison of crystalline syntomic and rigid syntomic cohomology for
 V. Ertl (Regensburg Univ.) strictly semistable log schemes 10
Kazuki Yamada (Keio Univ.) Comparison of crystalline syntomic and rigid syntomic cohomology for
 Veronika Ertl (Regensburg Univ.) strictly semistable log schemes

概要 We prove that Nekovář and Nizioł's syntomic cohomology and log rigid syntomic cohomology are isomorphic for a strictly semistable log scheme having a nice compactification and for non-negative twist. Key points of the proof are a generalization of Große-Klönne's log rigid cohomology theory and the compatibility of crystalline and rigid Hyodo-Kato maps on Frobenius eigenspaces.

- 24 植松哲也 (名城大理工) 3次フェルマー曲線の Brauer 群の 3-ねじれ部分について 10
 Tetsuya Uematsu (Meijo Univ.) 3-torsion part of the Brauer group of Fermat curves of degree 3

概要 The Brauer group of varieties has various application to algebraic geometry and number theory. Chernousov and Guletskii studied the 2-torsion part of the Brauer group of elliptic curves, especially its explicit generators represented by norm residue symbols and their relations. Using their method, we study the 3-torsion part of the Brauer group of diagonal cubic curves. In this presentation, we will explain our result in the case of Fermat curves of degree 3.

- 25 國府田玄基 (東京理大理) D_8 を含む同質類に含まれる群をガロア群に持つガロア拡大について ... 10
木田雅成 (東京理大理)
Genki Koda (Tokyo Univ. of Sci.) Certain Galois extensions whose Galois groups are isoclinic to D_8
Masanari Kida (Tokyo Univ. of Sci.)

概要 The notion of isoclinism was introduced by P. Hall in 1940 to classify finite p -groups. We show that this notion also plays an important role to classifying Galois groups of number fields.

- 26 星明考 (新潟大理) Degree three unramified cohomology groups and Noether's problem for
 Ming-chang Kang (Nat. Taiwan Univ.) groups of order 243 10
山崎愛一 (京大理)
Akinari Hoshi (Niigata Univ.) Degree three unramified cohomology groups and Noether's problem for
 Ming-chang Kang (Nat. Taiwan Univ.) groups of order 243
Aiichi Yamasaki (Kyoto Univ.)

概要 Let k be a field, G be a finite group and G act on the rational function field $k(x_g : g \in G)$ by k -automorphisms defined by $h \cdot x_g = x_{hg}$ for any $g, h \in G$. Define $k(G) = k(x_g : g \in G)^G$. Noether's problem asks whether $k(G)$ is rational over k . The unramified cohomology groups $H_{\text{nr}}^i(\mathbb{C}(G), \mathbb{Q}/\mathbb{Z})$ are obstructions to the rationality of $\mathbb{C}(G)$. Theorem 1. Let G be a group of order 3^5 . Then $H_{\text{nr}}^3(\mathbb{C}(G), \mathbb{Q}/\mathbb{Z}) \neq 0$ if and only if G belongs to the isoclinism family Φ_7 . Moreover, if $H_{\text{nr}}^3(\mathbb{C}(G), \mathbb{Q}/\mathbb{Z}) \neq 0$, then $H_{\text{nr}}^3(\mathbb{C}(G), \mathbb{Q}/\mathbb{Z}) \simeq \mathbb{Z}/3\mathbb{Z}$. Theorem 2. Let G be a group of order p^5 where $p = 5$ or $p = 7$. Then $H_{\text{nr}}^3(\mathbb{C}(G), \mathbb{Q}/\mathbb{Z}) \neq 0$ if and only if G belongs to the isoclinism family Φ_6 , Φ_7 or Φ_{10} . Moreover, if $H_{\text{nr}}^3(\mathbb{C}(G), \mathbb{Q}/\mathbb{Z}) \neq 0$, then $H_{\text{nr}}^3(\mathbb{C}(G), \mathbb{Q}/\mathbb{Z}) \simeq \mathbb{Z}/p\mathbb{Z}$. Theorem 3. Let G be a group of order 243. Then $\mathbb{C}(G)$ is \mathbb{C} -rational if and only if G belongs to the isoclinism family Φ_i where $1 \leq i \leq 6$ or $8 \leq i \leq 9$.

- 27 星 明 考 (新潟大理) Computation of degree three unramified cohomology groups using GAP
 Ming-chang Kang (Nat. Taiwan Univ.) 10
 山崎 愛一 (京大理)
 Akinari Hoshi (Niigata Univ.) Computation of degree three unramified cohomology groups using GAP
 Ming-chang Kang (Nat. Taiwan Univ.)
 Aiichi Yamasaki (Kyoto Univ.)

概要 We explain how to compute unramified cohomology group $H_{nr}^3(\mathbb{C}(G), \mathbb{Q}/\mathbb{Z})$ of degree three using Saltman–Peyre method and GAP.

Some algorithms are available from <https://www.math.kyoto-u.ac.jp/~yamasaki/Algorithm/UnramDeg3/>.

3月19日(月) 第Ⅲ会場

9:15~12:00

- 28 丸山文綱^b On a certain method for determining the non-singularity of an integral
 豊泉正男 (東洋大理工) matrix 10
 出口洋三
 Fumitsuna Maruyama On a certain method for determining the non-singularity of an integral
 Masao Toyozumi (Toyo Univ.) matrix
 Yozo Deguchi

概要 It is a joint work with M. Toyozumi and Y. Deguchi. The aim of our study is to give another algorithm to determine the singularity or non-singularity of an integral matrix. Our method can give the result on large order integral matrices within a practical time.

- 29 入江佑樹 (千葉大理)^b p 飽和マヤゲームと対称群の既約表現 10
 Yuki Irie (Chiba Univ.) p -Saturations of Welter’s game and the irreducible representations of
 symmetric groups

概要 We establish a relation between the Sprague–Grundy function of p -saturations of Welter’s game and the degrees of the ordinary irreducible representations of symmetric groups. We present a theorem on these degrees, and using this theorem we obtain an explicit formula for the Sprague–Grundy function of p -saturations of Welter’s game.

- 30 小田文仁 (近畿大理工) Axiomatic theory of Burnside rings I 10
 竹ヶ原裕元 (室蘭工大工)
 吉田知行 (北星学園大経済)
 Fumihito Oda (Kindai Univ.) Axiomatic theory of Burnside rings I
 Yugen Takegahara
 (Muroran Inst. of Tech.)
 Tomoyuki Yoshida
 (Hokusei Gakuen Univ.)

概要 We propose a construction of the Burnside ring of an essentially finite category admitting an epi-mono factorization and enough coequalizers. The main result of our talk is a vast generalization of the embedding theorem of the classical Burnside ring in its ghost ring, with finite cokernel of obstructions. This unifies many constructions and results relative to similar rings, such as various generalizations of the classical Burnside ring (monomial Burnside ring, section Burnside ring, crossed Burnside ring, slice Burnside ringetc), but also some of a seemingly different nature (such as the Möbius algebra of a poset).

- 31 若竹昌洋 (近畿大総理工) The unit group of a partial Burnside ring of a reducible Coxeter group
 小田文仁 (近畿大理工) of type A 10
 Masahiro Wakatake (Kindai Univ.) The unit group of a partial Burnside ring of a reducible Coxeter group
 Fumihito Oda (Kindai Univ.) of type A

概要 In this talk, I will discuss the structure of the unit group of the partial Burnside ring relative to the set of parabolic subgroups of a finite reducible Coxeter group of type A.

- 32 青影一哉 (有明工高専) 対称群のスピン表現におけるテンソル積 10
 Kazuya Aokage Tensor product of the spin representations for the symmetric groups
 (Ariake Nat. Coll. of Tech.)

概要 We describe the multiplicity of the irreducible components of tensor products in even numbers for spin representations.

- 33 上岡修平 (京大情報) 対称平面分割の母関数の精密化 10
 Shuhei Kamioka (Kyoto Univ.) A refinement of generating functions for symmetric plane partitions

概要 For symmetric plane partitions we have generating functions which can be nicely factored, such as the size generating function, half-the-size generating function and Gansner–Nakada’s generating function that respects the diagonal sums. In this talk we show two conjectural formulas which refine the above nice generating functions. Pfaffian expressions for those formulas are also given.

- 34 元良直輝 (京大数理研) \mathcal{W} 代数の余積構造 10
 Naoki Genra (Kyoto Univ.) Coproducts for \mathcal{W} -algebras

概要 The \mathcal{W} -algebras are vertex algebras defined by the generalized Drinfeld–Sokolov reductions. Using the Wakimoto representations of affine Lie algebras, we describe the explicit formulae of the screening operators for the \mathcal{W} -algebras with generic level. As applications, we show that the \mathcal{W} -algebras of type A have the “coproduct” structures related to affine Yangians.

- 35 藤田遼 (京大理) アフィン最高ウェイト圏における傾加群と荒川–鈴木関手 10
 Ryo Fujita (Kyoto Univ.) Tilting modules in affine highest weight categories and the Arakawa–Suzuki functor

概要 The notion of affine highest weight category introduced by Kleshchev generalizes the notion of highest weight category and axiomatizes certain homological structures of some non-semisimple abelian categories of Lie theoretic origin. In this talk, we see the existence of a special kind of tilting module in an affine highest weight category with a large categorical center. As an application, we can prove that a block of the BGG category of $\mathfrak{gl}_m(\mathbb{C})$ is embedded fully faithfully into the module category of finite-dimensional modules over the degenerate affine Hecke algebra of GL_n by the Arakawa–Suzuki functor.

- 36 藤田 遼 (京大 理) アフィン最高ウェイト圏と Dynkin 籠型量子アフィン Schur–Weyl 双対性

Ryo Fujita (Kyoto Univ.) Affine highest weight categories and quantum affine Schur–Weyl duality of Dynkin quiver types

概要 For a Dynkin quiver Q (of type ADE), Hernandez–Leclerc defined a good monoidal subcategory \mathcal{C}_Q inside the category of finite-dimensional modules over the quantum loop algebra $U_q(L\mathfrak{g})$ based on its relationship with the Auslander–Reiten quiver of Q . By using the geometric construction of $U_q(L\mathfrak{g})$ -modules with quiver varieties due to Nakajima, we see that a completion of the category \mathcal{C}_Q has a structure of affine highest weight category. As an application, we can prove that Kang–Kashiwara–Kim’s generalized quantum affine Schur–Weyl duality functor gives an equivalence of monoidal categories between the category of finite-dimensional modules over the quiver Hecke (KLR) algebra associated to Q and Hernandez–Leclerc’s category \mathcal{C}_Q .

- 37 櫻井 太朗 (千葉大 理) On characterizations of small block algebras 10

越谷 重夫
 (千葉大先進科学センター・千葉大*)

Taro Sakurai (Chiba Univ.) On characterizations of small block algebras
 Shigeo Koshitani
 (Chiba Univ./Chiba Univ.*)

概要 In 1941, Brauer–Nesbitt established a characterization of a block with trivial defect group as a block B with $k(B) = 1$. In 1982, Brandt established a characterization of a block with defect group of order two as a block B with $k(B) = 2$. These correspond to the cases when the block is Morita equivalent to the one-dimensional algebra and to the non-semisimple two-dimensional algebra respectively. In this talk, we redefine $k(A)$ to be the codimension of the commutator subspace $K(A)$ of a finite-dimensional algebra A and show analogous statements for arbitrary finite-dimensional algebras.

- 38 佐々木 洋城 (信州大 教育)* デフェクト群が extraspecial p -群である ブロック・イデアルのコホモロ
 ジー環 10

Hiroki Sasaki (Shinshu Univ.) Cohomology rings of block ideals with extraspecial defect groups

概要 Let b be a block ideal of the finite group algebra kG with a defect group S which is extraspecial of order p^3 and of exponent p . Let i be a source idempotent. We shall examine the module structure of the source algebra $ikGi$ of the block ideal b and show that the image of the transfer map $t : H^*(S, k) \rightarrow H^*(S, k)$ induced by $ikGi$ coincides with the cohomology ring of the block b with respect to the source idempotent i .

- 39 越谷 重夫 ^b スコット加群のブラウアー直既約性 10

(千葉大先進科学センター・千葉大*)

ケサールラダー

(City, Univ. of London)

リンケルマンマーカス

(City, Univ. of London)

Shigeo Koshitani

Brauer indecomposabilities of the Scott modules

(Chiba Univ./Chiba Univ.*)

Radha Kessar (City, Univ. of London)

Markus Linckelmann

(City, Univ. of London)

概要 We shall be talking about Brauer indecomposabilities of the Scott modules. The Scott module is an indecomposable p -permutation kG -module which contains the trivial kG -module (where G is a finite group and k is an algebraically closed field of characteristic $p > 0$). This plays an important role to try to prove so-called global-local conjecture in the representation theory of finite groups.

- 40 越谷重夫 ^b 自明な加群の射影被覆についてのある注意 10
 (千葉大先進科学センター・千葉大*)
 ミュラーユルゲン (Wuppertal Univ.)
Shigeo Koshitani A remark on the projective cover of the trivial module
 (Chiba Univ./Chiba Univ.*)
 Jürgen Müller (Wuppertal Univ.)

概要 We shall be talking on the projective cover of the trivial module for the group algebra kG where G is a finite group and k is a field of characteristic $p > 0$. Especially we will be interested in the case when p is odd and G is a finite simple group of Lie type defined over a finite field of the same characteristic p .

- 41 越谷重夫 ^b 有限群代数における単純加群の Auslander–Reiten 箭における位置 10
 (千葉大先進科学センター・千葉大*)
 ラシュールカロリーヌ
 (Kaiserslautern Univ.)
Shigeo Koshitani Locations of simple modules for finite group algebras in the Auslander–
 (Chiba Univ./Chiba Univ.*) Reiten quivers
 Caroline Lassueur
 (Kaiserslautern Univ.)

概要 We will be talking on the positions of simple modules for finite group algebras in connected components of the stable Auslander–Reiten quivers to which the simple modules belong.

13:15~14:15 特別講演

- 清水健一 非半単純なモジュラーテンソル圏に関する最近の展開
 (芝浦工大システム理工)
 Kenichi Shimizu Recent developments of ‘non-semisimple’ modular tensor categories
 (Shibaura Inst. of Tech.)

概要 A modular tensor category is usually defined as a semisimple ribbon category satisfying a certain non-degeneracy condition. Nevertheless, with motivation coming from CFT and TQFT, it is important and interesting to drop the semisimplicity assumption from the definition of a modular tensor category. Lyubashenko has formulated such a ‘non-semisimple’ modular tensor category and showed that, as in the semisimple case, a ‘non-semisimple’ modular tensor category yields an invariant of closed 3-manifolds and a projective representation of the surface mapping class groups. In this talk, while introducing several category-theoretical techniques that are important in the recent study of tensor categories, I will review recent developments of ‘non-semisimple’ modular tensor categories. I will, especially, mention the recent result that a ribbon finite tensor category is modular if and only if its Müger center is trivial. This criterion yields several new examples of modular tensor categories and factorizable (quasi-)Hopf algebras.

3月20日(火) 第三会場

9:15~12:00

- 42 松岡拓男 (Takuo Matsuoka) Filtration of a stable infinity 1-category 10
 Takuo Matsuoka Filtration of a stable infinity 1-category

概要 **Stability** is the appropriate notion of linearity in *homotopical algebra*, which supersedes *Abelianness* in classical algebra. Thus, stable homotopical algebra is useful e.g., for refining the method of *derived categories*. While homotopical algebra can conveniently be formulated in a $(\infty, 1)$ -category, to be simply called “**category**” here, stable homotopical algebra can be understood as algebra in a “stable” such. We discuss the theory of *filtration* of a stable category, which is useful for controlling behaviour of limits in stable homotopical algebra, and can be applied e.g., for the study of the *Koszul duality* and *higher Morita categories*. Examples of a filtration include a *t-structure*, and a natural filtration on the category of *filtered objects* in a stable category. Both may come with a compatible symmetric monoidal structure in practice.

- 43 毛利出 (静岡大理) 非可換射影空間の圏論的特徴付け 10
 上山健太 (弘前大教育) Izuru Mori (Shizuoka Univ.) A categorical characterization of noncommutative projective spaces
 Kenta Ueyama (Hirosaki Univ.)

概要 The noncommutative projective scheme associated to an AS-regular algebra is considered as a noncommutative projective space, and has been studied deeply and extensively in noncommutative algebraic geometry. In this talk, we will characterize a k -linear abelian category \mathcal{C} such that \mathcal{C} is equivalent to the noncommutative projective scheme associated to some AS-regular algebra.

- 44 板場綾子 (東京理大理) 楕円曲線に対応する3次元 quadratic AS 正則環 10
 松野仁樹 (静岡大理) Ayako Itaba (Tokyo Univ. of Sci.) 3-dimensional quadratic AS-regular algebras corresponding to elliptic
 Masaki Matsuno (Shizuoka Univ.) curves

概要 In this talk, we determine the automorphism group of an elliptic curve E in \mathbb{P}^2 depending on its j -invariant $j(E)$. By using the automorphism σ of E , we calculate the defining relations of a 3-dimensional quadratic AS-regular algebra corresponding to the pairs (E, σ) . By this calculations, we find a counterexample to the conjecture that any 3-dimensional quadratic AS-regular algebra Λ corresponding to an elliptic curve is isomorphic to a twist A^φ of a Sklyanin algebra A by $\varphi \in \text{Aut } A$.

- 45 板場綾子 (東京理大理) 道多元環上のある無限生成非射影加群について 10
 D. A. Mejía (静岡大理) Ayako Itaba (Tokyo Univ. of Sci.) On some non-projective infinitely generated modules over path algebras
 依岡輝幸 (静岡大理) Diego Alejandro Mejía (Shizuoka Univ.)
 Teruyuki Yorioka (Shizuoka Univ.)

概要 It is known that for any finite dimensional algebra Λ of finite global dimension and any finitely generated Λ -module M , if $\text{Ext}_\Lambda^{\geq 1}(M, \Lambda) = 0$, then M is projective. Let Q be the following quiver:

$$\circ \rightarrow \circ \rightarrow \cdots \circ \rightarrow \circ \rightarrow$$

and K an algebraically closed field. Using the compactness theorem of Mathematical Logic, we prove that for any finitely dimensional K -representation \mathcal{K} , if $\text{Ext}_{KQ}^1(\mathcal{K}, KQ) = 0$, then \mathcal{K} is projective. It is also proved that, under Martin’s Axiom (which is a combinatorial statement consistent with Axiomatic Set Theory), there exists a non-projective KQ -module M such that $\text{Ext}_{KQ}^1(\mathcal{K}, KQ) = 0$.

- 46 鯉江秀行 (東京理大理) On presentations of Hochschild extension algebras for a class of self-injective Nakayama algebras 10
 板垣智洋 (東京理大理)
 眞田克典 (東京理大理)
 Hideyuki Koie (Tokyo Univ. of Sci.) On presentations of Hochschild extension algebras for a class of self-injective Nakayama algebras
 Tomohiro Itagaki (Tokyo Univ. of Sci.)
 Katsunori Sanada (Tokyo Univ. of Sci.)

概要 For a bound quiver algebra satisfying the condition that the every oriented cycles in the quiver are vanished in the algebra, Fernádez and Platzeck determined the bound quiver algebra which is isomorphic to the trivial extension algebra. In this paper, we consider a Hochschild extension algebra which is a generalization of a trivial extension algebra. Our aim is to determine the bound quiver algebras which are isomorphic to Hochschild extension algebras of some finite dimensional self-injective Nakayama algebras.

- 47 板垣智洋 (東京理大理) Batalin–Vilkovisky algebra structures on the Hochschild cohomology of self-injective Nakayama algebras 10
 Tomohiro Itagaki (Tokyo Univ. of Sci.) Batalin–Vilkovisky algebra structures on the Hochschild cohomology of self-injective Nakayama algebras

概要 In this talk, we determine the Batalin–Vilkovisky algebra structure on the Hochschild cohomology of self-injective Nakayama algebras over an algebraically closed field.

- 48 浅井聡太 (名大多元数理)* 前射影的多元環のブリック 10
 Sota Asai (Nagoya Univ.) Bricks over preprojective algebras

概要 Mizuno gave an isomorphism of lattices from a Coxeter group of Dynkin type to the set of torsion-free classes in the module category of the corresponding preprojective algebra. Combining it with my bijection on semibricks, we obtain a bijection from the Coxeter group to the set of semibricks over the preprojective algebra. My aim is to explicitly describe the semibrick associated to each element in the Coxeter group in this bijection. In this process, a combinatorial notion “canonical join representations” introduced by Reading, is very useful. I observed that the canonical join representation of an element in the Coxeter group gives the decomposition of the corresponding semibrick into bricks. I will talk about such theoretic strategies to determine the semibrick.

- 49 中村力 (岡山大自然) Bousfield localization and cosupport in derived categories of commutative Noetherian rings 10
 Tsutomu Nakamura (Okayama Univ.) Bousfield localization and cosupport in derived categories of commutative Noetherian rings

概要 This talk is based on joint work with Yuji Yoshino. Let R be a commutative Noetherian ring. We denote by \mathcal{D} the unbounded derived category of R . An exact functor $\lambda : \mathcal{D} \rightarrow \mathcal{D}$ is called a localization functor if there is a morphism $\eta : \text{id}_{\mathcal{D}} \rightarrow \lambda$ such that $\lambda\eta$ is invertible and $\lambda\eta = \eta\lambda$. This notion was introduced by A. K. Bousfield in his topological work (1979). In this talk, we give a concrete way to compute localization functors on \mathcal{D} by using the notions of cosupport and Čech complexes. As an application, we can obtain a functorial way to construct pure-injective resolutions for complexes of flat R -modules and complexes of finitely generated R -modules.

- 50 松井 紘樹 (名大多元数理) Singular equivalences and reconstruction of singular loci 10
 Hiroki Matsui (Nagoya Univ.) Singular equivalences and reconstruction of singular loci

概要 The singularity category $D_{sg}(R)$ of a commutative Noetherian ring R is a triangulated category which measures singularity of R . Two commutative Noetherian rings R and S are said to be singularly equivalent if their singularity categories are equivalent as triangulated categories. Singular equivalence have deeply been studied in non-commutative setting and various examples are known, while in commutative setting, only a few examples of singular equivalence are known. The aim of this talk is to give a necessary condition for singular equivalence by using singular loci. The key tool to prove our main result is the support theory for triangulated categories without tensor structure.

- 51 小林 稔周 (名大多元数理) Syzygies of Cohen–Macaulay modules and endomorphism ring of the maximal ideal 10
 Toshinori Kobayashi (Nagoya Univ.) Syzygies of Cohen–Macaulay modules and endomorphism ring of the maximal ideal

概要 I will talk on syzygies of (maximal) Cohen–Macaulay modules over one dimensional Cohen–Macaulay local rings. We compare these modules to Cohen–Macaulay modules over the endomorphism ring of the maximal ideal. After this comparison, we give several characterizations of almost Gorenstein rings in terms of syzygies of Cohen–Macaulay modules.

- 52 早坂 太 (岡山大環境)* 多重次数付き加群の節減と単項式イデアルの整閉性 10
 Futoshi Hayasaka (Okayama Univ.) Complete reductions of multigraded modules and normality of monomial ideals

概要 Reduction of ideals introduced by Northcott and Rees plays an important role in the study of local rings, especially the multiplicity theory of ideals. This notion is extended to the multigraded modules and used in the study of asymptotic properties of them. In this talk, I will give a result about existence of certain complete reductions of multigraded modules. By applying the result to multi-Rees algebras of finitely many ideals, we obtain a result on normality of monomial ideals, which extends and improves several known results on this topic.

- 53 神代 真也 (千葉大理) イデアル化 $R \times I$ の almost Gorenstein 性 10
 後藤 四郎 (明大*)
 Shinya Kumashiro (Chiba Univ.) When is $R \times I$ an almost Gorenstein local ring?
 Shiro Goto (Meiji Univ.*)

概要 Let R be a Noetherian local ring and M be a nonzero finitely generated R -module. The notion of almost Gorenstein rings is one of the generalization of the notion of Gorenstein rings. The purpose of this talk is to explore the question of when the idealization $R \times M$ of M is an almost Gorenstein local rings. Although this problem was investigated by S. Goto, R. Takahashi, and N. Taniguchi, it is still open.

- 54 磯部 遼太郎 (千葉大 理) Characterization of generalized Gorenstein rings 10
 後藤 四郎 (明大*)
 谷口 直樹 (早大 GEC)
 神代 真也 (千葉大 理)
 Ryotaro Isobe (Chiba Univ.) Characterization of generalized Gorenstein rings
 Shiro Goto (Meiji Univ.*)
 Naoki Taniguchi (Waseda Univ.)
 Shinya Kumashiro (Chiba Univ.)

概要 The notion of a generalized Gorenstein local ring (GGL ring for short) is one of the generalizations of Gorenstein rings. Similarly for almost Gorenstein local rings, the notion is given in terms of a certain specific embedding of the rings into their canonical modules. In this talk, we give a characterization of GGL rings in terms of their canonical ideals and related invariants.

14:15~14:30 2018年度(第21回)日本数学会代数学賞授与式

14:30~15:30 2018年度(第21回)日本数学会代数学賞受賞特別講演

佐藤 周友 (中大 理工) 数論的スキームに対する新しいコホモロジー理論とその応用

Kanetomo Sato (Chuo Univ.) A new cohomology theory for arithmetic schemes and its applications

概要 In this talk, I would like to explain an approach to motivic cohomology in the étale topology with $\mathbb{Z}/p^n\mathbb{Z}$ -coefficients of an arithmetic scheme X which has good or semistable reduction at all primes dividing p . We construct a complex of étale sheaves $\mathbb{Z}/p^n\mathbb{Z}(r)$ on X for $r \geq 0$ by gluing the r -fold tensor power of the locally constant sheaf μ_{p^n} on $X[p^{-1}]$ with a certain differential sheaf on the fiber over p via the boundary map of Galois cohomology groups due to K. Kato. In the good reduction case, the object $\mathbb{Z}/p^n\mathbb{Z}(r)$ is already considered by J. S. Milne and P. Schneider about 30 years ago. What I did on this object is that I defined it in the semistable reduction case and proved a global duality result for étale cohomology with coefficients in $\mathbb{Z}/p^n\mathbb{Z}(r)$, which had not been unknown even in the good reduction case. I will also talk about a few applications of $\mathbb{Z}/p^n\mathbb{Z}(r)$ to the study of algebraic cycles on arithmetic schemes.

15:40~16:40 2018年度(第21回)日本数学会代数学賞受賞特別講演

内藤 聡 (東工大 理工) 量子アフィン代数の表現論

Satoshi Naito (Tokyo Tech) Representation theory of quantum affine algebras

概要 In this talk, I will explain the following aspects of level-zero representations of quantum affine algebras:
 1) explicit combinatorial realization, by semi-infinite Lakshmibai–Seshadri paths, of crystal bases of Demazure submodules of level-zero extremal weight modules over quantum affine algebras;
 2) explicit relation of graded characters of level-zero Demazure submodules with the specializations of nonsymmetric Macdonald polynomials at $t = 0$ and $t = \infty$;
 3) algebro-geometric interpretation of graded characters of level-zero Demazure submodules via Borel–Weil–Bott type theorem for semi-infinite flag manifolds.

16:50~17:50 2018年度(第21回)日本数学会代数学賞受賞特別講演

日比 孝之 (阪大 情報) 単項式イデアルと二項式イデアル

Takayuki Hibi (Osaka Univ.) Monomial ideals and binomial ideals

概要 Creating new trends of mutual development with combinatorics on convex polytopes, simplicial complexes, finite partially ordered sets and finite graphs and with statistics on contingency tables and experimental designs, commutative algebra on monomial ideals and binomial ideals is rapidly and dramatically growing by making the best use of modern techniques on, for example, Gröbner bases. In addition, plenty of fascinating problems remain unsolved. In the present draft, current streams of monomial ideals and binomial ideals will be surveyed quickly and their prospects will be predicted.

3月21日(水) 第三会場

9:15~12:00

- 55 東平光生(明大研究・知財) Sequentially generalized Cohen–Macaulay となる二部グラフについて · 10
Hiroataka Higashidaira (Meiji Univ.) On sequentially generalized Cohen–Macaulay bipartite graphs

概要 Let S be the polynomial ring in n variables over a field K and H a bipartite graph with n vertices. We denote by $I(H)$ the edge ideal of H . In 2003, N. T. Cuong and L. T. Nhan introduced the notion of sequentially generalized Cohen–Macaulay. H is called sequentially generalized Cohen–Macaulay when so is $S/I(H)$.

In this talk, we consider properties of sequentially generalized Cohen–Macaulay bipartite graphs. In particular, we investigate the behavior of edges of H when $\text{depth}(S/I(H)) \leq 3$. Consequently, we give a characterization of sequentially generalized Cohen–Macaulay graphs of essential dimension 3.

- 56 土谷昭善(阪大情報) パーフェクトグラフに付随する Gorenstein Fano 凸多面体 ····· 10
日比孝之(阪大情報)
Akiyoshi Tsuchiya (Osaka Univ.) Gorenstein Fano polytopes arising from perfect graphs
Takayuki Hibi (Osaka Univ.)

概要 Gorenstein Fano polytopes form one of the distinguished classes of lattice polytopes. Especially normal Gorenstein Fano polytopes are of interest. In this talk, we will give a new class of normal Gorenstein Fano polytopes arising from perfect graphs.

- 57 土谷昭善(阪大情報) 格子凸多面体のケーリー和の正規性と level 性 ····· 10
Akiyoshi Tsuchiya (Osaka Univ.) Normality and levelness of Cayley sums of lattice polytopes

概要 Normality or the integer decomposition property (IDP) is one of the most important properties on lattice polytopes. In fact, many authors have been studied the properties from view-points of combinatorics, commutative algebra and algebraic geometry. In this talk, we discuss when a Cayley sum is normal. Moreover, we consider when a Cayley sum is level.

- 58 松田一徳(阪大情報) 単項式イデアルの Castelnuovo–Mumford 正則度と h 多項式の次数につ
日比孝之(阪大情報) いて ····· 10
Kazunori Matsuda (Osaka Univ.) Regularity and h -polynomials of monomial ideals
Takayuki Hibi (Osaka Univ.)

概要 Let $S = K[x_1, \dots, x_n]$ denote the polynomial ring in n variables over a field K with each $\deg x_i = 1$ and $I \subset S$ a homogeneous ideal of S with $\dim S/I = d$. The Hilbert series of S/I is of the form $h_{S/I}(\lambda)/(1-\lambda)^d$, where $h_{S/I}(\lambda) = h_0 + h_1\lambda + h_2\lambda^2 + \dots + h_s\lambda^s$ with $h_s \neq 0$ is the h -polynomial of S/I . It is known that, when S/I is Cohen–Macaulay, one has $\text{reg}(S/I) = \deg h_{S/I}(\lambda)$, where $\text{reg}(S/I)$ is the (Castelnuovo–Mumford) regularity of S/I . In this talk, given arbitrary integers r and s with $r \geq 1$ and $s \geq 1$, a monomial ideal I of $S = K[x_1, \dots, x_n]$ with $n \gg 0$ for which $\text{reg}(S/I) = r$ and $\deg h_{S/I}(\lambda) = s$ will be constructed.

- 59 東谷章弘(京都産大理) 多項式環の Segre 積の generalized F-signature について ····· 10
中嶋祐介(東大IPMU)
Akihiro Higashitani On generalized F-signatures of Segre product of polynomial rings
(Kyoto Sangyo Univ.)
Yusuke Nakajima (Univ. of Tokyo)

概要 Let K be an algebraically closed field of positive characteristic. We set $S(t) = K[x_1, y_1] \# \dots \# K[x_t, y_t]$ and $R(r, s) = K[x_1, \dots, x_{r+1}] \# K[y_1, \dots, y_{s+1}]$. In this talk, we will compute generalized F-signatures of all modules belonging to the FFRT system of each of $S(t)$ and $R(r, s)$.

- 60 日比孝之(阪大情報) 膨らませた空単体の正則単模三角形分割の存在 10
 東谷章弘(京都産大理) 膨らませた空単体の正則単模三角形分割の存在 10
 吉田恒太郎(阪大情報) 膨らませた空単体の正則単模三角形分割の存在 10
 Takayuki Hibi (Osaka Univ.) Existence of regular unimodular triangulations of dilated empty simplices
 Akihiro Higashitani (Kyoto Sangyo Univ.) Existence of regular unimodular triangulations of dilated empty simplices
 Koutarou Yoshida (Osaka Univ.) Existence of regular unimodular triangulations of dilated empty simplices

概要 Given integers k and m with $k \geq 2$ and $m \geq 2$, let P be an empty simplex of dimension $(2k - 1)$ whose δ -polynomial is of the form $1 + (m - 1)t^k$. In this talk, the necessary and sufficient condition for the k -th dilation kP of P to have a regular unimodular triangulation will be presented.

- 61 宮崎充弘(京都教育大) On the anticanonical level property of a Hibi ring 10
 Mitsuhiro Miyazaki On the anticanonical level property of a Hibi ring
 (Kyoto Univ. of Edu.) On the anticanonical level property of a Hibi ring

概要 Let R be a Hibi ring and ω the canonical ideal of R . We denote the n -th power of ω in $D(R)$, the group of divisorial ideals of R , by $\omega^{(n)}$ for any integer n . R is by definition, anticanonical level if all the generators of $\omega^{(-1)}$ have the same degree. In this talk, we analyze the structure of $\omega^{(n)}$ for any integer n and state a criterion of anticanonical level property of R .

- 62 知念宏司(近畿大理工) Divisible formal weight enumerator に対する Mallows–Sloane bound の類似 10
 Koji Chinen (Kindai Univ.) Analogs of Mallows–Sloane bound for divisible formal weight enumerators

概要 The Mallows–Sloane bound is the inequality which estimates the minimum distance by the code length for a divisible self-dual code. Analogous inequalities for formal weight enumerators are completed.

- 63 中島規博(東京電機大情報) 超平面配置の高階自由性に関する Holm の問題の反例 10
 阿部拓郎(九大IMI) 超平面配置の高階自由性に関する Holm の問題の反例 10
 Norihiro Nakashima Counter examples of Holm’s questions for high order free hyperplane arrangements
 (Tokyo Denki Univ.) Counter examples of Holm’s questions for high order free hyperplane arrangements
 Takuro Abe (Kyushu Univ.) Counter examples of Holm’s questions for high order free hyperplane arrangements

概要 An m -free hyperplane arrangement is a generalization of a free arrangement. There are rich researches for free arrangements, but the behavior of m -freeness has not been well analyzed yet when $m > 2$. Some basic questions remain open. In particular, Holm asked the following: (1) Does m -free imply $(m + 1)$ -free for any arrangement? (2) Are all arrangements m -free for m large enough? In this talk, we characterize m -freeness for product arrangements and show that all localizations of an m -free arrangement are m -free. From these results, we give counter examples to Holm’s questions.

- 64 堀口達也 (阪大情報) ヘッセンバーク多様体と超平面配置 10
 阿部拓郎 (九大IM1)
 柘田幹也 (阪市大理)
 村井聡 (阪大情報)
 佐藤敬志 (阪市大数学研)
 Tatsuya Horiguchi (Osaka Univ.) Hessenberg varieties and hyperplane arrangements
 Takuro Abe (Kyushu Univ.)
 Mikiya Masuda (Osaka City Univ.)
 Satoshi Murai (Osaka Univ.)
 Takashi Sato (Osaka City Univ.)

概要 Hessenberg varieties are subvarieties of a flag variety. This subject makes connections with many research areas such as geometric representation theory, quantum cohomology of the flag variety, chromatic quasisymmetric functions of graph theory, and hyperplane arrangements. In this talk, I will explain the connection between Hessenberg varieties and hyperplane arrangements. More concretely, we show that a certain graded ring derived from the logarithmic derivation module of an ideal arrangement is isomorphic to the cohomology ring of a regular nilpotent Hessenberg variety, and the Weyl group invariant subring of the cohomology of a regular semisimple Hessenberg variety. This is joint work with Takuro Abe, Mikiya Masuda, Satoshi Murai, and Takashi Sato.

- 65 久保田絢子 (早大理工) Invariant Hilbert scheme resolution of Popov's $SL(2)$ -varieties 10
 Ayako Kubota (Waseda Univ.) Invariant Hilbert scheme resolution of Popov's $SL(2)$ -varieties

概要 Let X be an affine G -variety, where G is a reductive algebraic group. The invariant Hilbert scheme parametrizes closed G -subschemes of X whose coordinate rings have a prescribed decomposition as G -modules. One of the main usage of the invariant Hilbert scheme, together with so-called the Hilbert–Chow morphism, is to study singularities of affine quotient varieties. In this talk, we study Popov's $SL(2)$ -varieties by means of the invariant Hilbert scheme.

- 66 金鍾明 (名大多元数理) A freeness criterion for spherical twists 10
 Jong Myeong Kim (Nagoya Univ.) A freeness criterion for spherical twists

概要 Spherical twists along spherical objects are autoequivalences of a triangulated category defined by Seidel and Thomas as a categorical analogue of Dehn twists along simple closed curves. Spherical twists share many properties with Dehn twists. On the other hand, there is a classical result by Humphries which states that if a collection of simple closed curves admits a “complete partition” and does not bound a disk then the group generated by the Dehn twists along them is isomorphic to the free product of free abelian groups. In this talk, we give a categorical analogue of Humphries' argument.

- 67 土基善文 (高知大理) Non-commutative Kähler projective space: from commutative view-point 10
 Yoshifumi Tsuchimoto (Kochi Univ.) Non-commutative Kähler projective space: from commutative view-point

概要 We define “non-commutative Kähler projective space”. We describe fully the “infinitesimal deformation case” and discuss “finite (non-perturbative) deformation case”. We then examine the cohomology in the “infinitesimal deformation case” from a view point of commutative algebraic geometry.

14:20~17:00

- 68 安藤 哲哉 (千葉大理) 半代数多様体 10
 Tetsuya Ando (Chiba Univ.) Semialgebraic variety

概要 We give basic concepts of semialgebraic varieties. We will treat a semialgebraic subset of a real algebraic variety as a certain kind of abstract locally ringed space. We study a relation with semialgebraic varieties and complex algebraic varieties.

- 69 野口潤次郎 (東大*)^b Big Picard Theorem and the Manin–Mumford Conjecture 10
 Junjiro Noguchi (Univ. of Tokyo*) Big Picard Theorem and the Manin–Mumford Conjecture

概要 We will prove a theorem of Ax–Lindemann type for complex semi-abelian varieties as an application of a big Picard Theorem proved by the author in 1981, and then apply it to prove a theorem of the classical Manin–Mumford Conjecture for semi-abelian varieties, which was proved by M. Raynaud 1983, M. Hindry 1988, ..., and Pila–Zannier 2008 by a different method from others, which is most relevant to ours.

- 70 工藤 桃成 (九大IMI) Enumerating superspecial curves of genus 4 over prime fields 10
 原下 秀士 (横浜国大環境情報)
 Momonari Kudo (Kyushu Univ.) Enumerating superspecial curves of genus 4 over prime fields
 Shushi Harashita
 (Yokohama Nat. Univ.)

概要 A curve is called superspecial if its Jacobian is isomorphic to a product of supersingular elliptic curves. The purpose of this study is to enumerate superspecial curves of genus g over \mathbb{F}_q for a given g and for a given q . In 2016, the speakers gave an algorithm to enumerate superspecial curves of genus $g = 4$ over \mathbb{F}_q with $q > 5$. By executing the algorithm on a computer algebra system Magma, they also enumerated superspecial curves of genus 4 over \mathbb{F}_{25} and \mathbb{F}_{49} . In this talk, we present an improved algorithm, which works for any finite field \mathbb{F}_q with $q \geq 5$, and classifies the isomorphism classes of superspecial curves of genus 4. By our implementation of the improved algorithm over Magma, we newly enumerate superspecial curves of genus 4 over prime fields \mathbb{F}_p for $p \leq 11$.

- 71 千田 駿人 (横浜国大環境情報) Automorphism groups of superspecial curves of genus 4 over \mathbb{F}_{11} 10
 工藤 桃成 (九大IMI)
 原下 秀士 (横浜国大環境情報)
 Hayato Senda (Yokohama Nat. Univ.) Automorphism groups of superspecial curves of genus 4 over \mathbb{F}_{11}
 Momonari Kudo (Kyushu Univ.)
 Shushi Harashita
 (Yokohama Nat. Univ.)

概要 A nonsingular projective curve C over a field K of positive characteristic is called superspecial if its Jacobian is isomorphic to a product of supersingular elliptic curves. In 2016 and 2017, Kudo and Harashita enumerated nonhyperelliptic superspecial curves of genus 4 over \mathbb{F}_{25} , \mathbb{F}_{49} and \mathbb{F}_{11} . By implementing our algorithm over Magma, we determined the structure of the automorphism group of every nonhyperelliptic superspecial curve of genus 4 over \mathbb{F}_{11} .

- 72 白根竹人 (宇部工高専) 分解グラフと平面曲線の埋込位相 10
 Taketo Shirane Splitting graph and the embedded topology of plane curves
 (Ube Nat. Coll. of Tech.)

概要 We consider the embedded topology of plane curves in the complex projective plane, as in knot and link theory. In this talk, we introduce a new invariant, called the *splitting graph*, to distinguish the embedded topology of plane curves. This invariant is a generalization of the splitting number, which is an invariant not determined by the fundamental group of the complement of a plane curve. By using the splitting graph, we distinguish the embedded topology of plane curves consisting of one smooth curve and three lines.

- 73 坂内真三 (茨城工高専) Rational points of elliptic surfaces and the topology of cubic-line ar-
 徳永浩雄 (首都大東京理工) rangements 10
 山本桃果 (首都大東京理工)
 Shinzo Bannai Rational points of elliptic surfaces and the topology of cubic-line ar-
 (Ibaraki Nat. Coll. of Tech.) rangements
 Hiro-o Tokunaga (Tokyo Metro. Univ.)
 Momoko Yamamoto
 (Tokyo Metro. Univ.)

概要 In this talk we introduce some recent study of rational points of elliptic surfaces done from a geometric point of view, and apply it to construct interesting examples of plane curve arrangements of low degree which give rise to candidates for Zariski pairs. The constructed arrangements can be distinguished topologically by studying the arithmetic properties of the rational points used in the construction. Previously known examples of Zariski pairs consisting of cubic-line arrangements all involved flex tangent lines, but our new examples do not contain any flex tangents.

- 74 奥村克彦 (早大理工) Fano 多様体の直積上の単純正規交差対数的シンプレクティック構造 ... 10
 Katsuhiko Okumura (Waseda Univ.) SNC log symplectic structures on Fano products

概要 In the classification problem of Poisson structures, log symplectic structures which are generically symplectic Poisson structures with reduced degeneracy divisor is one of the most important class. Lima and Pereira studied log symplectic structures with simple normal crossing degeneracy divisor in the case that the variety is a Fano variety of Picard number 1 and they discovered a characterization of projective spaces. My research consider the case that the variety is a product of Fano varieties of Picard number 1. We will extend the result of Lima and Pereira and also give a better characterization of projective spaces.

- 75 南 範彦 (名工大) On covering by rational varieties 10
 Norihiko Minami On covering by rational varieties
 (Nagoya Inst. of Tech.)

概要 Taku Suzuki (arXiv.16060.9350) gave some sufficient condition for a smooth Fano manifold to be covered by rational N -folds. Results of this sort might be useful for future investigation of “higher connectivity” properties in the Morel–Voevodsky Motivic homotopy theory. Motivated by this, I shall report that a slight relaxation of the condition in this Suzuki’s theorem is possible.

- 76 柴田 康介 (東大数理) Bound of the multiplicity of complete intersection singularities 10
 Kohsuke Shibata (Univ. of Tokyo) Bound of the multiplicity of complete intersection singularities

概要 The multiplicity of a point on a variety is a fundamental invariant to estimate how bad the singularity is. It is introduced in a purely algebraic context. On the other hand, we can also attach to the singularity the log canonical threshold, which is introduced in a birational theoretic context. In this talk, we show bounds of the multiplicity by functions of this birational invariant for a singularity of locally a complete intersection. As an application, we obtain the affirmative answer to Watanabe's conjecture on the multiplicity of canonical singularity of locally a complete intersection up to dimension 32.

- 77 橋詰 健太 (京大理) 極小モデル予想と非消滅予想 10
 Kenta Hashizume (Kyoto Univ.) On the minimal model conjecture and the non-vanishing conjecture

概要 The minimal model theory is a fundamental method to classify higher-dimensional algebraic varieties. Today the theory is not completed. It consists of the minimal model conjecture and the abundance conjecture. On the other hand, the non-vanishing conjecture is also an important open problem in the minimal model theory. In fact, the minimal model conjecture and the abundance conjecture implies the non-vanishing conjecture, and it is known by Birkar that the non-vanishing conjecture implies the minimal model conjecture. In this talk I focus on the minimal model conjecture and the simplest case of the non-vanishing conjecture, that is, the non-vanishing conjecture for smooth varieties. I explain that the non-vanishing conjecture for smooth varieties implies the minimal model conjecture.

- 78 福間 慶明 (高知大理工) 偏極多様体の随伴束の大域切断のなす次元について 10
 Yoshiaki Fukuma (Kochi Univ.) On the dimension of global sections of adjoint bundles for polarized manifolds

概要 Let X be a smooth complex projective variety of dimension n , and let L be an ample line bundle on X . Then the pair (X, L) is called a polarized manifold. In my short talk, I will talk about some problems related with the dimension of the global sections of adjoint bundles for polarized manifolds.

- 79 渡邊 健太 (日大理工)* 偏極 K3 曲面上の ACM 直線束について 10
 Kenta Watanabe (Nihon Univ.) On ACM line bundles on polarized K3 surfaces

概要 An ACM bundle on a polarized algebraic variety is defined as a vector bundle whose intermediate cohomology vanishes. We are interested in ACM bundles of rank one with respect to a very ample line bundle on a K3 surface. In this session, we give a necessary and sufficient condition for a non-trivial line bundle M on X with $|M| \neq \emptyset$ and $M^2 \geq L^2 - 6$ to be an ACM and initialized line bundle with respect to L , for a given K3 surface X and a very ample line bundle L on X .

- 80 岩見 智宏 (九工大工)* An analogue of Miyaoka–Yau type inequality for extremal contractions of type (IIA) with special regards to the associated third Chern classes 10
 Tomohiro Iwami (Kyushu Inst. of Tech.) An analogue of Miyaoka–Yau type inequality for extremal contractions of type (IIA) with special regards to the associated third Chern classes

概要 Threefold semistable extremal neighborhoods, as of type (IIA), have important properties as rationality criterion of \mathbb{Q} -conic bundles, or the existence of semistable flips by division algorithm by S. Mori. In this talk, the author reports his study about inducing an analogue of Miyaoka–Yau type inequality for threefold extremal contractions of type (IIA) with special regarding to the third Chern classes associated to the filtration of bi-anti-canonical divisors used in local-to-global deformation in the proof of [Mori1988], based on (non-)normal hyperplane section case of threefold extremal contractions of type (IIA) [Mori–Prokhorov 2016, 2017].

幾何学

3月18日(日) 第I会場

9:15~11:35

- 1 永野哲也 (長崎県立大) 逆線形平行移動を与える曲線の存在について 15
Tetsuya Nagano (Univ. of Nagasaki) On the existence of the curve to give the inverse linear parallel displacement

概要 The existence of the curve to give the inverse linear parallel displacement is not known for a linear parallel displacement of Finsler space generally. A purpose of the study is to establish a mathematical method to find such a curve, but I don't have it, yet. Today, I report a necessary and sufficient condition for two curves that they are to be such curves to give other inverse linear parallel displacement each other.

- 2 竹内 司 (慶大経済) Construction of symplectic-Haantjes manifold of certain Hamiltonian
細川 聖理 systems 10
(日本医師会ORCA管理機構(株))
Tsukasa Takeuchi (Keio Univ.) Construction of symplectic-Haantjes manifold of certain Hamiltonian
Kiyonori Hosokawa systems
(ORCA Management Organization Co., Ltd.)

概要 Symplectic-Haantjes manifolds are constructed for several Hamiltonian systems following Tempesta-Tondo, which yields the complete integrability of systems.

- 3 森山 貴之 (三重大教育) Some examples of global Poisson structures on S^4 10
新田 貴士 (三重大教育)
Takayuki Moriyama (Mie Univ.) Some examples of global Poisson structures on S^4
Takashi Nitta (Mie Univ.)

概要 A Poisson structure is represented by a bivector whose Schouten bracket vanishes. We study a global Poisson structure on S^4 associated with a holomorphic Poisson structure on $\mathbb{C}P^3$. The space of the Poisson structures on S^4 is a real algebraic variety in the space of holomorphic Poisson structures on $\mathbb{C}P^3$. We generalize it to $\mathbb{H}P^n$ by using the twistor method. Furthermore, we provide examples of Poisson structures on S^4 associated with codimension one holomorphic foliations of degree 2 on $\mathbb{C}P^3$.

- 4 森山 貴之 (三重大教育) Splitting theorem for sheaves of holomorphic k -vectors on complex con-
新田 貴士 (三重大教育) tact manifolds 10
Takayuki Moriyama (Mie Univ.) Splitting theorem for sheaves of holomorphic k -vectors on complex con-
Takashi Nitta (Mie Univ.) tact manifolds

概要 A complex contact structure γ is defined by a system of holomorphic local 1-forms satisfying the completely non-integrability condition. The contact structure induces a subbundle $\text{Ker } \gamma$ of the tangent bundle and a line bundle L . In this paper, we prove that the sheaf of holomorphic k -vectors on a complex contact manifold splits into the sum of $\mathcal{O}(\wedge^k \text{Ker } \gamma)$ and $\mathcal{O}(L \otimes \wedge^{k-1} \text{Ker } \gamma)$ as sheaves of \mathbb{C} -module. The theorem induces the short exact sequence of cohomology of holomorphic k -vectors, and we obtain vanishing theorems for the cohomology of $\mathcal{O}(\wedge^k \text{Ker } \gamma)$.

- 5 澤井 洋 (沼津工高専) Examples of solvmanifolds without LCK structures 15
Hiroshi Sawai Examples of solvmanifolds without LCK structures
(Numazu Nat. Coll. of Tech.)

概要 Belgun proved that Inoue surface has no Vaisman structures. In this talk, we generalize this result and construct solvmanifolds without LCK structures. Note that these solvmanifolds have LCS structures.

- 6 五十嵐 雅之 (東京理大基礎工)* Hopf 曲面上に構成される Hermite–Liouville structures について 10
Masayuki Igarashi (Tokyo Univ. of Sci.) On Hermite–Liouville structures constructed on the Hopf surface

概要 In the last 20 years, the Hermite–Liouville structures on compact complex manifolds have been studied. In these studies, almost all of the non-Kählerian structures were obtained by deforming from the Kähler–Liouville structure. In this presentation, the speaker will illustrate the construction of the examples of the Hermite–Liouville structure on the Hopf surface, which leads to the complete integrability of its geodesic flow.

- 7 中村 聡 (東北大理) Hessian of the Ricci Calabi functional 15
Satoshi Nakamura (Tohoku Univ.) Hessian of the Ricci Calabi functional

概要 Ricci Calabi functional is a functional on the space of Kähler metrics of a Fano manifold. Its critical points are called generalized Kähler Einstein metrics. In this talk, we show that the Hessian of the Ricci Calabi functional is non-negative at a generalized Kähler Einstein metric.

- 8 中村 聡 (東北大理) Remarks on modified Ding functional for toric Fano manifolds 10
Satoshi Nakamura (Tohoku Univ.) Remarks on modified Ding functional for toric Fano manifolds

概要 We give a characterization of relative Ding stable toric Fano manifolds in terms of the behavior of the modified Ding functional. We call the corresponding behavior of the modified Ding functional the *pseudo-boundedness from below*. We also discuss the pseudo-boundedness of the Ding / Mabuchi functional of general Fano manifolds.

- 9 久本 智之 (名大多元数理) Ding 汎関数の勾配流と Fano 多様体の最適退化 15
Tomoyuki Hisamoto (Nagoya Univ.) Gradient flow of the Ding energy and optimal degeneration of a Fano manifold

概要 In order to study the optimal degeneration of a Fano manifold, we introduce the Ding flow as the gradient flow of the Ding energy functional on the space of Kahler metrics.

14:15~16:35

- 10 尾國 新一 (愛媛大理)* 粗幾何版アダマール・カルタンの定理について 15
Shin-ichi Oguni (Ehime Univ.) On a coarse Cartan–Hadamard theorem

概要 I will talk about one of coarse geometric versions of the so-called Cartan–Hadamard theorem, that is, a coarse Cartan–Hadamard theorem on coarse convex spaces. Also I will deal with coarse homotopy, open cones and the coarse Baum–Connes conjecture. This talk is based on a joint-work with Tomohiro Fukaya (Tokyo Metropolitan University); ‘Tomohiro Fukaya, Shin-ichi Oguni, A coarse Cartan–Hadamard theorem with application to the coarse Baum–Connes conjecture, preprint, 2017, arXiv:1705.05588’.

- 11 深谷 友宏 (首都大東京理工) 粗幾何学における非正曲率空間の新しい定式化とその境界 15
Tomohiro Fukaya (Tokyo Metro. Univ.) New formulation of nonpositively curved spaces from the view point of coarse geometry and its boundary

概要 In the joint work with Shin-ichi Oguni, we introduced a new class of metric spaces which we call “coarsely convex spaces”. This is a new formulation of “nonpositively curved spaces” from the view point of coarse geometry. This class includes Gromov hyperbolic spaces, CAT(0)-spaces, and systolic complexes. This class is closed under quasi-isometry, and direct product. The idea of the definition is “convexity of metric” and its coarsification. We also construct an ideal boundary for coarsely convex spaces. The construction is based on that of Gromov hyperbolic spaces. Due to time constraints, in this talk, we will not explain on an application to the coarse Baum–Connes conjecture, which is the original motivation of this work.

- 12 豊田 哲 (鈴鹿工高専) Gromov の $\text{Cycl}_k(0)$ 条件の一般化と $\text{CAT}(0)$ 空間の 5 点部分集合の内在的特徴付け 15
 Tetsu Toyoda A generalization of Gromov's $\text{Cycl}_k(0)$ condition and an intrinsic characterization of five points in a $\text{CAT}(0)$ space
 (Suzuka Nat. Coll. of Tech.)

概要 In this talk, we present the result that the $\text{Cycl}_4(0)$ condition implies the $\text{Cycl}_k(0)$ condition for any integer $k \geq 4$. We also present the result that a five-point metric space embeds isometrically into a $\text{CAT}(0)$ space if and only if it satisfies the \boxtimes -inequalities.

- 13 数川 大輔 (東北大理) エネルギー汎関数の収束と曲率次元条件の安定性のための新しい条件 .. 15
 Daisuke Kazukawa (Tohoku Univ.) A new condition for convergence of energies and stability of Ricci curvature bounds

概要 Gigli, Mondino, and Savaré introduced the pmG-convergence on the space of pointed metric measure spaces and studied the stability of the curvature-dimension condition and the Mosco convergence of Cheeger energies under the pmG-convergence. We introduce a new condition generalizing the pmG-convergence and then prove similar results under this condition. Our study is also related to the study by García, Kell, Mondino, and Sosa for quotient spaces by actions of compact groups.

- 14 中島 啓貴 (東北大理) Isoperimetric rigidity and distributions of 1-Lipschitz functions 15
 塩谷 隆 (東北大理)
 Hiroki Nakajima (Tohoku Univ.) Isoperimetric rigidity and distributions of 1-Lipschitz functions
 Takashi Shioya (Tohoku Univ.)

概要 We prove that if a geodesic metric measure space satisfies a comparison condition for the isoperimetric profile and if the observable variance is maximal, then the space is foliated by minimal geodesics, where the observable variance is defined to be the supremum of the variance of 1-Lipschitz functions on the space. Our result can be considered as a variant of Cheeger–Gromoll's splitting theorem and also of Cheng's maximal diameter theorem. As an application, we obtain an isometric splitting theorem for a complete weighted Riemannian manifold with positive Bakry–Émery Ricci curvature.

- 15 中島 啓貴 (東北大理) Lipschitz order with an additive error and normal law à la Lévy on the Hamming cubes 10
 Hiroki Nakajima (Tohoku Univ.) Lipschitz order with an additive error and normal law à la Lévy on the Hamming cubes

概要 M. Gromov introduced the Lipschitz order relation on the set of metric measure spaces and developed a rich theory. For a metric measure space X , we consider the set of the distributions of 1-Lipschitz functions on X and we call it the 1-measurement of X . We also define Lipschitz order on the 1-measurement naturally. The existence of the maximum of 1-measurement is deeply related to the isoperimetric inequality of X . In fact if X is an n -dimensional sphere, we obtain the maximum of 1-measurement by the isoperimetric inequality. However, if X is a n -dimensional Hamming cube, the maximum of 1-measurement does not exist because of discreteness. We solve this problem by generalizing the definition of Lipschitz order with an error. On the spheres case, we have Normal law à la Lévy by considering the weak limit. We have the Hamming cubes version of it as an application of the main theorem.

- 16 山田大貴 (東北大理) The Ricci curvature and the Laplacian on edges of graphs 10
 J. Jost (Max Planck Inst.)
Taiki Yamada (Tohoku Univ.) The Ricci curvature and the Laplacian on edges of graphs
 Jürgen Jost (Max Planck Inst.)

概要 We define the distance between edges of graphs and study the coarse Ricci curvature on edges. We consider the Laplacian on edges based on the definition of the Laplacian on simplicial complexes. As one of our main results, we obtain an estimate of the first non-zero eigenvalue of the Laplacian by the Ricci curvature for a regular graph.

- 17 三石史人 (福岡大理) アレクサンドロフ空間の鈍角定数と体積 15
山口孝男 (京大理)
Ayato Mitsuishi (Fukuoka Univ.) Obtuse constant and volume of Alexandrov spaces
 Takao Yamaguchi (Kyoto Univ.)

概要 We introduce a new geometric invariant called the obtuse constant of spaces with curvature bounded below, defined in terms of comparison angles. We first find relations between this invariant and volume. We discuss the case of maximal obtuse constant equal to $\pi/2$, where we prove some rigidity for spaces. Although we consider Alexandrov spaces with curvature bounded below, the results are new even in the Riemannian case.

16:45~17:45 特別講演

- 田中亮吉 (東北大理)^b 離散群上のポテンシャル論における距離埋め込みの方法
 Ryokichi Tanaka (Tohoku Univ.) Potential theory on discrete groups and metric embedding

概要 We discuss countable groups, mainly focusing on problems related to existence of (bounded) harmonic functions. In particular we study questions such as which group does not admit any non-constant bounded harmonic function (Liouville property), and when it admits such a function, how all such functions are obtained (Poisson boundary). This problem often requires deep understanding on geometry of underlying groups as well as quantitative behavior of random walks. I will try to present this subject with several explicit key examples, emphasizing on importance of combining different ideas and techniques.

3月19日(月) 第I会場

9:00~11:45

- 18 蛭子井博孝 共点, 共線の定理の発見 15
 (幾何数学研究センター)
 Hirotaka Ebisui Discovery of concurrent, collinear Theorems
 (Geomathes Res. Center)

概要 In the history, there are several elementary theorems by figure. We try to find such theorems, and found new 6 theorems. We show entire structures in figures of our text. We explain briefly them by naming as (1) star-star theorem (2) quadrangle Steiner theorem (3) 6 perpendicular-lines theorem (these 3 are concurrent theorems), (4) 10 lines theorem (5) 2 circles system theorem (6) Hexagon Theorem (these 3 are Collinear theorems). These are not proved. But, these theorems include important theoretical structures, and, it is interesting to follow the drawing orders and to consider on the compositions. Anyway, we show all figures of Theorems in our text. Please enjoy many strangeness of theorems. We will be able to speak some relations of theorems in later.

- 19 今井 淳 (千葉大理) \mathbb{R}^n の部分多様体の留数と内在的体積 15
 Jun O'Hara (Chiba Univ.) Residues and intrinsic volumes of submanifolds of \mathbb{R}^n

概要 We show that the intrinsic volumes of compact bodies in the Euclidean spaces of dimension two and three can be obtained from the residues of the (relative) Brylinski beta functions.

- 20 大橋 佳歩 (日本女大理) Delzant 多面体のモジュライ空間上の距離関数の構成 15
 Kaho Ohashi (Japan Women's Univ.) Construction of a metric on the moduli space of Delzant polytopes

概要 A. Pelayo–A. R. Pires–T. S. Ratiu–S. Sabatini defined a metric on the set of Delzant polytopes. They studied structures of the metric space and the moduli space with respect to the action of the integral affine transformations. The definition of the metric is natural, though, it does not induce a metric on the moduli space. In this talk we would like to try to define a metric on the moduli space. We can show that it actually defines a metric on the moduli space for 2-dimensional case.

- 21 佐藤 健治 (玉川大工) 双曲空間の単体の垂心の存在 15
 Kenzi Satô (Tamagawa Univ.) The existence of orthocenters of simplices of hyperbolic spaces

概要 We consider orthocenters of simplices of hyperbolic spaces. Unlike the cases of Euclidean spaces or spheres, the similar condition does not always imply the existence of orthocenters. In this talk, we give characterizations of the existence of orthocenters.

- 22 武富雄一郎 (広島大理) 左不変計量の moduli 空間の特異点及び対応する計量の性質について ... 15
 Yuichiro Taketomi (Hiroshima Univ.) Singularities of the moduli space of left-invariant metrics and the properties of the corresponding metrics

概要 The moduli space of left-invariant metrics on a Lie group is defined as the orbit space of the action of the group of automorphisms and scalings on the space of left-invariant metrics, and has been studied actively. In this talk, we focus on some kinds of singular points which arise in the moduli space. We show that if an equivalent class of a left-invariant metric is the “most singular” point in the moduli space, then the left-invariant metric has nice properties.

- 23 粕谷 直彦 (京都産大理) Knots and links of complex tangents 15
 高瀬 将道 (成蹊大理工)
 Naohiko Kasuya (Kyoto Sangyo Univ.) Knots and links of complex tangents
 Masamichi Takase (Seikei Univ.)

概要 We show that a link in a closed orientable 3-manifold can be realized as the set of complex tangents of a smooth embedding of the 3-manifold into the complex 3-space if and only if it represents the trivial integral homology class in the 3-manifold.

- 24 印南 信宏 (新潟大理) トーラス上の測地円の漸近挙動 15
 Nobuhiro Innami (Niigata Univ.) The asymptotic behavior of geodesic circles in a 2-torus

概要 We prove that in any 2-torus T^2 for any point $p \in T^2$ and for any $\varepsilon > 0$ there exists a number $R > 0$ such that the geodesic circles with center p and radii t are ε -dense in T^2 for all $t > R$.

- 25 近藤 慶 (山口大理) Hopf のピンチング予想から微分異種球面定理へ 15
 Kei Kondo (Yamaguchi Univ.) From Hopf's curvature pinching conjecture to a differentiable exotic sphere theorem

概要 We prove that for an arbitrarily given compact Riemannian manifold M admitting a point $p \in M$ with a single cut point, every compact Riemannian manifold N admitting a point $q \in N$ with a single cut point is diffeomorphic to M if the radial curvature of N at q are sufficiently close in the sense of L^1 -norm to that of M at p .

- 26 庄田敏宏 (佐賀大教育) ラプラシアン第1固有値を最大化する種数2閉曲面上の計量 15
 納谷 信 (名大多元数理)
 Toshihiro Shoda (Saga Univ.) Metrics on a closed surface of genus two which maximize the first eigen-
 Shin Nayatani (Nagoya Univ.) value of the Laplacian

概要 We will consider the upper bound for the first eigenvalue of the Laplacian on a closed surface. For the genus two case, we obtain a singular metric which maximize the first eigenvalue. This result was conjectured by Jakobson–Levitin–Nadirashvili–Nigam–Polterovich. It is joint work with Shin Nayatani, Nagoya University.

13:15~14:15 特別講演

- 笹平裕史 (九大数理)* Seiberg–Witten 方程式とその応用
 Hirofumi Sasahira (Kyushu Univ.) The Seiberg–Witten equations and applications

概要 Since the Seiberg–Witten equations were introduced by Witten in 1994, the equations have produced many significant applications to 3 and 4 dimensional geometry. In this talk, we will discuss some of them and recent progress.

3月20日(火) 第I会場

9:15~11:45

- 27 Jong Taek Cho (Chonnam Nat. Univ.) Realizations of some contact metric manifolds as Ricci soliton real hy-
 橋永貴弘 (北九州工高専) persurfaces 15
 久保 亮 (広島修道大経済)
 武富雄一郎 (広島大理)
 田丸博士 (広島大理)
 Jong Taek Cho (Chonnam Nat. Univ.) Realizations of some contact metric manifolds as Ricci soliton real hy-
 Takahiro Hashinaga persurfaces
 (Kitakyushu Nat. Coll. of Tech.)
 Akira Kubo (Hiroshima Shudo Univ.)
 Yuichiro Taketomi (Hiroshima Univ.)
 Hiroshi Tamaru (Hiroshima Univ.)

概要 Ghosh and Sharma have studied (κ, μ) -spaces, which are contact metric spaces with certain nullity conditions. Especially, they gave a necessary condition for (κ, μ) -spaces to be nongradient Ricci soliton manifolds. In this talk, we prove that in the connected, simply-connected and complete case such (κ, μ) -spaces can be realized as homogeneous real hypersurfaces in noncompact real two-plane Grassmannians. Consequently, we also prove that such spaces are actually Ricci soliton.

- 28 小林和志 (千葉大理) トーラス上のホモロジー的ミラー対称性に関するいくつかの注意点 15
 Kazushi Kobayashi (Chiba Univ.) Some remarks on the homological mirror symmetry for tori

概要 Let us consider an n -dimensional complex torus whose period matrix is (I_n, T) . Here, I_n is the identity matrix of order n and T is a complex matrix of order n whose imaginary part is positive definite. In particular, when we consider the case of $n = 1$, i.e., a one-dimensional complex torus, the corresponding complexified symplectic form of the mirror partner of the one-dimensional complex torus is defined by $-\frac{1}{T}$ or T . However, if we assume $n \geq 2$ and that T is a singular matrix, we can not define the mirror partner of the complex torus as the natural generalization of the case of $n = 1$ to the higher dimensional case. In this talk, we propose a way to avoid this problem, and discuss the homological mirror symmetry.

- 29 伊藤 光弘 (筑波大数理物質) 超幾何型調和多様体と球 Fourier 変換論 10
 佐藤 弘康 (日本工大工)
 Mitsuhiro Itoh (Univ. of Tsukuba) Harmonic manifolds of hypergeometric type and spherical Fourier trans-
 Hiroyasu Satoh (Nippon Inst. of Tech.) form

概要 We define a class of harmonic Hadamard manifolds of hypergeometric type. This class of harmonic manifolds includes all Damek–Ricci spaces and also all rank one symmetric spaces of non-compact type as particular cases. Using a hypergeometric description of spherical functions on each harmonic Hadamard manifold X belonging to this class, we discuss harmonic analysis of radial functions on X . In this talk we would like to present the inversion formula, Plancherel theorem and Paley–Wiener type theorem for the spherical Fourier transform on a Hadamard harmonic manifold which is of hypergeometric type.

- 30 坊向 伸隆 (大分大理工) 楕円軌道上の等質正則直線束にまつわる実半単純リー群の既約ユニタリ表現 10
 Nobutaka Boumuki (Oita Univ.) Irreducible representations of real semisimple Lie groups related to homogeneous holomorphic line bundles over elliptic orbits

概要 In this talk I consider a homogeneous holomorphic line bundle over a certain elliptic (adjoint) orbit, and set a representation of real semisimple Lie group on a complex vector subspace of the complex vector space of holomorphic cross-sections of the bundle. Then, I state that the representation is irreducible unitary.

- 31 渡部 淳 (東大数理)^b Fibered cusp b -pseudodifferential operators and its applications 15
 Jun Watanabe (Univ. of Tokyo) Fibered cusp b -pseudodifferential operators and its applications

概要 Let X be a smooth compact manifold with corners which has two embedded boundary hypersurfaces $\partial_0 X, \partial_1 X$, and suppose a fiber bundle $\phi : \partial_0 X \rightarrow Y$ is given. We define a pseudodifferential calculus $\Psi_{\Phi, b}^*(X)$ generalizing the Φ -calculus of Mazzeo–Melrose and the b -calculus of Melrose. We investigate the Fredholm condition and the index of an operator $P \in \Psi_{\Phi, b}^*(X)$. And as its application, we prove the index theorem of “non-closed” \mathbb{Z}/k -manifolds.

- 32 小磯 憲史 (九大IMI・阪大*)^b Riemann 多様体上の弾性曲線の波動運動方程式 15
 Norihito Koiso A wave equation of motion of an elastic wire on a Riemannian manifold
 (Kyushu Univ./Osaka Univ.*)

概要 A wave equation of motion of an elastic wire on a Riemannian manifold has a solution for any initial data.

- 33 梶ヶ谷 徹 (MathAM-OIL) 重み付きハミルトン安定性と変形ラグランジュ平均曲率流 15
 國川 慶太 (東北大AIMR)
 Toru Kajigaya (MathAM-OIL) Hamiltonian stability for weighted measure and generalized Lagrangian
 Keita Kunikawa (Tohoku Univ.) mean curvature flow

概要 We generalize several results for the Hamiltonian stability and the mean curvature flow of Lagrangian submanifolds in a Kähler–Einstein manifold to more general Kähler manifolds including a Fano manifold equipped with a Kähler form $\omega \in 2\pi c_1(M)$. Namely, we consider a variational problem for Lagrangian submanifolds in a Kähler manifold M w.r.t. a weighted volume functional. Moreover, we introduce the generalized Lagrangian mean curvature flow in a Fano manifold, and we show that if the initial Lagrangian is a small Hamiltonian deformation of a minimal and Hamiltonian stable Lagrangian w.r.t. the weighted volume functional, then the generalized MCF converges exponentially fast to a minimal Lagrangian submanifold.

- 34 小磯 深 幸 (九 大 I M I) 特異点を持つ非等方的平均曲率一定閉超曲面の非一意性と非等方的平均曲率流方程式への応用 15
Miyuki Koiso (Kyushu Univ.) Non-uniqueness of closed non-smooth hypersurfaces with constant anisotropic mean curvature

概要 An anisotropic surface energy is a generalization of the area of surfaces. It is the integral of an energy density function which depends on the surface normal over the considered surface, and it serves as a mathematical model of energy of crystals. The absolute minimizer of an anisotropic surface energy functional among all closed surfaces enclosing the same volume is unique and it is called the Wulff shape. In this talk, we show that, if the energy density function is not “convex”, there exist closed equilibrium surfaces of the anisotropic surface energy for volume-preserving variations which are not the Wulff shape. By applying this result, it is shown that the uniqueness for closed self-similar solutions with genus zero for anisotropic mean curvature flow does not hold in general. These concepts and results are naturally generalized to higher dimensions.

- 35 小磯 深 幸 (九 大 I M I) 特異点を持つ安定な非等方的平均曲率一定閉超曲面の一意性 10
Miyuki Koiso (Kyushu Univ.) Uniqueness of stable closed non-smooth hypersurfaces with constant anisotropic mean curvature

概要 An anisotropic surface energy is a generalization of the area of surfaces. It is the integral of an energy density function which depends on the surface normal over the considered surface, and it serves as a mathematical model of energy of crystals. The absolute minimizer of an anisotropic surface energy functional among all closed surfaces enclosing the same volume is unique and it is called the Wulff shape. In this talk, we show that, if the energy density function is of C^3 and “convex”, then any stable closed equilibrium surface of the anisotropic surface energy for volume-preserving variations is (up to homothety and translation) the Wulff shape, here an equilibrium surface is said to be stable if the second variation of the energy for all admissible variations is nonnegative. The result holds also for hypersurfaces in any Euclidean space.

14:15~16:30

- 36 榎 真 (弘前大理工)* 5次元空間形の極小曲面の変換 10
Makoto Sakaki (Hirosaki Univ.) Transforms for minimal surfaces in 5-dimensional space forms

概要 We discuss transforms for minimal surfaces in 5-dimensional Riemannian space forms, and Lorentzian minimal surfaces in the 5-dimensional semi-Euclidean space of index 2.

- 37 榎 真 (弘前大理工)* 3次元ユークリッド空間の非共形調和曲面の変換と表現公式 15
Makoto Sakaki (Hirosaki Univ.) Transforms and a representation formula for non-conformal harmonic surfaces in the Euclidean 3-space

概要 We give transforms and a representation formula for non-conformal harmonic surfaces in the Euclidean 3-space.

- 38 馬場 蔵人 (東京理大理工) キャリブレーション等式と超極作用 15
井川 治 (京都工繊大工芸)
笹木 集夢 (東海大理)
Kurando Baba (Tokyo Univ. of Sci.) Calibrated equalities and hyperpolar actions
Osamu Ikawa (Kyoto Inst. Tech.)
Atsumu Sasaki (Tokai Univ.)

概要 In this talk, we give a proof for calibrated equalities from the viewpoint of group actions. The essential part of our proof is to describe the orbit spaces for certain group actions on oriented Grassmann manifolds.

- 39 馬場 蔵人 (東京理大理工) 擬リーマン幾何学におけるキャリブレーション不等式と双対性 10
 井川 治 (京都工繊大工芸)
 笹木 集夢 (東海大理)
 Kurando Baba (Tokyo Univ. of Sci.) Calibrated inequalities in pseudo-Riemannian geometry and a duality
 Osamu Ikawa (Kyoto Inst. Tech.)
 Atsumu Sasaki (Tokai Univ.)

概要 In this talk, we give a proof for Mealy's calibrated inequalities from the view point of group actions. As a typical example, we prove Wirtinger's inequality in terms of the duality which was introduced by the speakers.

- 40 本田 淳史 (横浜国大工)* 混合型曲面の等長変形 10
 Atsufumi Honda Isometric deformations of mixed type surfaces
 (Yokohama Nat. Univ.)

概要 For a regular surface in Lorentz–Minkowski 3-space, a point is called a lightlike point if the first fundamental form is degenerate at the point. In this talk, we prove that any analytic surface admits non-trivial isometric deformations around a non-flat and non-degenerate lightlike point.

- 41 木村 太郎 (鶴岡工高専) Classification of Cartan embeddings which are austere submanifolds .. 15
 間下 克哉 (法政大理工)
 Taro Kimura Classification of Cartan embeddings which are austere submanifolds
 (Nat. Inst. of Tech., Tsuruoka Coll.)
 Katsuya Mashimo (Hosei Univ.)

概要 In this talk, we will classify austere submanifolds in compact simple Lie groups. In particular, we proved that Cartan embeddings which defined by inner automorphisms of finite order $k > 2$ of compact simple Lie groups are not austere.

- 42 井川 治 (京都工繊大工芸) 複素旗多様体内の二つの実形の Floer ホモロジー 15
 入江 博 (茨城大理)
 奥田 隆幸 (広島大理)
 酒井 高司 (首都大東京理工)
 田崎 博之 (筑波大数理物質)
 Osamu Ikawa (Kyoto Inst. Tech.) Lagrangian Floer homology of two real forms in a complex flag manifold
 Hiroshi Iriyeh (Ibaraki Univ.)
 Takayuki Okuda (Hiroshima Univ.)
 Takashi Sakai (Tokyo Metro. Univ.)
 Hiroyuki Tasaki (Univ. of Tsukuba)

概要 An adjoint orbit M of a compact connected semisimple Lie group G is called a complex flag manifold. The intersection of two real forms L_0 and L_1 in a complex flag manifold M is an antipodal set of M . Applying the antipodal structure of the intersection $L_0 \cap L_1$, we calculate the Lagrangian Floer homology $HF(L_0, L_1 : \mathbb{Z}_2)$, when M has a G -invariant Kähler–Einstein metric and when two involutions of G defining L_0 and L_1 commute with each other.

- 43 小森 洋平 (早大教育) Construction of pseudo-Anosov automorphisms whose dilatations are
 2-Salem numbers 15
 Yohei Komori (Waseda Univ.) Construction of pseudo-Anosov automorphisms whose dilatations are
 2-Salem numbers

概要 We construct a pseudo-Anosov automorphism whose dilatation is a 2-Salem number by means of the spectrum radius of the bicolored Coxeter element of a bipartite Coxeter system.

16:45~17:45 特別講演

宮地 秀樹 (阪大 理) Toward complex analysis on Teichmüller space

Hideki Miyachi (Osaka Univ.) Toward complex analysis on Teichmüller space

概要 In this talk, I will overview the recent progress on my research on the complex analysis on Teichmüller space. The aim of this research is to give a unified treatment between the topological aspect and the complex analytical aspect in Teichmüller theory. I will discuss the infinitesimal deformation of singular Euclidean structures on a surface in aiming for developing the Teichmüller geometry (Extremal length geometry) on Teichmüller space. I also give a formula of the Levi form of the Teichmüller distance and the pluricomplex Green function on the Teichmüller space. If time permits, I will give an idea for unification and a conjecture on the pluricomplex Poisson kernel on the Bers slice.

函数論

3月18日(日) 第VII会場

9:45~11:45

- 1 尾和重義 (大和大教育)* On univalences for Alexander type integrals 15
 斎藤 齊 (群馬工高専)
 J. Sokol (Univ. of Rzeszow)
 布川 護 (群馬大*)
 Shigeyoshi Owa (Yamato Univ.) On univalences for Alexander type integrals
 Hitoshi Saitoh
 (Gunma Nat. Coll. of Tech.)
 Janusz Sokol (Univ. of Rzeszow)
 Mamoru Nunokawa (Gunma Univ.*)

概要 Let \mathcal{H} be the class of functions $f(z)$ which are analytic in the open unit disk \mathbb{U} . Also let \mathcal{A} be the subclass of functions $f(z)$ in \mathcal{H} with $f(0) = 0$ and $f'(0) = 1$. The object of the present talk is to show some properties of functions $f(z)$ in \mathcal{A} concerning with univalences for Alexander type integrals.

- 2 田中清喜 (大同大)^b 多重調和ベルグマン空間上のテプリッツ作用素 15
 Kiyoki Tanaka (Daido Univ.) Toeplitz operators on the polyharmonic Bergman space

概要 In this talk, we consider the polyharmonic Bergman spaces on the unit ball. As previous results, we obtain the estimates for the reproducing kernel of the polyharmonic Bergman space. By using the estimates for this kernel, we give a characterization of bounded positive Toeplitz operators on the polyharmonic Bergman spaces.

- 3 山崎亮介 (学習院高) The realization problem for Jørgensen numbers 15
 山下 靖 (奈良女大理)
 Ryosuke Yamazaki The realization problem for Jørgensen numbers
 (Gakushuin Boys' Senior High School)
 Yasushi Yamashita
 (Nara Women's Univ.)

概要 Hiroki Sato defined the Jørgensen number of a two-generator Kleinian group as a generalization of Jørgensen's inequality. Oichi-Sato asked the following natural problem: for any real number $r \geq 1$, when is there a Kleinian group whose Jørgensen number is equal to r ? In this talk, we will give a complete solution for this realization problem.

- 4 梶野直孝 (神戸大理) Weyl's eigenvalue asymptotics for the Laplacian on circle packing limit sets of certain Kleinian groups 15
 Naotaka Kajino (Kobe Univ.) Weyl's eigenvalue asymptotics for the Laplacian on circle packing limit sets of certain Kleinian groups

概要 This talk will present the author's recent results on the construction of a "canonical" Laplacian on circle packing fractals invariant under the action of certain Kleinian groups and on the asymptotic behavior of its eigenvalues. In the simplest case of the *Apollonian gasket*, Teplyaev (2004) constructed a Laplacian with respect to which the coordinate functions on the gasket are harmonic, and the author has recently proved its uniqueness and discovered an explicit expression of it in terms of the circle packing structure of the gasket, which immediately extends to general circle packing fractals and defines (a candidate of) a "canonical" Laplacian on them. Then the author has further proved *Weyl's asymptotic formula* for the eigenvalues of this Laplacian, when the circle packing fractal is the limit set of certain Kleinian groups.

- 5 雪田友成 (早大教育) 増大度が Perron 数の 4 次元双曲理想コクセター多面体の無限系列の構成
 15
 Tomoshige Yukita (Waseda Univ.) Construction of infinite series of ideal hyperbolic Coxeter 4-polytopes
 whose growth rates are Perron numbers.

概要 By glueing the ideal hyperbolic Coxeter 4-pyramids, we construct new infinite series of non-simple ideal hyperbolic Coxeter 4-polytopes. In this way, we provide a first example of such a non-compact infinite polyhedral series and prove that their growth rates are Perron numbers.

- 6 柳下剛広 (山口大工) p 乗可積分タイヒミュラー空間上の p -Weil-Petersson 計量の構成について
 15
 Masahiro Yanagishita Construction of p -Weil-Petersson metric on p -integrable Teichmüller
 (Yamaguchi Univ.) space

概要 If $p \geq 2$ and a Riemann surface R satisfies Lehner's condition, then the p -integrable Teichmüller space $T^p(R)$ has a complex Banach manifold structure modeled on p -integrable harmonic Beltrami differentials on R . When $p = 2$, then $T^2(R)$ has a complex Hermitian metric, which is called the Weil-Petersson metric. It was shown that this metric is Kähler and has the negative holomorphic sectional curvature and negative Ricci curvature.

In this talk, we construct the p -Weil-Petersson metric on $T^p(R)$ similarly to the Weil-Petersson metric on $T^2(R)$. In particular, we will say that this metric is smooth and strongly pseudoconvex when p is an even number.

- 7 佐々木東容 (早大理工) 測地カレントとサブセットカレントの稠密性問題 15
 Dounnu Sasaki (Waseda Univ.) The denseness problem of geodesic currents and subset currents

概要 For a (Gromov) hyperbolic group G , which is a generalization of free groups (of finite rank) and surface groups, we can define the space $GC(G)$ of geodesic currents on G and the space $SC(G)$ of subset currents on G . The space $GC(G)$ is proved by Bonahon to be the completion of the set of conjugacy classes of cyclic subgroups of G with positive real weight. The space $SC(G)$ is expected to be the completion of the set of conjugacy classes of quasi-convex subgroups of G with positive real weight, which is still an open problem in general. In the case that G is a free group, Kapovich-Nagnibeda solved the problem. We solve the problem in the case of a surface group.

14:15~15:15 特別講演

- 角大輝 (京大人間環境) ランダム複素力学系におけるランダム性誘起現象とその応用
 Hiroki Sumi (Kyoto Univ.) Randomness-induced phenomena in random holomorphic dynamical systems and their applications

概要 We consider random holomorphic dynamical systems. There are many new phenomena caused by the effect of randomness in random dynamical systems which cannot hold in deterministic dynamical systems. Such phenomena are called randomness-induced phenomena (or noise-induced phenomena). In this talk, we see some randomness-induced phenomena in random holomorphic dynamical systems and some applications of such phenomena to random relaxed Newton's methods.

3月19日(月) 第VII会場

10:00~11:40

- 8 稲山 貴大 (東大数理) 特異 Hermite 計量を持つベクトル束に関する L^2 評価式とコホモロジーの消滅定理について 15

Takahiro Inayama (Univ. of Tokyo) L^2 estimates and vanishing theorems for holomorphic vector bundles equipped with singular Hermitian metrics

概要 We investigate singular Hermitian metrics on vector bundles, especially strictly Griffiths positive ones. L^2 estimates and vanishing theorems usually require an assumption that vector bundles are Nakano positive, however there is no general definition of the Nakano positivity in the singular settings. In this talk, we show some L^2 estimates and vanishing theorems by assuming that the vector bundle is strictly Griffiths positive and the base manifold is projective.

- 9 細野 元気 (東大数理) L^2 拡張定理に関する最近の結果 15

Genki Hosono (Univ. of Tokyo) On recent topics on L^2 extension theorems

概要 We give a L^2 -extension theorem of jets with a sharp constant using the method of Berndtsson–Lempert. We explain the result of a jet L^2 -extension theorem obtained by McNeal–Varolin. We also present a method for sharper estimates.

- 10 濱野 佐知子 (阪市大理) 有限種数開リーマン面の \mathbf{a} -スパンと擬凸領域 15

柴 雅和 (広島大*)

山口 博史 (滋賀大*)

Sachiko Hamano (Osaka City Univ.) Variation of the \mathbf{a} -span of an open Riemann surface and pseudoconvexity

Masakazu Shiba (Hiroshima Univ.*)

Hiroshi Yamaguchi (Shiga Univ.*)

概要 Let R be a marked open Riemann surface of finite genus. If there exists a conformal embedding of R into a closed Riemann surface \tilde{R} of the same genus with prescribed homological types of surfaces, \tilde{R} is called a closing of R . A closing of R induces the Riemann's period matrix T of \tilde{R} . Shiba–Yamaguchi investigated the set of all closings of R , and showed that each diagonal element of T is a closed disk \mathfrak{M} in the upper half plane.

We shall study variation of the period matrices $T(t)$ of the closings of an open Riemann surface $R(t)$ with complex parameter t , and show the rigidity of hyperbolic diameter of $\mathfrak{M}(t)$ under the pseudoconvex variation of $R(t)$.

- 11 児玉 秋雄 (金沢大*)^b A localization principle for biholomorphic mappings between the Fock–Bargmann–Hartogs domains 15

Akio Kodama (Kanazawa Univ.*) A localization principle for biholomorphic mappings between the Fock–Bargmann–Hartogs domains

概要 In this talk, we announce that a localization principle for biholomorphic mappings between equidimensional Fock–Bargmann–Hartogs domains holds. As an application of this, we can show that any proper holomorphic mapping between two equidimensional Fock–Bargmann–Hartogs domains satisfying some condition is necessarily a biholomorphic mapping.

- 12 泊 昌 孝 (日 大 文 理)* 2次元正規2重点の極大イデアル因子と基本因子 15
 都 丸 正 (群 馬 大*)
 Masataka Tomari (Nihon Univ.) maximal ideal cycle and fundamental cycle of normal two dimensional
 Tadashi Tomaru (Gunma Univ.*) double points

概要 Let Z be the Artin fundamental cycle of a resolution of normal complex singularity of the form $z^2 - f(x, y) = 0$. Once studied by Laufer around 1980's, $z^2 = y(x^4 + y^6)$ is famous as the example of the case $Z^2 = -1$ where the maximal ideal cycle M does not equals Z in all the resolution of singularities. We characterize the condition $Z^2 = -1$ completely by means of equisingular class of $f(x, y) = 0$ the numerical characters including a Puiseux pairs and other invariants. For the equality $Z = M$ problem, we can extend Dixon's theorems to the complete criterion about $Z = M$ in terms of numerical characters of $f(x, y) = 0$.

- 13 鍋 島 克 輔 (徳 島 大 理 工) 収束冪級数環での integral number の計算 II 10
 田 島 慎 一 (筑波大数理物質)
 Katsusuke Nabeshima An algorithm for computing integral numbers in a ring of convergent
 (Tokushima Univ.) power series II
 Shinichi Tajima (Univ. of Tsukuba)

概要 The notion of integral closure of an ideal is a key concept in commutative algebra and in singularity theory. Integral numbers, the degrees of integral dependence relations, are also of considerable importance. Effective methods for computing integral numbers w.r.t. an ideal are required. In this talk, first we give an algorithm for computing integral numbers w.r.t. an ideal in a ring of convergent power series. Second, we report some integral numbers of Malgrange's singularities and Skoda-Briançon's singularities.

13:15~14:15 特別講演

- 小 池 貴 之 (阪 市 大 理) 数値的半正な正則直線束のエルミート計量と部分多様体の近傍
 Takayuki Koike (Osaka City Univ.) Hermitian metrics on numerically effective line bundles and neighborhoods of complex submanifolds

概要 Let X be a complex manifold and $Y \subset X$ be a compact complex submanifold of X . Our main interest is in the complex analytical structure of a tubular neighborhood of Y . A motivation comes from a study of Hermitian metrics on a holomorphic line bundle L on X , especially when L has a semi-positivity property in numerical (i.e. intersection-theoretical) sense, called "numerically effective (nef)". We will explain the relationship between the study of neighborhoods of complex submanifolds and of numerically effective line bundles. We will also talk on other applications of the study of neighborhoods of complex submanifolds.

関 数 方 程 式 論

3月18日(日) 第IX会場

9:15~12:00

- 1 M. Adimy (Univ. of Lyon) 時間遅れを持つ非局所的な造血幹細胞ダイナミクスの数理モデル解析 .. 10
 A. Checkroun (Univ. of Tlemcen)
 國谷紀良 (神戸大システム情報)
 Mostafa Adimy (Univ. of Lyon) Analysis of a mathematical model for hematopoietic stem cells with a
 Abdennasser Checkroun nonlocal term and time delay
 (Univ. of Tlemcen)
 Toshikazu Kuniya (Kobe Univ.)

概要 In this study, we investigate a mathematical model of hematopoietic stem cells. The model is described by a system of partial differential equations, which depend on space and age. By applying the method of characteristics, we reformulate the model into a reaction-diffusion equation with a nonlocal spatial term and time delay. We prove the existence, uniqueness and positivity of the solution, and obtain a threshold condition for the global asymptotic stability of the trivial equilibrium. In addition, we obtain sufficient conditions for the existence of nontrivial equilibrium and the uniform persistence of the system.

- 2 泉 英明 (千葉工大情報) 次元数を用いた関数方程式の解法 10
 Hideaki Izumi (Chiba Inst. of Tech.) Solving functional equations by using dimensioned numbers

概要 In this talk, we introduce the notion of dimensioned numbers and extended dimensioned numbers. The notion of dimensioned numbers originates in geometric measurements, and they can be used to describe iterated exponential functions of a single variable. We apply the theory of dimensioned numbers to solving some functional equations of a single variable.

- 3 松永秀章 (阪府大工) 時間遅れをもつ非線形差分方程式の振動条件と相平面解析 10
 土井パティ
 杉江実郎 (島根大総合理工)
 Hideaki Matsunaga (Osaka Pref. Univ.) Oscillation criteria for a nonlinear delay difference equation by phase
 Doi Pati plane analysis
 Sugie Jitsuro (Shimane Univ.)

概要 The oscillation problem of a nonlinear delay difference equation is studied. Sufficient conditions for all solutions of the equation to be oscillatory and for the existence of a nonoscillatory solution are established. Our main results are proved by use of the phase plane analysis which is developed in a similar way to Sugie and Ono [4] in 2004.

- 4 藤田安啓 (富山大理) 病的関数を初期値とする Hamilton–Jacobi flow について 10
 浜向直 (北大理)
 アントニオシコノルフィ
 (Sapienza Univ. of Rome)
 山口範和 (富山大人間発達)
 Yasuhiro Fujita (Univ. of Toyama) On a Hamilton–Jacobi flow starting from a pathological function
 Nao Hamamuki (Hokkaido Univ.)
 Antonio Siconolfi
 (Sapienza Univ. of Rome)
 Norikazu Yamaguchi (Univ. of Toyama)

概要 We consider a Hamilton–Jacobi flow starting from a pathological function. Here, a function on \mathbf{R} is said to be pathological, if it is everywhere continuous but nowhere differentiable.

- 5 竹内 慎吾 (芝浦工大システム理工) 一般化三角関数の非局所境界値問題への応用 10
 Shingo Takeuchi (Shibaura Inst. of Tech.) Applications of generalized trigonometric functions to a nonlocal boundary value problem

概要 Generalized trigonometric functions (GTFs) are simple generalization of the classical trigonometric functions. GTFs are deeply related to the p -Laplacian, which is known as a typical nonlinear differential operator, and there are a lot of works on GTFs concerning the p -Laplacian. However, few applications to differential equations unrelated to the p -Laplacian are known. We will apply GTFs with two parameters to a nonlinear nonlocal boundary value problem without p -Laplacian.

- 6 藤本 皓大 (阪府大工) $p(t)$ -Laplacian を含む 2 階非線形微分方程式の振動定数 10
 山岡 直人 (阪府大工)
 Kodai Fujimoto (Osaka Pref. Univ.) Oscillation constants for second-order nonlinear differential equations
 Naoto Yamaoka (Osaka Pref. Univ.) with $p(t)$ -Laplacian

概要 We consider the oscillation problem for nonlinear differential equation $(|x'|^{p(t)-2}x')' + (\lambda/t^{p(t)})|x|^{p(t)-2}x = 0$, where λ is a positive constant and $p(t) > 1$ is a nondecreasing and smooth function. Using Riccati technique and function sequence technique, we obtain sufficient conditions for this equation to be (non)oscillatory. The obtained results show that there exists a critical value for this problem.

- 7 梶木屋 龍治 (佐賀大理工) A complete classification of bifurcation diagrams for a class of (p, q) -
 I. Sim (Univ. of Ulsan) Laplace equations 10
 田中 敏 (岡山理大理)
 Ryuji Kajikiya (Saga Univ.) A complete classification of bifurcation diagrams for a class of (p, q) -
 Inbo Sim (Univ. of Ulsan) Laplace equations
 Satoshi Tanaka (Okayama Univ. of Sci.)

概要 We study the bifurcation of positive solutions for the one-dimensional (p, q) -Laplace equation with nonlinear term u^{r-1} . There are five types of order relations for (p, q, r) . We study the exact shape of the bifurcation curve in each type of the order relation. Furthermore, we investigate the asymptotic profile of the normalized solution $u(x)/\|u\|_\infty$ as $\|u\|_\infty \rightarrow 0$ or $\|u\|_\infty \rightarrow \infty$, where $\|u\|_\infty$ denotes the L^∞ -norm of u .

- 8 鬼塚 政一 (岡山理大理) Box dimension of solution curves for a class of two-dimensional linear
 田中 敏 (岡山理大理) differential systems 10
 Masakazu Onitsuka (Okayama Univ. of Sci.) Box dimension of solution curves for a class of two-dimensional linear
 Tanaka Satoshi (Okayama Univ. of Sci.) differential systems

概要 This talk deals with the two-dimensional linear differential system

$$x' = y, \quad y' = -x - h(t)y$$

on $[t_0, \infty)$, where $h \in C^1[t_0, \infty)$ and $h(t) > 0$ for $t \geq t_0$. Criteria to obtain the box dimension of graphs of solution curves are established.

- 9 田中 敏 (岡山理大理) Characteristic equation for autonomous planar half-linear differential systems 10
 鬼塚 政一 (岡山理大理)
 Satoshi Tanaka (Okayama Univ. of Sci.) Characteristic equation for autonomous planar half-linear differential systems
 Masakazu Onitsuka
 (Okayama Univ. of Sci.)

概要 The autonomous planar half-linear differential system is considered, which is a generalization of the autonomous planar linear system. It is well-known that the autonomous planar linear system can be solved by eigenvalues, that is, roots of the characteristic equation. In this talk, the characteristic equation for the autonomous planar half-linear differential system is introduced, and the asymptotic behavior of its solutions is established by roots of the characteristic equation.

- 10 西口 純矢 自然な計量をもたない力学系に対する大域アトラクタについて 10
 (東北大数理学連携研究センター)
 Junya Nishiguchi (Tohoku Univ.) On global attractors for dynamical systems without natural metrics

概要 A *global attractor* is a notion for a topological semi-dynamical system whose phase space is a metric space. In particular, this notion is important when the phase space is an infinite-dimensional Banach space by the finiteness of its fractal dimension. However, it should be noticed that the notion of a global attractor depends on the specific choice of a metric. In this talk, we “define” global attractors in the context of the “non-existence of natural metrics” of the phase space and study those properties. This includes a case where the phase space is a Fréchet space, which is motivated by differential equations with unbounded delay. We obtain sufficient conditions for the existence, which will be applied to such equations.

- 11 竹井 優美子 (神戸大理) 2変数超幾何微分方程式の Voros 係数の位相的漸化式による表示とその応用 10
 Yumiko Takei (Kobe Univ.) On the expression of Voros coefficients for hypergeometric differential equations with two variables in terms of the topological recursion and its applications

概要 Voros coefficients are important objects in exact WKB analysis to study global behavior of solutions of differential equations. In this talk I will report that the Voros coefficients for hypergeometric differential equations with two variables are given by the generating functions of free energies defined in terms of Eynard and Orantin’s topological recursion.

- 12 齋藤 三郎 (群馬大*再生核研)* Incompleteness of the theory of differential equations and open problems
 H. Akca (Abu Dhabi Uni.) 10
 S. Pinelas
 (Military Acad. Portugal)
 Saburou Saitoh Incompleteness of the theory of differential equations and open problems
 (Gunma Univ.*/Inst. of Reproducing Kernels)
 Haydar Akca (Abu Dhabi Uni.)
 Sandra Pinelas
 (Military Acad. Portugal)

概要 From the viewpoint of the division by zero ($0/0 = 1/0 = z/0 = 0$) and the division by zero calculus ($\tan(\pi/2) = 0$), we will show some incompleteness of the theory of differential equations in an undergraduate level and we will propose fundamental open problems as the results.

Other topics in this talk: Differential equations with singularities; Continuation of solution; Singular solutions; Solutions with singularities; Solutions with an analytic parameter; Special reductions by division by zero of solutions; Partial differential equations; Introduction of $\log 0 = \log \infty = 0$; and Applications of $\log 0 = 0$; $e^0 = 1, 0$.

- 13 齋藤三郎 (群馬大*・再生核研)* Division by zero calculus and singular integrals 10
 松浦勉 (群馬大工)
 Saburo Saitoh Division by zero calculus and singular integrals
 (Gunma Univ.*/Inst. of Reproducing Kernels)
 Tsutomu Matsuura (Gunma Univ.)

概要 Here, we will give the interpretation for the Hadamard finite part of singular integrals by means of the division by zero calculus - $\log 0 = \log \infty = 0$ (not as limiting values) in the meaning of the one point compactification of Aleksandrov.

14:15~16:15

- 14 原宇信 p -優調和関数に対する Carleson 評価について 10
 (北大理・首都大東京理工)
 Takanobu Hara A Carleson-type estimate for p -superharmonic functions
 (Hokkaido Univ./Tokyo Metro. Univ.)

概要 Let Ω be a domain in \mathbb{R}^n with $n \geq 2$, and let u be a nonnegative p -superharmonic function in Ω . Kilpeläinen and Malý proved that there exists a constant $C > 0$ such that

$$u(x) \leq C \left(\inf_{B(x,R)} u + \mathbf{W}_p^\mu(x, 2R) \right)$$

whenever $B(x, 2R) \subset \Omega$, where μ is the Riesz measure of u and $\mathbf{W}_p^\mu(x, 2R)$ is the Wolff potential of μ . In this paper, we extend this inequality to near the boundary of Ω . More precisely, we give a pointwise estimate for p -superharmonic functions which vanish on the boundary and a global integrability estimate of p -superharmonic functions. Combining the two estimates, we give an analog of the Carleson estimate.

- 15 劉曉静 (茨城大理) The equivalences among p -capacity, p -Laplace-capacities and Hausdorff
 堀内利郎 (茨城大理) measure 10
 Xiaojing Liu (Ibaraki Univ.) The equivalences among p -capacity, p -Laplace-capacities and Hausdorff
 Toshio Horiuchi (Ibaraki Univ.) measure

概要 Let Ω be a smooth bounded domain of \mathbf{R}^N . In this paper, we study the equivalences among p -capacity, p -Laplace-capacities and Hausdorff measure. Firstly we present the equivalence between p -capacity $C_p(K)$ and p -Laplace-capacity $C_{\Delta_p}(K)$ relative to Ω for a given compact set $K \subset \Omega$. Secondly we establish the equivalence between p -Laplace capacity $C_p(K, \partial\Omega)$ relative to $\partial\Omega$ and Hausdorff measure $\mathcal{H}^{N-1}(K)$ on $\partial\Omega$ for a given compact set $K \subset \partial\Omega$.

- 16 A. R. Mulet (北大理) 細い直線的な弾性体の固有振動 10
 Albert Rodríguez Mulet Eigenfrequencies of a thin straight elastic body
 (Hokkaido Univ.)

概要 The Lamé operator is a 2nd order linear elliptic differential operator frequently used to describe the oscillations that take place in a uniform isotropic elastic body. When the oscillations are time-periodic, the differential equation can be simplified to the spectral analysis of the Lamé operator. In our case, we study the case of a thin straight elastic body such that it has its ends fixed. In particular we provide results about the asymptotic behaviour of the eigenvalues and eigenfunctions as the domain gets thinner.

- 17 古屋貴士 (名大多元数理) An application of the factorization method for some inverse acoustic scattering problems 10
 Takashi Furuya (Nagoya Univ.) An application of the factorization method for some inverse acoustic scattering problems

概要 We consider the inverse scattering problem of time-harmonic acoustic plane waves by multiple impenetrable obstacles. For the purpose, we derive the factorization method of Kirsch, which is a sampling method for solving certain kinds of inverse problems where the shape and location of a domain have to be reconstructed. We introduce new results to reconstruct the unknown obstacles by the factorization method. The main idea is to modify the original factorization method by using a priori known outer and inner estimations for a part of unknown obstacles. By our work, we can expand the application of the factorization method for some inverse acoustic scattering problems.

- 18 中島主恵 (東京海洋大海洋) Multiple existence of indefinite nonlinear diffusion problem in population genetics 10
 Kimie Nakashima Multiple existence of indefinite nonlinear diffusion problem in population genetics
 (Tokyo Univ. of Marine Sci. and Tech.)

概要 We consider Lou–Nagylaki conjecture (2002) on a stationary problem of some reaction diffusion equation in population genetics. We deal with the case where spatial dimension is 1. In this case a stationary problem of the equation satisfies $du'' + g(x)u^2(1-u) = 0$ with Neumann zero boundary condition. Under the condition $\int_{\Omega} g(x) dx \geq 0$ and some additional condition on $g(x)$, uniqueness of a nontrivial solution has been already shown. In this talk we construct many nontrivial solutions for some $g(x)$ satisfying $\int_{\Omega} g(x) dx < 0$.

- 19 大枝和浩 (早大 GEC) Classification of steady states to a prey-predator model with population flux by attractive transition 10
 久藤衡介 (電通大情報理工) flux by attractive transition
 Kazuhiro Oeda (Waseda Univ.) Classification of steady states to a prey-predator model with population flux by attractive transition
 Kousuke Kuto (Univ. of Electro-Comm.)

概要 This talk is concerned with the stationary problem of a diffusive Lotka–Volterra prey-predator model with population flux by attractive transition. We analyze two limiting systems as the nonlinear diffusion coefficient approaches infinity. A main result reveals the global bifurcation structure of positive solutions of one of the limiting systems.

- 20 宮本安人 (東大数理) あるスケール変換による極限方程式とその応用 10
 Yasuhito Miyamoto (Univ. of Tokyo) A limit equation and bifurcation diagrams of semilinear elliptic equations with general supercritical growth

概要 We study radial solutions of the semilinear elliptic equation $\Delta u + f(u) = 0$ under rather general growth conditions on f . We construct a radial singular solution and study the intersection number between the singular solution and a regular solution. An application to bifurcation problems of elliptic Dirichlet problems is given. To this end, we derive a certain limit equation from the original equation at infinity, using a generalized similarity transformation. Through a certain transformation, all the limit equations can be reduced into two canonical cases, i.e., $\Delta u + u^p = 0$ and $\Delta u + e^u = 0$.

- 21 宮本安人 (東大数理) Exact eigenvalues and eigenfunctions for a one-dimensional Gel'fand
若狭徹 (九工大工) problem 10
Yasuhito Miyamoto (Univ. of Tokyo) Exact eigenvalues and eigenfunctions for a one-dimensional Gel'fand
Tohru Wakasa (Kyushu Inst. of Tech.) problem

概要 It is known that every positive solution of a one-dimensional Gel'fand problem can be written explicitly. In this talk we give exact expressions of all the eigenvalues and eigenfunctions of the linearized eigenvalue problem at each solution. We study asymptotic behaviors of eigenvalues and eigenfunctions as the L^∞ -norm of the solution goes to the infinity. We also study the problem $u'' + \lambda e^{-u} = 0$ and the associated linearized problem.

- 22 梅津健一郎 (茨城大教育) Loop components of nontrivial nonnegative solutions for indefinite concave-
U. Kaufmann convex problems 10
(Univ. Nacional de Córdoba)
H. Ramos Quoirin
(Univ. de Santiago de Chile)
Kenichiro Umezū (Ibaraki Univ.) Loop components of nontrivial nonnegative solutions for indefinite concave-
Uriel Kaufmann convex problems
(Univ. Nacional de Córdoba)
Humberto Ramos Quoirin
(Univ. de Santiago de Chile)

概要 In this talk, we discuss the existence of a loop component of nontrivial nonnegative solutions for a concave-convex elliptic problem with the Neumann boundary condition. Positivity for solutions on the loop is also discussed. Our approach relies on bifurcation analysis.

16:30~17:30 2017年度(第16回)日本数学会解析学賞受賞特別講演

竹井義次 (同志社大理工) 楕円函数やパレルヴェ特殊函数のインスタントン展開をめぐって —パレルヴェ方程式の完全 WKB 解析最終章

Yoshitsugu Takei (Doshisha Univ.) On the instanton-type expansions of elliptic functions and Painlevé transcendents —The final stage of the exact WKB analysis for Painlevé equations

概要 The exact WKB analysis is based on the Borel resummation method (or the Borel–Laplace method) and, as its consequence, the instanton-type expansions of Painlevé transcendents play an important role in the exact WKB analysis of Painlevé equations. However, such instanton-type expansions are very wild objects and we cannot expect their convergence in general. In this talk, I would like to propose a new approach to handle the instanton-type expansions of Painlevé transcendents. A key idea of this approach is to use the instanton-type expansions of elliptic functions. The structure of instanton-type expansions of elliptic functions will be also discussed in the talk.

3月19日(月) 第IX会場

9:15~12:00

- 23 豊田 洋平 (阪大基礎工) The behavior of blow-up solutions for mean field equation with multi-intensities 10
 鈴木 貴 (阪大MMDS)
 Yohei Toyota (Osaka Univ.) The behavior of blow-up solutions for mean field equation with multi-intensities
 Takashi Suzuki (Osaka Univ.)

概要 In this talk we consider the minimizing sequence for some energy functional of an elliptic equation associated with the mean field limit of the point vortex distribution one-sided Borel probability measure. If such a sequence blows up, we derive some estimate which is related to the behavior of solution near the blow-up point. Moreover, we study the two-intensities case to derive the sufficient condition for this estimate.

- 24 内免 大輔 (室蘭工大工) Blow-up analysis for sign-changing radial solutions in Trudinger–Moser critical equations in \mathbb{R}^2 10
 M. Grossi (Sapienza Univ. of Roma)
 Daisuke Naimen (Muroran Inst. of Tech.) Blow-up analysis for sign-changing radial solutions in Trudinger–Moser critical equations in \mathbb{R}^2
 Massimo Grossi (Sapienza Univ. of Roma)

概要 In this talk we consider low energy sign-changing radial solutions to a elliptic problem related to the Trudinger–Moser inequality. We study the asymptotic behaviour of them. As a result, we show that when the solution has k interior zeros, it exhibits a multiple blow-up behaviour in the first k nodal sets while it converges to the least energy solution of a critical problem in the $(k+1)$ -th one. We also prove that in each concentration set, with an appropriate scaling, the solution converges to the solution of the classical Liouville problem in \mathbb{R}^2 .

- 25 石関 彩 (千葉大理) 一般化された O'Hara エネルギーの分解について 10
 長澤 壯之 (埼玉大理工)
 Aya Ishizeki (Chiba Univ.) Decomposition of generalized O'Hara's energies
 Takeyuki Nagasawa (Saitama Univ.)

概要 O'Hara defined some knot energies for finding canonical configuration of a knot in a given knot type. One of them is known as the Möbius energy. The authors showed that the Möbius energy can be decomposed into three parts; the first one measures how the curve is bent, the second one does how the curve is twisted, and the third one is an absolute constant. The authors define generalized O'Hara's energies, and announce that they can be decomposed in a similar way under suitable assumptions including the case of O'Hara's $(\alpha, 1)$ energies with $\alpha \in [2, 3)$, which are self-repulsive and bounded for any smooth curves without self-intersections.

- 26 佐藤 洋平 (埼玉大理工) Infinitely many solutions for a nonlinear Schrödinger equation with general nonlinearity 10
 柴田 将敬 (東工大理工)
 Yohei Sato (Saitama Univ.) Infinitely many solutions for a nonlinear Schrödinger equation with general nonlinearity
 Shibata Masataka (Tokyo Tech)

概要 We prove the existence of infinitely many solutions for $-\Delta u + V(x)u = f(u)$ in \mathbb{R}^N , $u \in H^1(\mathbb{R}^N)$, where $V(x)$ satisfies $\lim_{|x| \rightarrow \infty} V(x) = V_\infty > 0$ and some conditions. We require conditions of $f(u)$ only around 0 and at ∞ .

- 27 舘山翔太 (東北大理) Fully nonlinear parabolic equations with unbounded ingredients ····· 10
小池茂昭 (東北大理)
A. Świąch (Georgia Tech)
Shota Tateyama (Tohoku Univ.) Fully nonlinear parabolic equations with unbounded ingredients
Shigeaki Koike (Tohoku Univ.)
Andrzej Świąch (Georgia Tech)

概要 The weak Harnack inequality for L^p -viscosity supersolutions of fully nonlinear second-order uniformly parabolic partial differential equations with unbounded coefficients and inhomogeneous terms is established. It is shown that Hölder continuity of L^p -viscosity solutions is derived from the weak Harnack inequality for L^p -viscosity supersolutions. Furthermore, the local maximum principle for L^p -viscosity subsolutions is shown. By these properties, the Harnack inequality for L^p -viscosity solutions is obtained. Several further remarks are presented.

- 28 舘山翔太 (東北大理) The Phragmén–Lindelöf theorem for fully nonlinear parabolic equations with unbounded ingredients ····· 10
Shota Tateyama (Tohoku Univ.) The Phragmén–Lindelöf theorem for fully nonlinear parabolic equations with unbounded ingredients

概要 In 1957, A. Friedman proved the Phragmén–Lindelöf theorem for classical solutions of linear parabolic equations in cones, whose axis is the positive t -axis and whose vertex is the origin of \mathbb{R}^{n+1} . We establish the Phragmén–Lindelöf theorem for fully nonlinear uniformly parabolic equations with unbounded coefficients in same domains.

- 29 梶原直人 (東大数理) Time periodic solutions on real interpolation spaces and its applications
M. Hieber (TU Darmstadt) to some electrophysiological models ····· 10
K. Kress (TU Darmstadt)
P. Tolksdorf (TU Darmstadt)
Naoto Kajiwara (Univ. of Tokyo) Time periodic solutions on real interpolation spaces and its applications
Matthias Hieber (TU Darmstadt) to some electrophysiological models
Klaus Kress (TU Darmstadt)
Patrick Tolksdorf (TU Darmstadt)

概要 We consider the time periodic abstract linear parabolic evolution equation $\partial_t u + Au = f, t \in \mathbb{R}, u(t+T) = u(t)$. We construct the general theory on a real interpolation spaces $D_A(\theta, p)$. It corresponds to time periodic version of the DaPrato–Grisvard maximal $L^p - D_A(\theta, p)$ regularity theorem. Moreover as its application we prove that the nonlinear bidomain equations which is the electrophysiological model have unique time periodic solution near the stable solution by fixed point theorem if the periodic data is sufficiently small.

- 30 梶原直人 (東大数理) Time periodic solutions to the bidomain equations subject to arbitrary
儀我美一 (東大数理) large force ····· 10
K. Kress (TU Darmstadt)
Naoto Kajiwara (Univ. of Tokyo) Time periodic solutions to the bidomain equations subject to arbitrary
Yoshikazu Giga (Univ. of Tokyo) large force
Klaus Kress (TU Darmstadt)

概要 We consider the bidomain equations with Fitzhugh–Nagumo type nonlinear term. We prove that there exists a periodic solution if the data is periodic. Here we do not assume the smallness of the data and the nonlinear term is super-linear. The proof is based on a weak-strong uniqueness argument. To construct the weak periodic solution, we use the Brouwer’s fixed point theorem for the Poincaré map.

- 31 水上雅昭 (東京理大理) The parabolic-elliptic Keller–Segel case as a limit of the fully parabolic Keller–Segel system 10
 Masaaki Mizukami The parabolic-elliptic Keller–Segel case as a limit of the fully parabolic
 (Tokyo Univ. of Sci.) Keller–Segel system

概要 The subject of this work is to construct a new approach to a parabolic-elliptic Keller–Segel system from its parabolic-parabolic case, and to use the parabolic-parabolic case as a step to establish new results in the parabolic-elliptic case. Our aim is, by considering that the parabolic-elliptic case is as a limit of its parabolic-parabolic system, to establish a result such that, only dealing with the parabolic-parabolic Keller–Segel system is enough to obtain the new properties for solutions of its parabolic-elliptic case. In this talk we consider fast signal diffusion limit in a Keller–Segel system, which namely is convergence of a solution for the parabolic-parabolic Keller–Segel system to that for its parabolic-elliptic version.

- 32 Xinru Cao (Paderborn Univ.) Global existence and stabilization in a 3D two-species Keller–Segel–
 来 間 俊 介 (東京理大理) Stokes system with competitive kinetics 10
 水上雅昭 (東京理大理)
 Xinru Cao (Paderborn Univ.) Global existence and stabilization in a 3D two-species Keller–Segel–
 Shunsuke Kurima (Tokyo Univ. of Sci.) Stokes system with competitive kinetics
 Masaaki Mizukami
 (Tokyo Univ. of Sci.)

概要 This talk deals with a 3D two-species Keller–Segel–Stokes system with competitive kinetics. Recently, in a 3D two-species chemotaxis-Stokes system Cao–K.–Mizukami proved global existence and asymptotic behaviour of classical solutions under some conditions. However, the same argument as in the previous work could not be applied to the present problem. The present work asserts global existence and asymptotic behaviour of classical solutions for the Keller–Segel setting.

- 33 石田祥子 (千葉大理) Finite-time blow-up in a fully parabolic Keller–Segel system with de-
 横田智巳 (東京理大理) generate diffusion 10
 柱 貴裕 (東京理大理)
 Sachiko Ishida (Chiba Univ.) Finite-time blow-up in a fully parabolic Keller–Segel system with de-
 Tomomi Yokota (Tokyo Univ. of Sci.) generate diffusion
 Takahiro Hashira (Tokyo Univ. of Sci.)

概要 In this talk we will consider the initial boundary problem for degenerate Keller–Segel systems. For the non-degenerate systems, it is known that $q = m + \frac{2}{N}$ (m denotes the intensity of diffusion, q denotes a nonlinearity and N is the space dimension) is the critical condition for boundedness and blow-up. It is expected that the case of degenerate diffusion has the same critical condition. However, for the blow-up results, the previous paper gave only the existence of unbounded solutions, which includes the blow-up in infinite time. This talk hence gives finite-time blow-up of energy solutions from the initial data with a large negative energy.

- 34 小川卓克 (東北大理) 質量臨界指数における退化移流拡散方程式の解の非有界性と球対称解の
和久井洋司 (東北大理) 凝集質量の評価 10
Takayoshi Ogawa (Tohoku Univ.) Unboundedness of solutions to a degenerate drift-diffusion equation
Hiroshi Wakui (Tohoku Univ.) equation with the mass critical exponent and estimates of the concentration quantity of radially symmetric solutions

概要 We show unboundedness of solutions to a degenerate drift-diffusion equation with the mass critical exponent and estimates of the concentration quantity of radially symmetric solutions. When a given initial datum has finite second moment and the energy functional of it is initially negative, it is known that the corresponding solution blows up in finite time. We prove that solutions to our problem do not remain bounded in the energy space even if we do not impose such a weight condition under the negative energy condition. In particular, if the solution is radially symmetric and the energy functional of the initial datum is negative, then we can eliminate the possibility of growing up. Moreover, we give lower estimates of the mass for radially symmetric blow-up solutions.

13:15~14:15 2017年度(第16回)日本数学会解析学賞受賞特別講演

- 柴田徹太郎 (広島大工) 非線形楕円型方程式の固有値問題の漸近解析と逆分岐問題の解析
Tetsutaro Shibata (Hiroshima Univ.) Asymptotic analysis of eigenvalue problems for nonlinear elliptic equations and analysis of inverse bifurcation problems

概要 We first study the global and local behavior of bifurcation curves for elliptic nonlinear eigenvalue problems in L^q -framework ($q > 1$). We consider the case where λ is parameterized by the L^q -norm $\alpha = \|u_\lambda\|_q$ of the solution u_λ corresponding to λ and is represented as a continuous function $\lambda = \lambda(\alpha)$. Especially, we restrict our attention to the asymptotic behavior of $\lambda(\alpha)$ as $\alpha \rightarrow \infty$ and $\alpha \rightarrow 0$. We establish several precise asymptotic expansion formulas for $\lambda(\alpha)$ as $\alpha \rightarrow \infty$ and $\alpha \rightarrow 0$ to understand well the total structures of the bifurcation curves. We next consider the inverse bifurcation problems. Especially, we will show some results for inverse problems by using a variational method and asymptotic expansion formulas for $\lambda(\alpha)$ for $\alpha \gg 1$.

3月20日(火) 第IX会場

9:15~12:00

- 35 中村謙太 (九大数理) p -Sobolev 流の解の存在とその性質について 10
三沢正史 (熊本大先端科学)
Kenta Nakamura (Kyushu Univ.) The existence and properties of a solution to the p -Sobolev flow
Masashi Misawa (Kumamoto Univ.)

概要 In this talk, we study a doubly nonlinear parabolic equation, called the p -Sobolev flow, which is the classical Yamabe flow on a bounded domain in Euclidean space. We show the existence of a weak solution to the p -Sobolev flow without geometrical assumption and present properties of its solution.

- 36 L. Cavallina (東北大情報) 不変等流面を有する二相熱伝導体の幾何学的形状 10
R. Magnanini (Univ. of Florence)
 坂口 茂 (東北大情報)
Lorenzo Cavallina (Tohoku Univ.) On the shape of a two-phase heat conductor with a surface of the constant flow property
Rolando Magnanini (Univ. of Florence)
Shigeru Sakaguchi (Tohoku Univ.)

概要 We study how the presence of a surface of the constant flow property influences the shape of a two-phase heat conductor. The existence of a surface that satisfies the constant flow property at every moment in time is a very strong requirement: we show that this condition implies the radial symmetry of our heat conductor. In addition, we study the difference behaviour of two-phase heat conductors satisfying an analogous overdetermined elliptic problem. In this case we are able to construct a family of non radially symmetric solutions.

- 37 可香谷隆 (九大IMI) 非凸領域における Neumann 境界条件付き Allen–Cahn 方程式に対する特異極限問題 10
Takashi Kagaya (Kyushu Univ.) A singular limit problem of Allen–Cahn equation with Neumann boundary condition on non-convex domains

概要 We study a singular limit problem of the Allen–Cahn equation with the homogeneous Neumann boundary condition on non-convex domains with smooth boundaries under suitable assumptions for initial data. The main result is the convergence of the time parametrized family of the diffused surface energy to Brakke’s mean curvature flow with a generalized right angle condition on the boundary of the domain.

- 38 浜向直 (北大理) On a dynamic boundary condition for singular degenerate parabolic equations in a half space 10
 儀我美一 (東大数理)
Nao Hamamuki (Hokkaido Univ.) On a dynamic boundary condition for singular degenerate parabolic equations in a half space
Yoshikazu Giga (Univ. of Tokyo)

概要 We consider a dynamic boundary value problem for singular degenerate parabolic equations in a half space. In the context of viscosity solutions, we establish a comparison principle and prove existence of solutions together with Lipschitz regularity of the unique solution. A relationship with a Dirichlet or Neumann condition is also studied.

- 39 関行宏 (九大数理) 球面に値を取る調和写像流方程式の解の爆発について 10
P. Biernat (Bonn Univ.)
Yukihiro Seki (Kyushu Univ.) On blow-up of solutions for harmonic map heat flow with values in a sphere
Paweł Biernat (Bonn Univ.)

概要 In this talk we will discuss blow-up of a harmonic map heat flow from R^d to S^d , where S^d denotes a unit sphere in R^d . Our main result yields a constructive examples of Type II blow-up solutions for $d \geq 7$. These blow-up solutions satisfy various point-wise estimates in some space-time regions.

- 40 松澤 寛 (沼津工高専) ある Fisher-KPP 方程式の自由境界問題について 10
 Hiroshi Matsuzawa (Numazu Nat. Coll. of Tech.) A free boundary problem for the Fisher-KPP equation with a moving boundary

概要 In this talk, we consider a free boundary problem of Fisher-KPP equation $u_t = u_{xx} + u(1 - u)$, $t > 0$, $ct < x < h(t)$. The number $c > 0$ is a given constant, $h(t)$ is a free boundary which is determined by the Stefan-like condition. This model may be used to describe the spreading of a non-native species over a one dimensional habitat. The free boundary $x = h(t)$ represents the spreading front. In this model, we impose zero Dirichlet condition at left moving boundary $x = ct$. This means that the left boundary of the habitat is a very hostile environment and that the habitat is eroded away by the left moving boundary at constant speed c . In this talk, I will give a trichotomy result, that is, for any initial data, exactly one of the three behaviours, vanishing, spreading and transition, happens.

- 41 内藤雄基 (愛媛大理) Asymptotic behavior of global solutions for semilinear heat equations with slowly decaying initial data 10
 Yūki Naito (Ehime Univ.) Asymptotic behavior of global solutions for semilinear heat equations with slowly decaying initial data

概要 We consider the Cauchy problem for semilinear heat equation in R^N . We study the case where initial data have polynomial decay rate at the spatial infinity, and investigate the convergence property of the global solutions to the forward self-similar solutions.

- 42 板倉恭平 (神戸大理) Spectral theory for repulsive Hamiltonians 10
 Kyohei Itakura (Kobe Univ.) Spectral theory for repulsive Hamiltonians

概要 For repulsive Hamiltonians we obtained Rellich's theorem, the radiation condition and the limiting absorption principle. Our setting include the case of the inverted harmonic oscillator. In the proofs, we mainly use a commutator argument. This argument simple and elementary, and does not employ energy cut-offs or the microlocal analysis.

- 43 一ノ瀬 弥 (信州大理) 連続量子測定 of Feynman 経路積分と非自己共役 Schrödinger 方程式 ... 10
 Wataru Ichinose (Shinshu Univ.) The Feynman path integrals of continuous quantum measurements and the non-self-adjoint Schrödinger equations

概要 My talk is concerning on the the mathematical theory of quantum measurements. Especially, we will consider the continuous quantum measurements of the position of particles during a finite time. The probability amplitude of particles just after the measurement is given in the form of the weighted Feynman path integrals, WFPI, or the restricted Feynman path integrals, according to Feynman and Mensky's theory of quantum measurements. We will show that WFPI for it are defined mathematically in L^2 and the weighted Sobolev spaces, and satisfy the non-self-adjoint Schrödinger equations.

- 44 道久 寛載 (広島大理) 消散波動方程式の波動効果と拡散構造 10
 Hironori Michihisa (Hiroshima Univ.) Diffusive structures and wave effects of damped wave equations

概要 We obtain higher order expansions of evolution operators corresponding to the Cauchy problem of the linear damped wave equation in \mathbf{R}^n . Established hyperbolic part of expansion seems to be new in the sense that the order of the expansion of the hyperbolic part depends on the spatial dimension.

- 45 森 直文 (福岡工大) The S & K mixed condition for symmetric hyperbolic systems with non-symmetric relaxations 10
 Naofumi Mori (Fukuoka Inst. of Tech.) The S & K mixed condition for symmetric hyperbolic systems with non-symmetric relaxations

概要 We consider the new structural conditions to show the decay property of the linear symmetric hyperbolic systems from the viewpoint of the dissipative structure. Especially, we are concerned with the model systems which have a non-symmetric relaxation and therefore their decay estimate is of regularity-loss. Recently, since the Shizuta–Kawashima stability theory cannot be applicable to such a system, the structural conditions to show the decay property of regularity-loss have been investigated. However, we have the last question that the structural condition which can be applied to the Timoshenko–Cattaneo system has not been developed yet. In this talk, we introduce the new structural condition which will be the first condition applicable to the Timoshenko–Cattaneo system and the other systems of the same weakest dissipative mechanism.

- 46 中村 謙太 (九大数理) Asymptotic stability of rarefaction waves for a model system of hyperbolic balance laws 10
 中村 徹 (熊本大工)
 川島 秀一 (九大数理)
 Kenta Nakamura (Kyushu Univ.) Asymptotic stability of rarefaction waves for a model system of hyperbolic balance laws
 Tohru Nakamura (Kumamoto Univ.)
 Shuichi Kawashima (Kyushu Univ.)

概要 In this talk, we addressed with rarefaction waves for a hyperbolic system of balance laws in the whole space or half space. We shall prove a priori estimate of a solution and the asymptotic stability of rarefaction waves by using L^2 -energy method and standard calculus.

- 47 側島 基宏 (東京理大理工) Life-span of blowup solutions to semilinear wave equation with space-dependent critical damping 10
 池田 正弘 (理化学研・慶大理工)
 Motohiro Sobajima Life-span of blowup solutions to semilinear wave equation with space-dependent critical damping
 (Tokyo Univ. of Sci.)
 Masahiro Ikeda (RIKEN/Keio Univ.)

概要 We consider the semilinear wave equation with space-dependent critical damping term in \mathbb{R}^N ($N \geq 3$). The equation is of the form $\partial_t^2 u - \Delta u + V_0 |x|^{-1} \partial_t u = |u|^p$. If $V_0 = 0$, then small data blowup for $p \leq p_0(N)$ and small data global existence for $p > p_0(N)$ are proved with well-known Strauss exponent $p_0(N)$. In this talk we will show that a similar blowup phenomenon occurs for $\frac{N}{N-1} < p < p_0(N + V_0)$.

14:15~16:15

- 48 新村 貴之 (北大理) 梁方程式における回転慣性がアトラクタに与える影響について 10
 Takayuki Niimura (Hokkaido Univ.) Attractors and their stability with respect to rotational inertia for a nonlocal extensible beam equation

概要 We will study the initial-boundary value problem for the Kirchhoff type plate equation with rotational inertia. In particular, we consider the long-time behavior of the solution of the equation and show the existence of attractors and clarify their properties. Furthermore, we will also cover the influences of the rotational inertia on the long-time dynamics.

- 49 池田正弘 (理化学研 AIP・慶大理工) 線形消散型波動方程式の解の時間減衰評価, 可積分でない初期値を持つ非線形問題の臨界指数 10
 若杉勇太 (愛媛大工)
 戌亥隆恭 (東京理大理)
 岡本葵 (信州大工)
Masahiro Ikeda (RIKEN/Keio Univ.) Time decay estimates for the linear damped wave equation and the
Yuta Wakasugi (Ehime Univ.) critical exponent for the semilinear problem with slowly decaying data
Inui Takahisa (Tokyo Univ. of Sci.)
Mamoru Okamoto (Shinshu Univ.)

概要 We study the Cauchy problem of the damped wave equation

$$\partial_t^2 u - \Delta u + \partial_t u = 0$$

and give L^p - L^q estimates of the solution for $1 \leq q \leq p < \infty$ ($p \neq 1$) with derivative loss. We apply this estimate to the nonlinear problem

- 50 池田正弘 (理化学研 AIP・慶大理工) スケール臨界な時間変数に依存した係数を摩擦項に持つ波動方程式の解の最大存在時刻について 10
 側島基宏 (東京理大理工)
Masahiro Ikeda (RIKEN/Keio Univ.) Life-span of solutions to semilinear wave equation with time-dependent
Motohiro Sobajima critical damping
 (Tokyo Univ. of Sci.)

概要 We consider the Cauchy problem of the nonlinear wave equation with a scaling critical time-dependent damping $\mu(1+t)^{-1}u_t$. Here μ is non-zero constant. When $\mu = 0$, the equation becomes the usual wave equation and the critical power dividing small data global existence and blow-up is given by Strauss exponent $p_0(N)$ (N means spatial dimension). We give a small data blow-up result in the case $1 + 2/N < p \leq p_0(N + \mu)$ if μ is near 0.

- 51 吉田夏海 (立命館大 OIC 総合研究機構) Large time behavior of solutions toward a multiwave pattern to the Cauchy problem for the dissipative wave equation with partially linearly degenerate flux 10
Natsumi Yoshida (Ritsumeikan Univ.) Large time behavior of solutions toward a multiwave pattern to the Cauchy problem for the dissipative wave equation with partially linearly degenerate flux

概要 We study the large time asymptotics of solutions to the Cauchy problem for the one-dimensional dissipative wave equation where the far field states are prescribed. Especially, we deal with the case when the flux function is convex or concave but linearly degenerate on some interval. Then the Cauchy problem has a unique global in time solution which tends toward a multiwave pattern consists of rarefaction and viscous contact waves as time goes to infinity. The proof is given by a technical energy method and the careful estimates for the interactions between the nonlinear waves.

- 52 吉田夏海 (立命館大 OIC 総合研究機構) Global asymptotic stability of the rarefaction waves for a scalar conservation law with nonlinear viscosity 10
松村昭孝 (阪大*)
Natsumi Yoshida (Ritsumeikan Univ.) Global asymptotic stability of the rarefaction waves for a scalar conservation law with nonlinear viscosity
Akitaka Matsumura (Osaka Univ.*)

概要 We study the large time asymptotics of solutions to the Cauchy problem for a one-dimensional scalar conservation law with nonlinear viscosity where the far field states are prescribed. Especially, we deal with the case when the flux function is fully convex, and also the viscosity is a nonlinearly degenerate one. Then the Cauchy problem has a unique global in time solution which tends toward a rarefaction wave as time goes to infinity. The proof is given by a technical energy method, and a Sobolev type inequality motivated by an idea of Kanel'.

- 53 川原雄一朗 (同志社中高) Scattering problem for the fourth order nonlinear Schrödinger equation
林仲夫 (阪大理) 10
P. I. Naumkin (UNAM)
Yuichiro Kawahara (Doshisha Junior and Senior High School) Scattering problem for the fourth order nonlinear Schrödinger equation
Nakao Hayashi (Osaka Univ.)
Pavel I. Naumkin (UNAM)

概要 We consider the fourth order nonlinear Schrödinger equation

$$i\partial_t u - \frac{1}{4}\partial_x^4 u = f(u), \quad (t, x) \in \mathbf{R} \times \mathbf{R},$$

where $f(u)$ is the power nonlinearity of order $p > 5$. We show scattering operators are well defined in the neighborhood of the origin of a suitable weighted Sobolev space.

- 54 眞崎聡 (阪大基礎工) 排斥的デルタポテンシャルを持つ非線型シュレディンガー方程式の長距離
J. Murphy (Missouri Univ. of Sci. and Tech.) 離散乱 10
瀬片純市 (東大理)
Satoshi Masaki (Osaka Univ.) Long range scattering for nonlinear Schrödinger equation with repulsive
Jason Murphy (Missouri Univ. of Sci. and Tech.) delta potential
Jun-ichi Segata (Tohoku Univ.)

概要 We consider initial data problem for 1d cubic nonlinear Schrödinger equation with repulsive delta potential. We will show that if a data belongs to a weighted Sobolev space and is sufficiently small then solution decays in time in the same order as a free equation and asymptotically behaves like free solution with a logarithmic phase correction. Recently, long range scattering for nonlinear Schrödinger equation with a potential is extensively studied. In the previous results, a class of smooth and decaying potential is considered. In our case, we fully use explicit formulas which are available in the delta potential case.

- 55 宮崎隼人 (津山工高専) 斉次型臨界非線形項をもつ非線形 Schrödinger 方程式の非散乱について
 眞崎聡 (阪大基礎工) 10
 Hayato Miyazaki Nonexistence of scattering and modified scattering states for nonlinear
 (Tsuyama Nat. Coll. of Tech.) Schrödinger equations with critical homogeneous nonlinearity
 Satoshi Masaki (Osaka Univ.)

概要 We consider large time behavior of solutions to the nonlinear Schrödinger equation with a homogeneous nonlinearity of the critical order which is not necessarily a polynomial. We handle the case in which the nonlinearity contains non-oscillating factor $|u|^{1+2/d}$. It turns out that there is no solution which behaves like a free solution with or without any logarithmic phase corrections. We also prove nonexistence of an asymptotic free solution in the case that the gauge invariant nonlinearity is dominant, and give a small data finite time blow-up result.

- 56 村井宗二郎 (産業技術高専)* 外部領域における磁場付き Schrödinger 方程式の Strichartz 評価と非線形散乱理論 10
 Sojiro Murai Strichartz and scattering for magnetic Schrödinger equations in exterior
 (Tokyo Metropolitan Coll. of Indus. Tech.) domain

概要 In this talk we shall treat the Strichartz estimates for magnetic Schrödinger equations and its application to global existence of the solutions and scattering theory with power type nonlinear term in an exterior domain. The proof relies on similar argument to the case of whole space.

16:30~17:30 特別講演

- 二宮広和 (明大総合数理) 反応拡散方程式の進行波解と全域解
 Hirokazu Ninomiya (Meiji Univ.) Traveling wave solutions and entire solutions of reaction-diffusion equations

概要 When several stable states coexist, propagation phenomena are often observed in many fields including dissipative situations. To characterize the universal profiles of these phenomena, traveling wave solutions and entire solutions play important roles. Here traveling wave solution is meant by a solution of a partial differential equation that propagates with a constant speed, while it maintains its shape in space, and an entire solution is a solution defined for all space and time variables. In this talk we focus on the Allen-Cahn-Nagumo equation, which is a single reaction diffusion equation with bistable nonlinearity and explain how to construct entire solutions and the relation between traveling wave solutions and entire solutions.

3月21日(水) 第IX会場

9:15~12:00

- 57 清水一慶 (京大理) Schrödinger map 方程式の局所適切性について 10
 Ikkei Shimizu (Kyoto Univ.) Remarks on local well-posedness of Schrödinger map equation

概要 We consider the initial value problem for Schrödinger map equation. We provide some supplemental arguments for the work by Gustafson et al. (Duke Math. J. 145(3), 537–583, 2008), in which local well-posedness near the family of harmonic maps is asserted.

- 58 岡本葵 (信州大工) 分数階非線形 Schrödinger 方程式の初期値問題の非適切性 10
 Mamoru Okamoto (Shinshu Univ.) Ill-posedness of the Cauchy problem for the fractional Schrödinger equation

概要 We consider ill-posedness of the Cauchy problem for the fractional Schrödinger equation. More precisely, we prove norm inflation with general initial data. This argument with minor modifications also shows the ill-posedness for the generalized Boussinesq equations.

- 59 迫田大輔 (阪大 理) Small data global existence for a quadratic derivative nonlinear Schrödinger system in two space dimensions 10
 砂川秀明 (阪大 理)
 Daisuke Sakoda (Osaka Univ.) Small data global existence for a quadratic derivative nonlinear Schrödinger system in two space dimensions
 Hideaki Sunagawa (Osaka Univ.)

概要 We consider the initial value problem for a system of quadratic derivative nonlinear Schrödinger equations in two space dimensions with the masses satisfying a suitable resonance relation. We give a structural condition on the nonlinearity under which small data global existence holds. This is an extension of previous results by Hayashi–Li–Naumkin and Ikeda–Katayama–Sunagawa.

- 60 平山浩之 空間 2 次元上の Zakharov–Kuznetsov–Burgers 方程式の適切性について 10
 (宮崎大テニュアトラック推進機構)
 Hiroyuki Hirayama (Univ. of Miyazaki) Well-posedness for the Zakharov–Kuznetsov–Burgers equation in two space dimensions

概要 We consider the Cauchy problem of the Zakharov–Kuznetsov–Burgers equation (ZKB for short) in two space dimensions. By using the Fourier restriction norm with the effect of the dissipative term, we prove the well-posedness in the Sobolev space H^s for $s > -1/2$. It is interesting that ZKB has dissipative effect only x -direction, but the result for the regularity is better than the well-posedness of the Zakharov–Kuznetsov equation for both x and y -directions.

- 61 福田一貴 (北大 理) 一般化された KdV–Burgers 方程式の解の第 2 漸近形 10
 Ikki Fukuda (Hokkaido Univ.) Second asymptotic profile for the generalized KdV–Burgers equation

概要 We study the asymptotic behavior of global solutions to the initial value problem for the generalized KdV–Burgers equation. One can expect that the solution to this equation converges to a self-similar solution to the Burgers equation, due to earlier works related to this problem. Actually, we obtain the optimal asymptotic rate similar to those results and the second asymptotic profile for the generalized KdV–Burgers equation.

- 62 坂本祥太 (京大人間環境) 重み付き Chemin–Lerner 型空間におけるボルツマン方程式の解 10
 段仁軍 (香港中文大)
 Shota Sakamoto (Kyoto Univ.) Solution to the Boltzmann equation in velocity-weighted Chemin–Lerner type spaces
 Duan Renjun (Chinese Univ. of Hong Kong)

概要 We study the Boltzmann equation near global Maxwellians in the d -dimensional whole space. A unique global-in-time mild solution to the Cauchy problem of the equation is established in a Chemin–Lerner type space with respect to the phase variable (x, v) . Both hard and soft potentials with angular cutoff are considered. The new function space for global well-posedness is introduced to essentially treat the case of soft potentials, and the key point is that the velocity variable is taken in the weighted supremum norm, and the space variable is in the s -order Besov space with $s \geq d/2$ including the spatially critical regularity.

- 63 牧野哲 (山口大*) Einstein–Euler 方程式の軸対称解について 10
 Tetu Makino (Yamaguchi Univ.*) On axisymmetric solutions of the Einstein–Euler equations

概要 Axisymmetric solutions to the Einstein equations with the energy-momentum tensor of barotropic perfect fluid can be constructed mathematically as a post Newtonian approximation to slowly rotating axisymmetric solutions to the Euler–Poisson equation of gaseous stars, provided that the adiabatic exponent near the vacuum belongs to the interval $]6/5, 3/2[$.

- 64 橋本伊都子 (関西大システム理工・阪市大数学研) 多次元空間上におけるバーガーズ方程式の球対称問題の漸近挙動について 10
 松村昭孝 (阪大*)
 Itsuko Hashimoto (Kansai Univ./Osaka City Univ.) Asymptotic behavior toward nonlinear waves for radially symmetric solutions of multi-dimensional Burgers equation
 Akitaka Matsumura (Osaka Univ.*)

概要 We concern with the asymptotic behaviors of radially symmetric solutions for multi-dimensional Burgers equation on the exterior domain in \mathbb{R}^n , where the boundary and far field conditions are prescribed. In a case where the corresponding 1-D Riemann problem for the non-viscous part admits a shock wave, we show the solution tends toward a superposition of stationary wave and rarefaction wave as time goes to infinity. We also show the decay rate estimate. Furthermore, for $n = 3$, we give the complete classification of the asymptotic states, which includes even a superposition of stationary wave and viscous shock wave.

- 65 石垣祐輔 (九大数理) Global existence of solutions of the compressible viscoelastic fluid around parallel flow 10
 Yusuke Ishigaki (Kyushu Univ.) Global existence of solutions of the compressible viscoelastic fluid around parallel flow

概要 A system of equations for compressible viscoelastic fluid is considered in an infinite layer. When the external force has a suitable form, the system has a solution of parallel flow type. It is shown that the solution of the system exists globally in time if the initial data is sufficiently close to the one of the parallel flow, provided that the initial data for the parallel flow is sufficiently small and the viscosity coefficient and the shear wave speed are sufficiently large.

- 66 渡邊圭市 (早大理工) Maximal regularity of compressible-incompressible two-phase flows with phase transitions 10
 Keiichi Watanabe (Waseda Univ.) Maximal regularity of compressible-incompressible two-phase flows with phase transitions

概要 In this talk, we consider the free boundary problem for compressible-incompressible two-phase flows with phase transitions in isothermal case. Two fluids are separated by a sharp interface and a surface tension is taken into account. We use the Navier–Stokes–Korteweg equations for the compressible fluid and the Navier–Stokes equations for the incompressible fluid, whose model is thermodynamically correct. We show the maximal L_p - L_q regularity theorem with the help of the \mathcal{R} -bounded solution operators of the corresponding generalized resolvent problem and Weis’s operator-valued Fourier multiplier theorem.

- 67 柴田良弘 (早大理工) 表面張力付き自由境界問題について 10
 Yoshihiro Shibata (Waseda Univ.) Free boundary problem with surface tension

概要 It is presented the local well-posedness of free boundary problem for the Navier–Stokes equations with surface tension without any restriction of the size of initial data. Hanzawa transform is used to represent the free surface. Since the standard linearized procedure requires a smallness restriction for the initial data, to avoid such smallness assumption, I used a modified linearized problem, which was first proposed by V. A. Solonnikov.

- 68 柴田良弘 (早大理工) Global well-posedness for a \mathbb{Q} tensor model of Incompressible Nematic Liquid Crystals in \mathbb{R}^N 10
 Yoshihiro Shibata (Waseda Univ.) Global well-posedness for a \mathbb{Q} tensor model of Incompressible Nematic Liquid Crystals in \mathbb{R}^N

概要 I will talk about the global well-posedness for the Cauchy problem of a \mathbb{Q} tensor model of Incompressible Nematic Liquid Crystals in the N -dimensional Euclidean space. This is a joint work with Maria Schonbeck (Univ. California Santa Cruz). The proof is done by combination of L_p - L_q decay estimations and L_p - L_q maximal regularity for the heat equations and Stokes equations.

- 69 津田和幸 (阪大基礎工) Time decay estimate with diffusive property and smoothing effect for solution to the compressible Navier–Stokes–Korteweg system 10
 小林孝行 (阪大基礎工) solution to the compressible Navier–Stokes–Korteweg system 10
 Kazuyuki Tsuda (Osaka Univ.) Time decay estimate with diffusive property and smoothing effect for solution to the compressible Navier–Stokes–Korteweg system
 Takayuki Kobayashi (Osaka Univ.) solution to the compressible Navier–Stokes–Korteweg system

概要 Time decay estimate of a solution to the compressible Navier–Stokes–Korteweg system is studied. Concerning the linearized problem, the decay estimates with diffusive property for initial data are derived. As an application, the time decay estimates of a solution to the nonlinear problem are given. In contrast to the compressible Navier–Stokes system, for linear system regularities of initial data are lower and independent of the order of derivative of the solution owing to smoothing effect from the Korteweg tensor. Furthermore, for the nonlinear system diffusive properties are obtained with initial data having lower regularity than that of studies of the compressible Navier–Stokes system.

14:15~16:15

- 70 岸本展 (京大数理研) Global mild solution to Navier–Stokes equations with partial hyperviscosity 10
 齐木吉隆 (一橋大商) solution to the compressible Navier–Stokes–Korteweg system 10
 中井拳吾 (東大数理)
 米田剛 (東大数理)
 Nobu Kishimoto (Kyoto Univ.) Global mild solution to Navier–Stokes equations with partial hyperviscosity
 Yoshitaka Saiki (Hitotsubashi Univ.) solution to the compressible Navier–Stokes–Korteweg system 10
 Kengo Nakai (Univ. of Tokyo)
 Yoneda Tsuyoshi (Univ. of Tokyo)

概要 Our purpose is to clarify the energy cascade mechanism for the incompressible Navier–Stokes equations. For the first step, we study the NS equations with partial hyperviscosity (dissipation is removed from some of the low Fourier modes) with DNS. This study direction is highly related to the previous result by T. Elgindi, W. Hu, V. Šverák (2017). This is a joint work with Professors Kishimoto, Saiki and Yoneda.

- 71 三浦達彦 (東大数理) On singular limit equations for the Navier–Stokes equations in moving thin domains 10
 Tatsu-Hiko Miura (Univ. of Tokyo) On singular limit equations for the Navier–Stokes equations in moving thin domains

概要 We consider the incompressible Navier–Stokes equations in a three-dimensional moving thin domain. Under the assumption that the moving thin domain degenerates into a two-dimensional closed evolving surface as the width of the thin domain tends to zero, we give a formal derivation of limit equations on the degenerate evolving surface of the Navier–Stokes equations. We also compare our limit system with the Navier–Stokes equations on a stationary manifold, which is described in terms of the Levi–Civita connection.

- 72 前川 泰則 (京大理) On stability of physically reasonable solutions to the two-dimensional Navier–Stokes equations 10
 Yasunori Maekawa (Kyoto Univ.) On stability of physically reasonable solutions to the two-dimensional Navier–Stokes equations

概要 The flow past an obstacle is a fundamental object in fluid mechanics. In 1967 R. Finn and D. R. Smith proved the unique existence of stationary solutions, called the physically reasonable solutions, to the Navier–Stokes equations in a two-dimensional exterior domain modeling this type of flows when the Reynolds number is sufficiently small. In this talk we prove that the physically reasonable solutions constructed by Finn and Smith are asymptotically stable with respect to small and well-localized initial perturbations.

- 73 仲尾 光平 (信州大総合工) 3次元の領域における Navier–Stokes 方程式の Beale–Kato–Majda 型の
 谷内 靖 (信州大理) 解の接続定理 10
 Kohei Nakao (Shinshu Univ.) Beale–Kato–Majda type extension criterion of smooth solutions to the
 Yasushi Taniuchi (Shinshu Univ.) Navier–Stokes equations in three dimensional domains

概要 We shall establish a Beale–Kato–Majda type extension criterion of smooth solutions to the Navier–Stokes equations. It is known that if a smooth solution u to the Navier–Stokes equations on $(0, T)$ satisfies $\int_0^T \|\operatorname{rot} u(\tau)\|_{L^\infty} d\tau < \infty$, then u can be continued to the smooth solution on $(0, T')$ for some $T' > T$. In this talk, we shall slightly relax this condition for extension of smooth solutions to the 3D Navier–Stokes equations in not only the whole space but also the half space, bounded domains and exterior domains with smooth boundary.

- 74 小 菌 英雄 (早大基幹理工)* A remark on Liouville-type theorem for the nonstationary Navier–Stokes
 寺澤 祐高 (名大多元数理) equations in two dimensional domains 10
 若杉 勇太 (愛媛大理工)
 Hideo Kozono (Waseda Univ.) A remark on Liouville-type theorem for the nonstationary Navier–Stokes
 Yutaka Terasawa (Nagoya Univ.) equations in two dimensional domains
 Yuta Wakasugi (Ehime Univ.)

概要 We consider the initial-boundary value problem for the Navier–Stokes equations in two dimensional domains. Under a certain condition on the asymptotic behavior of the vorticity at infinity, we prove that the vorticity and its gradient of solutions are both globally square integrable. As their applications, Liouville-type theorems are obtained.

- 75 岡田 晃 (京大人間環境) 斉次 Besov 空間に初期値をとる Navier–Stokes 方程式の解の一意存在と
 小 菌 英雄 (早大理工) その解析性 10
 清水 扇丈 (京大人間環境)
 Akira Okada (Kyoto Univ.) Spatial analyticity of solutions to the Navier–Stokes equations with
 Hideo Kozono (Waseda Univ.) initial data in homogeneous Besov spaces
 Senjo Shimizu (Kyoto Univ.)

概要 There are mild solutions to the Navier–Stokes equations in Serrin class with initial data in scale invariant homogeneous Besov spaces. We show the solution is uniformly analytic in $x \in \mathbb{R}^n$ whose convergence radius is in proportion to \sqrt{t} .

- 76 岡部 考宏 (弘前大教育) Remark on the strong solvability of the Navier–Stokes equations in the weak L^n space 10
筒井 容平 (信州大理) Remark on the strong solvability of the Navier–Stokes equations in the weak L^n space
Takahiro Okabe (Hirosaki Univ.) Remark on the strong solvability of the Navier–Stokes equations in the weak L^n space
Youhei Tsutsui (Shinshu Univ.) Remark on the strong solvability of the Navier–Stokes equations in the weak L^n space

概要 We consider incompressible Navier–Stokes equations in the whole space \mathbb{R}^n under the non-trivial forces. In particular, we construct a strong solution to the Navier–Stokes equations in weak Lebesgue space. Firstly we introduce a maximal subspace where the Stokes semigroup is strongly continuous. Then we construct a local in time weak mild solution of the Navier–Stokes equations in $L^{n,\infty}$. Then uniqueness criterion is discussed.

- 77 鶴見 裕之 (早大理工) Solutions of the stationary Navier–Stokes equations in homogeneous Triebel–Lizorkin spaces 10
Hiroyuki Tsurumi (Waseda Univ.) Solutions of the stationary Navier–Stokes equations in homogeneous Triebel–Lizorkin spaces

概要 We consider the stationary Navier–Stokes equations in \mathbb{R}^n for $n \geq 3$. We show the existence and uniqueness of solutions in the homogeneous Triebel–Lizorkin space $\dot{F}_{p,q}^{-1+\frac{n}{p}}$ with $p \leq n$ for small external forces in $\dot{F}_{p,q}^{-3+\frac{n}{p}}$. These are shown by the boundedness of the Riesz transform, the para-product formula, and the embedding theorem in homogeneous Triebel–Lizorkin spaces. Moreover, it is proved that under some additional assumption on external forces, our solutions have more regularity.

- 78 鶴見 裕之 (早大理工) Ill-posedness of the stationary Navier–Stokes equations in homogeneous Besov spaces 10
Hiroyuki Tsurumi (Waseda Univ.) Ill-posedness of the stationary Navier–Stokes equations in homogeneous Besov spaces

概要 The solutions of the stationary Navier–Stokes equations in \mathbb{R}^n for $n \geq 3$ in the scaling invariant Besov spaces are investigated. It is proved that bounded smooth external forces whose $\dot{B}_{\infty,1}^{-3}$ norms are arbitrary small can produce bounded smooth solutions whose $\dot{B}_{\infty,\infty}^{-1}$ norms are arbitrary large. Such norm inflation phenomena are shown by constructing the sequence of external forces, as similar to those of initial data proposed by Bourgain–Pavlović in the non-stationary problem.

16:30~17:30 特別講演

岩 渕 司 (東北大理)^b 圧縮性 Navier–Stokes 方程式に対する不適切性について

Tsukasa Iwabuchi (Tohoku Univ.) On the ill-posedness for the compressible Navier–Stokes equations

概要 We consider the ill-posedness problem for the compressible Navier–Stokes system under the barotropic condition in the critical Besov spaces. It is known that the existence and the uniqueness of the solution hold in the homogeneous Besov spaces $\dot{B}_{p,1}^{\frac{n}{p}} \times \dot{B}_{p,1}^{\frac{n}{p}-1}$ with $1 \leq p < 2n$, where the density and the velocity belong to $\dot{B}_{p,1}^{\frac{n}{p}}$ and $\dot{B}_{p,1}^{\frac{n}{p}-1}$, respectively. On the other hand, if $p > 2n$, the solution does not depend on initial data continuously in general. In this talk, we show that for the critical case $p = 2n$ the system is ill-posed by showing the norm inflation.

実 函 数 論

3月20日(火) 第VII会場

9:00~12:00

- 1 川崎敏治 (日大工・玉川大工) Henstock–Kurzweil 主値積分について 15
 Toshiharu Kawasaki On the principal value of Henstock–Kurzweil integral
 (Nihon Univ./Tamagawa Univ.)

概要 There are the wide Denjoy integral, the approximately continuous Perron integral, the approximately continuous Henstock integral, the approximately continuous Denjoy integral and the distributional denjoy integral as the wider integrals than Denjoy–Perron–Henstock–Kurzweil integral. These integrals are defined by replacing derivative with approximately derivative and distributinal derivative. In this talk, we extend an integral by the Cauchy’s principal value and show the obtained results.

- 2 本田あおい (九工大情報工) 準劣加法的単調測度に関する弱 L_p 空間 $L^{p,\infty}$ 15
 岡崎悦明 (フuzzyシステム研)
 Aoi Honda (Kyushu Inst. of Tech.) Weak L_p space $L^{p,\infty}$ for quasi-subadditive monotone measure
 Yoshiaki Okazaki
 (Fuzzy Logic Systems Inst.)

概要 The weak L_p space $L^{p,\infty}(\mu)$ is introduced for the quasi-subadditive monotone measure μ . If μ is continuous from below, then $L^{p,\infty}(\mu)$ is a quasi-Banach space. As an application it is shown that there exists a real number $\alpha \in (0, 1]$ such that the power transform μ^α is uniformly quasi-subadditive. Furthermore there exists a subadditive monotone measure λ satisfying $\lambda \leq \mu^\alpha \leq 2\lambda$.

- 3 室伏俊明 (東工大情報理工) 非加法的測度論における強形の Egorov の定理の成立条件 15
 榎本直樹 (東工大情報理工)
 Toshiaki Murofushi (Tokyo Tech) Conditions for the strong form of the Egorov theorem in non-additive
 Naoki Enomoto (Tokyo Tech) measure theory

概要 The consequent of the strong form of the Egorov theorem in non-additive measure theory is that strong almost everywhere convergence implies strong almost uniform convergence. This paper shows that the conjunction of the uniform subadditive continuity and the order continuity of the non-additive measure is a sufficient condition for the consequent of the strong form of the Egorov theorem, and that the monotone continuity is a necessary condition.

- 4 高阪史明 (東海大理) 完備 CAT(1) 空間における凸関数に対する近接点法 15
 Fumiaki Kohsaka (Tokai Univ.) The proximal point algorithm for convex functions in complete CAT(1)
 spaces

概要 Using the recently introduced resolvent of a convex function in a complete CAT(1) space, we obtain existence and convergence theorems for the proximal point algorithm in such a space.

- 5 松下慎也 (秋田県立大) 作用素分割法の収束について 15
 Shin-ya Matsushita (Akita Pref. Univ.) On the convergence of an operator splitting method

概要 Let H be a real Hilbert space and let $f : H \rightarrow (-\infty, \infty]$ and $g : H \rightarrow (-\infty, \infty]$ be proper, lower semicontinuous and convex functions. This talk considers a problem of finding the resolvent $J_{\partial(f+g)}$ of the subdifferential $\partial(f+g)$. It is assumed that both the resolvents $J_{\partial f}$ and $J_{\partial g}$ of ∂f and ∂g can be easily computed. This enables us to consider the case in which a solution to the problem cannot be computed easily.

- 6 青 山 耕 治 (千葉大社会科学) 強擬非拡大写像について 15
Koji Aoyama (Chiba Univ.) Strongly quasi-nonexpansive mappings, II

概要 In this talk, we introduce and study a quasi-nonexpansive mapping, a strictly quasi-nonexpansive mapping, and a strongly quasi-nonexpansive mapping in an abstract space. In particular, we give some basic properties of such mappings.

- 7 厚 芝 幸 子 (山梨大教育) Weak and strong convergence theorems for a sequence of nonlinear operators 15
Sachiko Atsushiba Weak and strong convergence theorems for a sequence of nonlinear operators
(Univ. of Yamanashi)

概要 In this talk, we study the relations among k -acute points, attractive points and fixed points. Further, we apply these to rearrange proofs of some known convergence theorems and to prove new convergence theorems for nonlinear mappings in Hilbert spaces. Using the ideas of attractive points, acute points and fixed points, we also prove convergence theorems for nonlinear mappings in Banach spaces.

- 8 鈴 木 智 成 (九 工 大 工) ν -generalized metric space の 2 つ の 位 相 15
Tomonari Suzuki Two topologies on ν -generalized metric spaces
(Kyushu Inst. of Tech.)

概要 We will talk about two topologies on ν -generalized metric spaces.

- 9 富 澤 佑 季 乃 (新 潟 工 大 工) $\pi/2$ 回 転 不 変 ノ ル ム に よ る 幾 何 学 的 定 数 15
Yukino Tomizawa Geometric constants of $\pi/2$ -rotation invariant norms
(Niigata Inst. of Tech.)

概要 In this talk, we study the von Neumann–Jordan constant of $\pi/2$ -rotation invariant norms on R^2 . We know that any $\pi/2$ -rotation invariant normed space is isometrically isomorphic to some Day–James space. Since the von Neumann–Jordan constant is invariant under isometrically isomorphic, for characterization of the constant, it is enough to consider the Day–James space. From this fact, we can give some estimations of the constant.

- 10 国 定 亮 一 (早 大 教 育)^b 総 和 法 と \mathbb{R}^\times 上 の 調 和 解 析 15
Ryoichi Kunisada (Waseda Univ.) Summability methods and Fourier analysis on \mathbb{R}^\times

概要 We introduce a certain class of summability methods which are defined by the convolution operation in the group algebra $L^1(\mathbb{R}^\times)$ and study b -strongness and b -equivalence between them. In particular, this class contains an integral version of Cesàro summability method and we give a necessary and sufficient condition for a summability method in the class to equivalent to this one.

- 11 飯 田 毅 士 (福 島 工 高 専) The dual inequality of the boundedness for the Hardy–Littlewood maximal operator and the fractional integrals 15
Takeshi Iida The dual inequality of the boundedness for the Hardy–Littlewood maximal operator and the fractional integrals
(Fukushima Nat. Coll. of Tech.)

概要 In this talk, we consider the dual inequality for the Hardy–Littlewood maximal operator M and the fractional integral operator I_α . Since the fractional integral operator I_α has the property $\int I_\alpha f g dx = \int f I_\alpha g dx$ for $f \geq 0$ and $g \geq 0$, the weight norm inequality $I_\alpha : L^p(v^p) \rightarrow L^q(u^q)$ is equivalent to $I_\alpha : L^{q'}(u^{-q'}) \rightarrow L^{p'}(v^{-p'})$, where $\frac{1}{p} + \frac{1}{p'} = \frac{1}{q} + \frac{1}{q'} = 1$ ($1 < p < \infty$, $1 < q < \infty$) and let u and v be weights. Analogously, we consider whether the dual inequality for the Hardy–Littlewood maximal operator holds on weighted Lebesgue spaces. In particular, we verify the dual inequalities of $M : L^p(w) \rightarrow L^p(w)$ and $M : L^p(Mw) \rightarrow L^p(w)$.

14:15~16:15

- 12 新井龍太郎 (茨城大理) Compact commutators of Calderón–Zygmund and generalized fractional
中井英一 (茨城大理) integral operators with a function in Campanato spaces on generalized
Morrey spaces 15
- Ryutaro Arai (Ibaraki Univ.) Compact commutators of Calderón–Zygmund and generalized fractional
Eiichi Nakai (Ibaraki Univ.) integral operators with a function in Campanato spaces on generalized
Morrey spaces

概要 We discuss the compactness of the commutators $[b, T]$ and $[b, I_\rho]$ on generalized Morrey spaces with variable growth condition, where T is a Calderón–Zygmund operator, I_ρ is a generalized fractional integral operator and b is a function in generalized Campanato spaces with variable growth condition.

- 13 石明磊 (茨城大理工) Commutators of generalized fractional integral operators on Orlicz spaces
中井英一 (茨城大理) 15
- Minglei Shi (Ibaraki Univ.) Commutators of generalized fractional integral operators on Orlicz spaces
Eiichi Nakai (Ibaraki Univ.)

概要 Let \mathbb{R}^n be the n -dimensional Euclidean space. Let $b \in \text{BMO}(\mathbb{R}^n)$ and T be a Calderón–Zygmund singular integral operator. In 1976 Coifman, Rochberg and Weiss proved that the commutator $[b, T] = bT - Tb$ is bounded on $L^p(\mathbb{R}^n)$ ($1 < p < \infty$), that is,

$$\|[b, T]f\|_{L^p} = \|bTf - T(bf)\|_{L^p} \leq C\|b\|_{\text{BMO}}\|f\|_{L^p},$$

where C is a positive constant independent of b and f . For the fractional integral operator I_α , Chanillo proved the boundedness of $[b, I_\alpha]$ in 1982. These results were extended to Orlicz spaces by Fu, Yang and Yuan (2012, 2014). In this talk we discuss the boundedness of the commutator $[b, I_\rho]$ on Orlicz spaces, where I_ρ is a generalized fractional integral operator.

- 14 貞末岳 (大阪教育大数学教育) Commutators of fractional integrals on martingale Morrey spaces 15
中井英一 (茨城大理)
- Gaku Sadasue (Osaka Kyoiku Univ.) Commutators of fractional integrals on martingale Morrey spaces
Eiichi Nakai (Ibaraki Univ.)

概要 We study the boundedness of the commutator of fractional indegrals on martingale Morrey spaces. We give a necessary and sufficient condition on the boundedness in terms of martingale Campanato spaces.

- 15 岩瀨司 (東北大理)* Dirichlet Laplacian で生成される Besov 空間 15
谷口晃一 (中大理工)
松山登喜夫 (中大理工)
- Tsukasa Iwabuchi (Tohoku Univ.) Besov spaces generated by the Dirichlet Laplacian
Koichi Taniguchi (Chuo Univ.)
Tokio Matsuyama (Chuo Univ.)

概要 We define the Besov spaces on an arbitrary open set of \mathbb{R}^d . Based on the spectral theorem for the Dirichlet Laplacian, we introduce test function spaces and distributions to define the Besov spaces analogously to Peetre’s idea for the whole space case, which is by the dyadic decomposition of the spectrum. We will define the Besov spaces of the inhomogeneous type and the homogeneous type, and also show fundamental properties such as completeness, embedding, etc.

- 16 谷口晃一(中大理工) 領域上の Besov 空間における双線形評価式 15
 岩渕 司(東北大理)
 松山登喜夫(中大理工)
 Koichi Taniguchi (Chuo Univ.) Bilinear estimates in Besov spaces on domains
 Tsukasa Iwabuchi (Tohoku Univ.)
 Tokio Matsuyama (Chuo Univ.)

概要 The purpose of this talk is to establish bilinear estimates in Besov spaces generated by the Dirichlet Laplacian on a domain of Euclidian spaces. These estimates are proved by using the gradient estimates for heat semigroup together with the Bony paraproduct formula and the boundedness of spectral multipliers.

- 17 谷口晃一(中大理工) Schrödinger 作用素によって生成される Besov 空間 15
 岩渕 司(東北大理)
 松山登喜夫(中大理工)
 Koichi Taniguchi (Chuo Univ.) Besov spaces generated by Schrödinger operators
 Tsukasa Iwabuchi (Tohoku Univ.)
 Tokio Matsuyama (Chuo Univ.)

概要 We define the Besov spaces on an arbitrary open set of \mathbb{R}^d , based on the spectral theorem for Schrödinger operators with potential of which negative part is of the Kato class. The purpose of this talk is to show the isomorphism relations among the Besov spaces generated by the Dirichlet Laplacian and the Schrödinger operators.

- 18 加藤睦也(阪大理) A remark on the Schrödinger operator on Wiener amalgam spaces ... 15
 富田直人(阪大理)
 Tomoya Kato (Osaka Univ.) A remark on the Schrödinger operator on Wiener amalgam spaces
 Naohito Tomita (Osaka Univ.)

概要 In this talk, we study the boundedness of the Schrödinger operator $e^{i\Delta}$ on Wiener amalgam spaces and determine its optimal condition.

16:30~17:30 特別講演

- 田中亮太郎(九大数理) Geometric techniques in Banach space theory: Challenges to Tingley's problem
 Ryotaro Tanaka (Kyushu Univ.) Geometric techniques in Banach space theory: Challenges to Tingley's problem

概要 Mathematical developments are always based on problems. Challenges to difficulties generate new ideas. In this talk, we focus on a 30 years old open problem in Banach space theory, so-called Tingley's problem, and present new geometric techniques (and results) derived from our challenges.

3月21日(水) 第VII会場

9:15~12:00

- 19 水上雅昭(東京理大理) The parabolic-elliptic chemotaxis case as a limit of the fully parabolic chemotaxis system with signal-dependent sensitivity 15
 Masaaki Mizukami (Tokyo Univ. of Sci.) The parabolic-elliptic chemotaxis case as a limit of the fully parabolic chemotaxis system with signal-dependent sensitivity

概要 This paper gives a first insight into making a mathematical bridge between the parabolic-parabolic signal-dependent chemotaxis system and its parabolic-elliptic version. To be more precise, this talk deals with convergence of a solution for the parabolic-parabolic chemotaxis system with strong signal sensitivity to that for the parabolic-elliptic chemotaxis system.

- 20 来間俊介 (東京理大理) Vanishing viscosity for a Cahn–Hilliard type system on unbounded domains 15
 Shunsuke Kurima (Tokyo Univ. of Sci.) Vanishing viscosity for a Cahn–Hilliard type system on unbounded domains

概要 This talk deals with vanishing viscosity for a Cahn–Hilliard type system on an unbounded domain with smooth bounded boundary. Colli–Gilardi–Rocca–Sprekels (2017) studied it in the case of a bounded domain using Aubin–Lions lemma. However, this lemma dose not work well in the case of unbounded domains. The present work asserts that we can discuss vanishing viscosity for the above system in an unbounded domain.

- 21 喜多航佑 (早大理工) Some threshold property for a reaction diffusion system arising from a
 大谷光春 (早大理工) nuclear reactor model 15
 坂本浩紀
 (日立GEニュークリア・エナジー(株))
 Kosuke Kita (Waseda Univ.) Some threshold property for a reaction diffusion system arising from a
 Mitsuharu Ôtani (Waseda Univ.) nuclear reactor model
 Hiroki Sakamoto
 (Hitachi-GE Nuclear Energy, Ltd.)

概要 We study an initial-boundary value problem for a reaction diffusion system, which consists of two real-valued unknown functions. This system describes diffusion phenomena of neutrons and heat in nuclear reactors, introduced by Kastenberg and Chambré. In this model, the unknown functions represent the neutron density and the temperature in nuclear reactors. We proved that this equation has at least one positive stationary solution in last MSJ Autumn Meeting 2017. In this talk, we show that the positive stationary solution plays a role of threshold to classify initial data into two groups; corresponding solutions of the equation blow up in finite time and exist globally.

- 22 都築 寛 (広島修道大経済) Existence for Vlasov–Poisson equations with angle error in magnetic field in a half-space 15
 Yutaka Tsuzuki Existence for Vlasov–Poisson equations with angle error in magnetic field in a half-space
 (Hiroshima Shudo Univ.)

概要 We deal with initial-boundary problems for Vlasov–Poisson equations in a half-space with external magnetic force horizontal to a wall. In 2013, Skubachevskii gives local-in-time solvability to the system. Moreover, in 2017, global-in-time solutions were obtained by effectively using the magnetic force whose direction is horizontal to the wall. This talk provides an existence result for the system where the magnetic force has angle error in the vertical direction.

- 23 元田大志 (京都教育大) 力学的境界条件下における Cahn–Hilliard 系の時間周期解について 15
 Taishi Motoda (Kyoto Univ. of Edu.) Time periodic solutions of Cahn–Hilliard system with dynamic boundary conditions

概要 The existence problem for Cahn–Hilliard system with dynamic boundary conditions and time periodic conditions is discussed. We apply the abstract theory of evolution equations by using viscosity approach and the Schauder fixed point theorem in the level of approximate problem. One of the key point is the assumption for maximal monotone graphs with respect to their domains. Thanks to this, we obtain the existence result of the weak solution by using the passage to the limit.

- 24 中屋敷亮太 (千葉大理) 力学的境界条件を含む準線形型 Kobayashi–Warren–Carter システムの数学解析 15
 Ryota Nakayashiki (Chiba Univ.) Kobayashi–Warren–Carter system subject to quasi-linear diffusions and dynamic boundary conditions

概要 In this talk, we consider a coupled system of two parabolic type initial-boundary value problems, called the Kobayashi–Warren–Carter model of grain boundary motion in a polycrystal. The systems are denoted by $(S)_\varepsilon$ with arguments $\varepsilon \geq 0$. The characteristic point of our systems is to assume the dynamic boundary conditions in one problem. Now, the focus of this talk is to address the three assignment concerned with the qualitative results of the systems. The first is the existence of solutions to the systems, including the representation of solutions. The second is the continuous dependence of the systems to $(S)_\varepsilon$ for the variations of $\varepsilon \geq 0$. The third is the large-time behavior of solutions.

- 25 渡邊 紘 (大分大理工) ある退化放物型-放物型方程式系の可解性 15
 Hiroshi Watanabe (Oita Univ.) Solvability of degenerate parabolic-parabolic systems

概要 We consider the initial value problem (CP) for degenerate parabolic-parabolic systems with variable coefficients. The systems are coupled with strongly degenerate parabolic equations and nonhomogeneous heat equations. Strongly degenerate parabolic equations are regarded as a linear combination of the time-dependent conservation laws (quasilinear hyperbolic equations) and the porous medium type equations (nonlinear degenerate parabolic equations). Thus, the equation has both properties of hyperbolic equations and those of parabolic equations. In this talk, we discuss the solvability for (CP).

- 26 中村 誠 (山形大理) On the Cauchy problem of a semilinear diffusion equation in an expanding space 10
 佐藤 祐也 (山形大理) ing space 10
 Makoto Nakamura (Yamagata Univ.) On the Cauchy problem of a semilinear diffusion equation in an expanding space
 Yuya Sato (Yamagata Univ.) ing space

概要 We consider the Cauchy problem for a semilinear diffusion equation in an expanding space, and we show global solutions for small initial data.

- 27 中村 誠 (山形大理) Remarks on the derivation of Navier–Stokes equations and elastic wave equations in uniform and isotropic spacetimes 10
 Makoto Nakamura (Yamagata Univ.) Remarks on the derivation of Navier–Stokes equations and elastic wave equations in uniform and isotropic spacetimes

概要 We consider the extension of the Navier–Stokes equations and the elastic wave equations in the Minkowski spacetime to the equations in uniform and isotropic spacetimes.

- 28 儀我美一 (東大数理) 有界平均振動関数空間におけるストークス半群の解析性 10
 M. Bolkart
 (ダルムシュタット工科大)
 鈴木 拓也
 ((株)先端力学シミュレーション研)
 Yoshikazu Giga (Univ. of Tokyo) Analyticity of the Stokes semigroup in BMO
 Martin Bolkart (TU Darmstadt)
 Takuya Suzuki
 (Adv. Simulation Tech. of Mechanics R&D, Co., Ltd.)

概要 We consider the Stokes semigroup in a large class of domains including bounded domains, the half-space and exterior domains. We will prove that the Stokes semigroup is analytic in a certain type of solenoidal subspaces of BMO.

14:15~16:00

- 29 深尾 武 史 (京 都 教 育 大) Log 型ポテンシャルを持つ GMS モデルにおける純粋相からの分離定理
 15

Takeshi Fukao (Kyoto Univ. of Edu.) A strict separation property from pure phases for GMS model with logarithmic potential

概要 The equation and dynamic boundary condition of Cahn–Hilliard type was introduced by Goldstein–Miranville–Schimperna (2011), this problem is called GMS model and it is similar to the general Cahn–Hilliard system. In this talk, the double well potential of logarithmic type is employed. A strict separation property from pure phases is considered.

- 30 山 崎 教 昭 (神 奈 川 大 工) Quasi-variational evolution equations governed by double time-dependent
 剣 持 信 幸 (ワルシャワ大ICM) subdifferentials 15
 白 川 健 (千 葉 大 教 育)

Noriaki Yamazaki (Kanagawa Univ.) Quasi-variational evolution equations governed by double time-dependent
 Nobuyuki Kenmochi (Univ. of Warsaw) subdifferentials
 Ken Shirakawa (Chiba Univ.)

概要 In this talk we introduce a new class of doubly nonlinear quasi-variational evolution equation governed by double time-dependent subdifferentials. The main aim of this talk is to show the existence of a solution to our equations.

- 31 伊 藤 昭 夫 ^b Evolution inclusion on a real Hilbert space with quasi-variational structure for inner product —Time-dependent convex functions— 15

Akio Ito Evolution inclusion on a real Hilbert space with quasi-variational structure for inner product —Time-dependent convex functions—

概要 Throughout our recent researches, we showed that the tumor invasion model with quasi-variational structure can be rewritten into an evolution inclusion on a suitable Hilbert space. Moreover, its norms has a quasi-variational structure in general. That is, they depend upon an unknown function, which is one of the functions (components) of a unique solution to the tumor invasion model. In this talk, an evolution inclusion on a real Hilbert space with quasi-variational structure for inner products is considered. And a main purpose is to give the continuity property of proper l.s.c. convex functions with quasi-variational structure which appear as subdifferentials operators in the evolution inclusions.

- 32 白 川 健 (千 葉 大 教 育) 非等方性を考慮する画像処理問題の支配エネルギーに対する連続系勾配流
 15

Ken Shirakawa (Chiba Univ.) A gradient system based on an anisotropic image processing

概要 In this talk, we consider a gradient flow of a non-convex functional, which was proposed by [Berkels et al, pp. 293–301, Vision Modeling and Visualization 2006 (2006)] as a possible governing energy for an anisotropic image processing on a bounded spatial domain $\Omega \subset \mathbb{R}^2$. Our gradient flow is described in a nonstandard form of *partial differential inclusions*, which contains a composition $\partial\gamma \circ R$ of: a (possibly) set-valued subdifferential $\partial\gamma$ of an anisotropic metric $\gamma \in W^{1,\infty}(\mathbb{R}^2)$; and a rotation matrix $R \in C^\infty(\mathbb{R}; \mathbb{R}^{2 \times 2})$. Under appropriate settings, some mathematical observations for the gradient system will be provided on the basis of the time-discretization approach.

- 33 熊崎 耕太 (苫小牧工高専) 多孔質媒体内で起こる水分膨張過程を表すある自由境界問題の可解性について 15
 Kota Kumazaki (Tomakomai Nat. Coll. of Tech.) Solvability of a free boundary problem describing moisture swelling process in porous materials

概要 In this talk, we propose a mathematical model for moisture swelling process in concrete materials. Moisture swelling process appear in, for instance, frost damage in concrete materials which is a nonlinear phenomenon to give rise to crack inside of concrete. Our model consists of a diffusion equation for moisture in a one microscopic hole of concrete and a free boundary problem for the front of the moisture region. In this talk, we discuss the existence and uniqueness of a time global solution for this problem, and moreover, some results of the behavior of the free boundary as time goes to infinity.

- 34 愛木 豊彦 (日本女大理) 水分吸着過程を記述する自由境界問題の弱解の存在 15
 Toyohiko Aiki (Japan Women's Univ.) Existence of a weak solution to a free boundary problem describing adsorption process

概要 Recently, we investigated a free boundary problem describing a adsorption process in a porous media. In the global existence result to this model we require the smallness condition $h < 1$ for the boundary function h to prevent that the free boundary touches the fixed boundary. This is a big obstacle to consider a multi-scale model consisting of nonlinear diffusion equation and the free boundary problem. To overcome this difficulty, we propose a new weak formulation to the free boundary problem. Also, we show existence of a weak solution without the smallness condition to the boundary data.

16:15~17:15 特別講演

- 内田 俊 (早大理工) 多孔質媒質中の二重拡散対流現象を記述する方程式系の可解性について
 Shun Uchida (Waseda Univ.) Solvability of a system describing double-diffusive convection phenomena in some porous medium

概要 We consider some equations describing double-diffusive convection phenomena of incompressible viscous fluid in a porous medium. Roughly speaking, this system consists of the Stokes equation and two advection-diffusion equations. Although the fluid equation in our model is linearized, the others still possess convection terms as non-monotone perturbations, which make it difficult to deal with this system. Main topic of this talk is to show the global solvability of this double-diffusive convection system. In particular, we focus on the existence of time periodic solutions to the system in the whole space domain for large data, i.e., without any smallness conditions for given external forces. In previous results for periodic problems of parabolic type equations with non-monotone perturbation terms (e.g., incompressible Navier–Stokes equations and Boussinesq system), it seems that either of the smallness of given data or the boundedness of space domain is essential. However, in spite of the presence of non-monotone terms, the solvability of our problem in the whole space is shown for large external forces via the convergence of solutions to approximate equations in bounded domains.

函数解析学

3月18日(日) 第IV会場

14:15~16:15

- 1 齋藤三郎 (群馬大*・再生核研)* What are reproducing kernels? 15
 松浦勉 (群馬大工)
 澤野嘉宏 (首都大東京理工)
Saburo Saitoh What are reproducing kernels?
 (Gunma Univ.*/Inst. of Reproducing Kernels)
 Tsutomu Matsuura (Gunma Univ.)
 Yoshihiro Sawano (Tokyo Metro. Univ.)

概要 Here, we will state simply a general meaning for reproducing kernels. We would like to answer for the general and essential question that: what are reproducing kernels? By considering the basic problem, we were able to obtain a general concept of the generalized delta function as a generalized reproducing kernel and, as a general reproducing kernel Hilbert space, we can consider all separable Hilbert spaces comprising functions.

- 2 齋藤三郎 (群馬大*・再生核研)* General integral transforms by the concept of generalized reproducing
 松浦勉 (群馬大工) kernels 15
 澤野嘉宏 (首都大東京理工)
Saburo Saitoh General integral transforms by the concept of generalized reproducing
 (Gunma Univ.*/Inst. of Reproducing Kernels) kernels
 Tsutomu Matsuura (Gunma Univ.)
 Yoshihiro Sawano (Tokyo Metro. Univ.)

概要 The general integral transforms in the framework of Hilbert spaces were combined with the general theory of reproducing kernels and many applications were developed. The basic assumption here that the integral kernels belong to some Hilbert spaces. However, as a very typical integral transform, in the case of Fourier integral transform, the integral kernel does not belong to $L_2(\mathbf{R})$, however, we can establish the isometric identity and inversion formula.

On the above situations, we will develop some general integral transform theory containing the Fourier integral transform case that the integral kernel does not belong to any Hilbert space, based on the general concept of generalized reproducing kernels.

- 3 青木貴史 (近畿大理工) Linear continuous operators acting on the space of entire functions of a
 石村隆一 (千葉大理) given order 15
 D. C. Struppa (Chapman Univ.)
内田匠風 (近畿大総合理工)
 Takashi Aoki (Kindai Univ.) Linear continuous operators acting on the space of entire functions of a
 Ryuichi Ishimura (Chiba Univ.) given order
 Daniele C. Struppa (Chapman Univ.)
Shofu Uchida (Kindai Univ.)

概要 We consider the relationship between linear continuous operators acting on the space of entire functions of one variable of a given order and linear differential operators of infinite order satisfying certain growth conditions for the coefficients. We found that these two classes of operators are equivalent. Our results can be extended to the case of several variables.

- 4 岩田 順 敬 (東工大科学技術創成研究院) 無限小生成作用素の対数表現に基づいた Banach 代数上の加群 15

Yoritaka Iwata (Tokyo Tech) Module over the Banach algebra defined by the logarithmic representation of infinitesimal generators

概要 The operator algebra is introduced based on the framework of logarithmic representation of infinitesimal generators. In conclusion a set of generally-unbounded infinitesimal generators is characterized as a module over the Banach algebra.

- 5 廣島 文生 (九大数理) Mass renormalization in the Nelson model 12
大澤 進 (九大数理)

Fumio Hiroshima (Kyushu Univ.) Mass renormalization in the Nelson model
Susumu Osawa (Kyushu Univ.)

概要 The asymptotic behavior of the effective mass $m_{\text{eff}}(\Lambda)$ of the so-called Nelson model in quantum field theory is considered, where Λ is an ultraviolet cutoff parameter of the model. Let m be the bare mass of the model. It is shown that for sufficiently small coupling constant $|\alpha|$ of the model, $m_{\text{eff}}(\Lambda)/m$ can be expanded as $m_{\text{eff}}(\Lambda)/m = 1 + \sum_{n=1}^{\infty} a_n(\Lambda)\alpha^{2n}$. A physical folklore is that $a_n(\Lambda) = O([\log \Lambda]^{(n-1)})(\Lambda \rightarrow \infty)$. It is rigorously shown that

$$0 < \lim_{\Lambda \rightarrow \infty} a_1(\Lambda) < C, \quad C_1 \leq \lim_{\Lambda \rightarrow \infty} a_2(\Lambda)/\log \Lambda \leq C_2$$

with some constants C , C_1 and C_2 .

- 6 廣島 文生 (九大数理) Renormalized Gibbs measures associated with the Nelson model 15
Fumio Hiroshima (Kyushu Univ.) Renormalized Gibbs measures associated with the Nelson model

概要 We can construct the Gibbs measure associated with the renormalized Nelson model in scalar quantum field theory. By using this Gibbs measure we investigate properties of the ground state of the Nelson Hamiltonian.

- 7 片岡 清 臣 (東大数理) 蔵本予想に関連する作用素の一般化固有値について 15
馬田 優 (東大数理)

Kiyoomi Kataoka (Univ. of Tokyo) On generalized eigenvalues of an operator related to Kuramoto conjecture
Yu Mada (Univ. of Tokyo)

概要 Kuramoto's famous conjecture is that there is a positive constant K_c such that the system of oscillators becomes synchronized only when the coupling constant $K > K_c$. H. Chiba proved this conjecture for the continuous version of Kuramoto model in 2015. His proof covers the cases that the initial distributions $g(\omega)$ of frequencies of oscillators are Gaussian or Cauchy. His key tool is the precise analysis of the generalized eigenvalues of some unbounded linear operator T related to $g(\omega)$ on some Hilbert space. We extended his method to the analytic distributions of form $g = e^{-P(\omega)}/M$, where $P(\omega)$ is an even polynomial which increases in $\omega > 0$, and M is a positive constant.

16:30~17:30 特別講演

尾畑 伸明 (東北大情報) グラフのスペクトル解析における量子確率論の手法
 Nobuaki Obata (Tohoku Univ.) Quantum probabilistic methods for spectral analysis of graphs

概要 Quantum (non-commutative) probability, tracing back to von Neumann who originally aimed at statistical study of quantum mechanics, has been penetrated into other branches of mathematics. In this lecture we focus on the interaction with spectral analysis of (growing) graphs developed during the last decade and mention some current topics.

In general, quantum probability is discussed in terms of an algebraic probability space (\mathcal{A}, φ) , where \mathcal{A} is a unital $*$ -algebra and $\varphi : \mathcal{A} \rightarrow \mathbb{C}$ is a state. Given a graph we consider the adjacency matrix (or other matrices such as Laplacian matrix, distance matrix, etc.) as a real random variable of an algebraic probability space, typically the adjacency algebra equipped with a state.

The first method is quantum decomposition. If the adjacency matrix A is decomposed into a sum of three operators $A = A^+ + A^- + A^\circ$, where A^+, A^-, A° are the creation, annihilation and preservation operators in an interacting Fock space, then the spectral distribution of A is obtained as the vacuum distribution of the canonical field operator of the interacting Fock space, where the theory of orthogonal polynomials is applied. This method is useful also for asymptotic spectral distributions of growing graphs.

The second method is based on various concepts of independence arising from non-commutative nature. If the adjacency matrix is expressible as a sum of independent random variables, typically when a graph admits a product structure, the spectral distribution is given by convolution of probability distributions. Hence for a growing graph, the asymptotic spectral distribution is obtained from the associated central limit theorem. The Cartesian, comb, star, free, Kronecker, lexicographic products of graphs are discussed along with quantum probability.

3月19日(月) 第IV会場

9:30~12:00

- 8 渡辺 純成 (東京学大教育)* 翻訳書の計量言語学における函数解析的視点 15
 Junsei Watanabe Functional analytic viewpoints in computational linguistics of translated
 (Tokyo Gakugei Univ.) literary works

概要 Statistical analysis of styles of translated versions of literary works is more difficult than that of the originals, because most of the variables used in the latter case indicate only characteristics of the authors, not of the translators. We replace probability measures related with the text of a literary work by conditional expectations related with the text of the original and translated versions of the work, and compare the numerical values of these conditional expectations. Results obtained by applying our framework to Mancu books on natural sciences, Neo-Confucianism or Catholicism published in the 17th and the 18th centuries are given.

- 9 宮島 静雄 (東京理大理) Characterization of closed balls via metric projections 10
 齊藤 功 (東京理大理)
 Shizuo Miyajima (Tokyo Univ. of Sci.) Characterization of closed balls via metric projections
 Isao Saito (Tokyo Univ. of Sci.)

概要 Consider the following property (P) for a bounded closed convex set C in a Banach space X : (P) For every $x \in X$, a positive-scalar multiple of x gives a nearest point in C to x . Then it is clear that a closed ball with its center at the origin has this property. The converse to this assertion is the subject of this talk, and it is proved that a bounded closed convex set $C \subset X$ with $0 \in \text{Int } C$ possessing property (P) is a closed ball with center 0, provided X is smooth and $\dim X > 1$. It is also proved that if a closed convex set $C \subset X$ with $0 \in \text{Int } C$ satisfies (P) , then X is smooth, provided the boundary of C is smooth in a weak sense.

- 10 高橋 眞映 (山形大*) 掛け算や足し算によって分配される実数上のある種の半群演算の決定 .. 15
 高木 啓行 (信州大理)
 三浦 毅 (新潟大理)
 岡 裕和 (茨城大工)
 Sin-Ei Takahasi (Yamagata Univ.*) Semigroup operations distributed by the ordinary multiplication or addition on the real numbers
 Hiroyuki Takagi (Shinshu Univ.)
 Takeshi Miura (Niigata Univ.)
 Hirokazu Oka (Ibaraki Univ.)

概要 Our motivation of this research is based on a certain refinement of an inequality that is induced by an abstract Jensen's inequality and on a characterization of a semigroup operations that is distributed by a vector addition in a plane. We determine cancellative, continuous operations on \mathbb{R} with the usual topology that is distributed by the usual multiplication and addition.

- 11 羽鳥 理 (新潟大自然) Surjective isometries on Banach algebras of Lipschitz maps 15
 Osamu Hatori (Niigata Univ.) Surjective isometries on Banach algebras of Lipschitz maps

概要 We show that a surjective isometry with respect to the sum norm between the Banach algebras of Lipschitz maps with the values in unital commutative C^* -algebras is canonical. When the unital commutative C^* -algebra is the complex plane, the result confirms the statement of Example 8 of the paper by Jarosz and Pathak.

- 12 示野 信一 (関西学院大理工) A 型の超幾何関数と Lauricella の超幾何級数 15
 玉岡 優一 (関西学院大理工)
 Nobukazu Shimeno (Kwansei Gakuin Univ.) The hypergeometric function of type A and the Lauricella hypergeometric series
 Yuichi Tamaoka (Kwansei Gakuin Univ.)

概要 Heckman–Opdam hypergeometric function for the root system of type A_{n-1} with a certain degenerate parameter can be expressed by the Lauricella hypergeometric function F_D .

- 13 織田 寛 (拓殖大工) ファイン K -タイプの球関数 (1 階不変微分作用素がある場合) 15
 示野 信一 (関西学院大理工)
 Hiroshi Oda (Takushoku Univ.) Spherical functions for fine K -types
 Nobukazu Shimeno
 (Kwansei Gakuin Univ.)

概要 For a real split Lie group $G = KAN$, Vogan introduced the notion of *fine* K -types. We define a similar class of K -types for general connected real semisimple Lie groups. For a K -type in this class, we study matrix-valued elementary spherical functions. If there exists an invariant differential operator of the first order acting on the spherical functions, we can explicitly write the elementary spherical functions using Opdam's non-symmetric hypergeometric functions.

- 14 笹木 集夢 (東海大理) A Cartan decomposition for spherical homogeneous spaces of reductive type 15
 Atsumu Sasaki (Tokai Univ.) A Cartan decomposition for spherical homogeneous spaces of reductive type

概要 In this talk, we announce the result on a generalization of Cartan decomposition for symmetric spaces to spherical homogeneous spaces of reductive type. Moreover, we deal with some examples concerning to our result and explain how to find abelian subgroups.

- 15 田内 大渡 (東大数理)^b Multiplicity of a degenerate principal series for homogeneous spaces with infinite orbits 15
 Taito Tauchi (Univ. of Tokyo) Multiplicity of a degenerate principal series for homogeneous spaces with infinite orbits

概要 Let X be a homogeneous space of a real reductive Lie group G . Then it is proved by T. Kobayashi and T. Oshima that the regular representation $C^\infty(X)$ contains each irreducible representation of G at most finitely many times if a minimal parabolic subgroup P of G has an open orbit on X , or equivalently, if the number of P -orbits on X is finite. Moreover, Kobayashi proved that for a general parabolic subgroup Q of G , there is a degenerate principal series representation induced from Q contained in $C^\infty(X)$ with infinite multiplicity if Q has no open orbit on X . In this article, we prove that there is a degenerate principal series representation induced from Q contained in $C^\infty(X)$ with infinite multiplicity if the number of orientable (or transverse orientable) Q -orbits on X is infinite even when there exists an open Q -orbit on X .

- 16 伊藤 稔 (鹿児島大理) トレース付き wreath 代数を用いたある不変式環の記述 15
 Minoru Itoh (Kagoshima Univ.) A description of an invariant theory using the notion of wreath algebra with trace

概要 We introduce the notion of wreath algebra with trace. This can be regarded as an abstraction of the wreath product of an algebra with trace and the infinite symmetric group. Using this notion, we describe an invariant theory simply.

13:15~14:15 特別講演

伊 師 英 之 凸錐上の Γ 型積分
 (名大多元数理・JST さきがけ)
 Hideyuki Ishi Gamma-type integrals over convex cones
 (Nagoya Univ./JST PRESTO)

概要 The Laplace transform of a power function on the positive half line is equal to a power function again up to a constant multiple, where the Gamma function appears as the coefficient. Replacing the half line by a homogeneous convex cone, Gindikin showed that the Laplace transform of a relatively invariant function over the homogeneous cone equals a relatively invariant function on the dual cone with an explicit Gamma factor. On the other hand, it is known in mathematical statistics that the Laplace transform of a power of the determinant function over the cone of positive definite real symmetric matrices with prescribed zero components equals a product of powers of minors with an explicit Gamma factor provided that the zero pattern is associated to a chordal graph. Although such a cone is not necessarily homogeneous, the integral formula is quite similar to Gindikin's formula eventually.

In this talk, we introduce a new cone consisting of positive definite real symmetric matrices with a specific block decompositions satisfying certain axioms. The class of this new cone contains all the homogeneous cones and the cones associated to chordal graphs. We establish Gamma-type integral formulas concerning the Laplace transform of a product of powers of minors over the new cones, so that the formulas mentioned above are obtained as special cases. Furthermore, considering the analytic continuation of the Gamma integral formula, we define the Riesz distribution whose Laplace transform equals a product of powers of minors of symmetric matrices, which gives a fundamental solution of a certain differential operator in a special case.

3月20日(火) 第IV会場

9:30~12:00

- 17 綿谷安男(九大数理)* テント写像による自己相似写像系の次元群の計算 15
 梶原毅(岡山大環境理工)
 Yasuo Watatani (Kyushu Univ.) A computation of the dimension group for the self-similar map given by
 Tsuyoshi Kajiwara (Okayama Univ.) the tent map

概要 We introduce a dimension group for a self-similar map as the K_0 -group of the core of the C^* -algebra associated with the self-similar map together with the canonical endomorphism. The key step for the computation is an explicit description of the core as the inductive limit using their matrix representations over the coefficient algebra, which can be described explicitly by the singularity structure of branched points. We compute that the dimension group for the tent map is the countably generated free abelian group together with the unilatral shift.

- 18 松本健吾(上越教育大)* スメール空間からできる Ruelle C^* -環と両側 Cuntz-Krieger 環について
 15
 Kengo Matsumoto Ruelle C^* -algebras associated with Smale spaces and bilateral Cuntz-Krieger algebras
 (Joetsu Univ. of Edu.)

概要 We will characterize topologically conjugate two-sided topological Markov shifts $(\bar{X}_A, \bar{\sigma}_A)$ in terms of the associated asymptotic Ruelle C^* -algebras \mathcal{R}_A with its commutative C^* -subalgebras $C(\bar{X}_A)$ and the canonical circle actions. We will also show that extended Ruelle algebras $\tilde{\mathcal{R}}_A$, which are purely infinite version of the asymptotic Ruelle algebras, with its C^* -subalgebras $C(\bar{X}_A)$ and the canonical torus actions γ^A are complete invariants for topological conjugacy of two-sided topological Markov shifts.

- 19 大坂博幸 (立命館大理工) Stable rank for crossed products by actions of finite groups on C^* -algebras 15
 Hiroyuki Osaka (Ritsumeikan Univ.) Stable rank for crossed products by actions of finite groups on C^* -algebras

概要 Let G be a finite group, A a unital separable finite simple nuclear C^* -algebra, and α an action of G on A . Assume that A absorbs the Jiang–Su algebra \mathcal{Z} , the extremal boundary of the trace space of A is compact and finite dimensional and that α fixes any tracial state of A . Then $\text{tsr}(A \rtimes_{\alpha} G) = 1$. In particular, when A has a unique tracial state, we conclude $\text{tsr}(A \rtimes_{\alpha} G) = 1$ without above conditions on a tracial state space of A .

- 20 長谷川慧 (九大数理) 自由積 C^* 環の境界剛性 15
 Kei Hasegawa (Kyushu Univ.) Boundary rigidity for free product C^* -algebras

概要 We prove a boundary rigidity result for the embedding of a reduced free product C^* -algebra into its associated “crossed product” C^* -algebra. This provides new examples of rigid embeddings of exact C^* -algebras into purely infinite simple nuclear C^* -algebras.

- 21 澤田友佑 (名大多元数理) 自己双対 CAR 環上の準自由状態のアフィン性 15
 Yusuke Sawada (Nagoya Univ.) The affine property of quasi-free states on self-dual CAR algebras

概要 There is a one-to-one correspondence between quasi-free states on a self-dual CAR algebra and covariance operators. The problem of when the correspondence preserve convex combinations is solved in the case when the Hilbert space which we treat has a finite dimension and covariance operators commutes.

- 22 武石拓也 (京大数理研) Reconstructing the Bost–Connes semigroup actions from K-theory ... 15
 Takuya Takeishi (Kyoto Univ.) Reconstructing the Bost–Connes semigroup actions from K-theory

概要 We discuss on the classification of Bost–Connes systems. We present that two Bost–Connes C^* -algebras for number fields are isomorphic if and only if the original semigroups actions are conjugate. Together with recent reconstruction results in number theory by Cornelissen–de Smit–Li–Marcolli–Smit, we conclude that two Bost–Connes C^* -algebras are isomorphic if and only if the original number fields are isomorphic. This is a joint work with Y. Kubota.

- 23 増田俊彦 (九大数理) Tannaka–Kreĭn–Woronowicz duality from the viewpoint of \mathbb{Q} -systems 15
 Toshihiko Masuda (Kyushu Univ.) Tannaka–Kreĭn–Woronowicz duality from the viewpoint of \mathbb{Q} -systems

概要 We discuss the Tannaka–Kreĭn duality theorem of Woronowicz from the viewpoint of \mathbb{Q} -systems in the sense of Fidaleo–Isola.

- 24 磯野優介 (京大数理研) 無限テンソル積因子環に対する一意テンソル分解定理について 15
 Yusuke Isono (Kyoto Univ.) Unique prime factorization for infinite tensor product factors

概要 In this talk, we study a unique prime factorization property for tensor product factors with infinitely many tensor components. We provide several examples of type II and III factors which satisfy this property, including all free product factors with diffuse free product components. Our proof is based on Popa’s intertwining techniques and the study of relative amenability on the continuous cores.

- 25 小沢登高 (京大数理研) Finite-dimensional representations constructed from random walks ... 15
Narutaka Ozawa (Kyoto Univ.) Finite-dimensional representations constructed from random walks

概要 Let a finitely generated amenable group G and a probability measure μ on it (that is finitely-supported, symmetric, and non-degenerate) be given. I will present a construction, via the μ -random walk on G , of a harmonic cocycle and the associated orthogonal representation of G . Then I describe when the constructed orthogonal representation contains a non-trivial finite-dimensional subrepresentation (and hence an infinite virtually abelian quotient), and some sufficient conditions for G to satisfy Shalom's property H_{FD} . (joint work with A. Erschler, arXiv:1609.08585)

14:15~16:00

- 26 渚 勝 (千葉大理) いくつかの作用素ノルム不等式 15
Masaru Nagisa (Chiba Univ.) Some operator norm inequalities

概要 We prove some formulas for operator norm inequalities related to operator means. As an example, we can get the following inequality (proved by H. Kosaki in 2014): For any positive integer $n \geq m$, it holds

$$\frac{1}{2^n} \left\| \sum_{i=0}^n {}_n C_i H^{i/n} X K^{(n-i)/n} \right\| \leq \frac{1}{2^m} \left\| \sum_{j=0}^m {}_m C_j H^{j/m} X K^{(m-j)/m} \right\|,$$

where $\|\cdot\|$ is an arbitrary unitarily invariant norm on $\mathbb{M}_N(\mathbb{C})$, $H, K, X \in \mathbb{M}_N(\mathbb{C})$ and $H, K \geq 0$.

- 27 藤井淳一 (大阪教育大) Hopf代数での弦表現の導入 15
Junichi Fujii (Osaka Kyoiku Univ.) Graphical approach to Hopf algebras

概要 Hopf algebra is one of a important structure to consider objects in the quantum information theory. But the Sweedler's convention, which is a basic tool of the calculation, is a little complicated for beginners. So, as another tool, we observe the graphical calculus of Hopf algebras. Though it is often used partially, we extend it for instance to the quantum double or universal R-matrix.

- 28 藤本将行 (大阪教育大) Mixed Schwarz inequalities via the matrix geometric mean 10
瀬尾祐貴 (大阪教育大)
Masayuki Fujimoto Mixed Schwarz inequalities via the matrix geometric mean
(Osaka Kyoiku Univ.)
Yuki Seo (Osaka Kyoiku Univ.)

概要 Furuta showed a weighted version of a mixed Schwarz inequality for any square matrices. In this talk, we show the following matrix version based on the Cauchy-Schwarz inequality for matrices: Let A, X and Y be matrices in \mathbb{M}_n and $U \in \mathbb{M}_n$ a unitary matrix in a polar decomposition of $Y^*AX = U|Y^*AX|$. Then

$$|Y^*AX| \leq X^*|A|^{2\alpha} X \# U^*Y^*|A^*|^{2\beta} YU$$

holds for all $\alpha, \beta \in [0, 1]$ with $\alpha + \beta = 1$.

- 29 瀬尾祐貴 (大阪教育大教育) 作用素幾何平均を用いた加重混合シュワルツ不等式 10
藤本将行 (大阪教育大教育)
Yuki Seo (Osaka Kyoiku Univ.) A weighted mixed Schwarz operator inequality via the geometric operator mean
Masayuki Fujimoto
(Osaka Kyoiku Univ.)

概要 In this talk, by virtue of the Cauchy-Schwarz operator inequality due to J. I. Fujii, we show a weighted mixed Schwarz operator inequality in terms of the geometric operator mean. As applications, we show Wielandt type operator inequalities via the geometric operator mean.

- 30 山崎 丈明 (東洋大理工) The Karcher equation, relative operator entropy and the Ando–Hiai inequality 15
 Takeaki Yamazaki (Toyo Univ.) The Karcher equation, relative operator entropy and the Ando–Hiai inequality

概要 Pálfi has been obtained that the Generalized Karcher Equation (GKE) has a unique positive solution, and he obtained a lot of nice properties of it as an operator mean. In this talk, we shall introduce relations among a solution of the Generalized Karcher Equation (GKE), representing function of an operator mean and relative operator entropy, firstly. Next, we shall introduce further extensions of the Ando–Hiai inequality.

- 31 和田 州平 (木更津工高専) 安藤–日合不等式が成り立つための条件 10
 Shuhei Wada When does Ando–Hiai inequality hold?
 (Nat. Inst. of Tech., Kisarazu Coll.)

概要 Let α be in $(0, 1)$ and $r > 0$ and $\#_\alpha$ stand for the weighted operator geometric mean. We consider the following statement:

$$A, B > 0, \quad A \#_\alpha B \geq I \Rightarrow A^r \#_\alpha B^r \geq I.$$

Ando and Hiai show that if $r \geq 1$, then this holds. In the present paper, we prove that the above statement holds only if $r \geq 1$. We try to find a characterization of the indices $p, q \in (0, 1)$ and $\mu, \lambda > 0$ satisfying

$$A, B > 0, \quad A \#_p B \geq I \Rightarrow A^\mu \#_q B^\lambda \geq I.$$

- 32 遠山 宏明 (前橋工科大) Some relations among the n -th relative operator entropies and the n -th
 伊佐 浩史 (前橋工科大) operator divergences 15
 亀井 栄三郎
 渡邊 雅之 (前橋工科大)
 Hiroaki Tohyama Some relations among the n -th relative operator entropies and the n -th
 (Maebashi Inst. of Tech.) operator divergences
 Hiroshi Isa (Maebashi Inst. of Tech.)
 Eizaburou Kamei
 Masayuki Watanabe
 (Maebashi Inst. of Tech.)

概要 Let A and B be bounded positive invertible operators on a Hilbert space \mathcal{H} . For each $n \in \mathbb{N}$, let $\Psi_{A,B}^{[1]}(x, y) \equiv \frac{\Psi_{A,B}(x) - \Psi_{A,B}(y)}{x - y}$ and $\Psi_{A,B}^{[n]}(x, y) \equiv \frac{\Psi_{A,B}^{[n-1]}(x, y) - \Psi_{A,B}^{[n-1]}(y, y)}{x - y}$ ($n \geq 2$), where $\Psi_{A,B}(t) \equiv A^{\frac{1}{2}}(A^{-\frac{1}{2}}BA^{-\frac{1}{2}})^t A^{\frac{1}{2}}$. So far, we have introduced the notions of the n -th operator divergences $D_{FK}^{[n]}(A|B)$ and $\Delta_1^{[n]}(x)$ which are generalizations of the Petz–Bregman divergence $D_{FK}(A|B) \equiv B - A - S(A|B)$ and $\Delta_1 \equiv T_x(A|B) - S(A|B)$, respectively. In this talk, we introduce the notions of the n -th relative operator entropies $T_x^{[n]}(A|B) \equiv \Psi_{A,B}^{[n]}(x, 0)$ and $S^{[n]}(A|B) \equiv \Psi_{A,B}^{[n]}(0, 0)$, and show some relations among the n -th relative operator entropies and the n -th operator divergences.

16:15~17:15 特別講演

日 合 文 雄 (東 北 大 情 報) 多変数の行列・作用素平均

Fumio Hiai (Tohoku Univ.) Multivariate matrix/operator means

概要 The notion of (two-variable) operator means was introduced in an axiomatic way by Kubo and Ando (1980). A long-standing problem since then had been to generalize it to more than two variables. A breakthrough happened when the definitions of multivariate geometric means of positive definite matrices were found by the iteration method of Ando, Li and Mathias (2004) and in the Riemannian geometry approach by Moakher (2005) and by Bhatia and Holbrook (2006). In this talk I survey recent developments on multivariate means of positive definite matrices/operators in the Riemannian geometry approach, mainly on the multivariate version of the weighted geometric and power means. A significant feature of the multivariate geometric mean is that it is understood as the Cartan barycenter in the Riemannian trace metric, so it is also characterized by the Karcher equation (the gradient zero equation). This approach using Karcher type equations can be extended to an even more general setting of probability measures on the positive definite matrices (also operators), where the Wasserstein distance plays a crucial role. I furthermore explain log-majorization for the geometric mean in the setting of probability measures and a recent result on deformation of multivariate operator means by a fixed point method.

統計数学

3月18日(日) 第VIII会場

9:15~12:00

- 1 山崎和俊 (関西大システム理工) Fluctuation theory for level-dependent Lévy processes 15
 I. Czarna (Univ. of Wrocław)
 J.-L. Perez
 (Centro de Investigación en Matemáticas)
 T. Rolski (Univ. of Wrocław)
- Kazutoshi Yamazaki (Kansai Univ.) Fluctuation theory for level-dependent Lévy processes
 Irmina Czarna (Univ. of Wrocław)
 Jose-Luis Perez
 (Centro de Investigación en Matemáticas)
 Tomasz Rolski (Univ. of Wrocław)

概要 We consider a level-dependent Lévy process that changes its drift depending on the position of the process. We first generalize the refracted Lévy process of Kyprianou and Loeffen (2010), which changes its drift above a given threshold, to the multi-refracted case. We then extend the results for more general level-dependent Lévy processes. We show how fluctuation identities of these processes can be expressed via scale functions.

- 2 野場啓 (京大理) 負スペクトル Lévy 過程における Poisson 的配当の最適化問題 15
 J.-L. Pérez (CIMAT)
山崎和俊 (関西大システム理工)
矢野孝次 (京大理)
- Kei Noba (Kyoto Univ.) On optimal periodic dividend strategies for Lévy risk processes
 José-Luis Pérez (CIMAT)
 Kazutoshi Yamazaki (Kansai Univ.)
 Kouji Yano (Kyoto Univ.)

概要 We study the optimal periodic dividend problem where dividend payments can only be made at the jump times of an independent Poisson process.

- 3 塩沢裕一 (阪大理) Upper rate functions of Brownian motion type for symmetric jump processes 15
 Jian Wang (Fujian Normal Univ.)
- Yuichi Shiozawa (Osaka Univ.) Upper rate functions of Brownian motion type for symmetric jump processes
 Jian Wang (Fujian Normal Univ.)

概要 We are concerned with upper rate functions for a symmetric jump process on the Euclidean space generated by a regular Dirichlet form. We give a condition on the jumping function for the process to enjoy upper rate functions of Brownian motion type. Our condition implies that the second moment of the jumping function is finite.

- 4 長谷部 高広 (北 大 理) 正規分布の自由自己分解可能性 10
 佐久間 紀佳 (愛知教育大教育)
 S. Thorbjørnsen (Aarhus Univ.)
 Takahiro Hasebe (Hokkaido Univ.) Freely selfdecomposability of the normal distributions
 Noriyoshi Sakuma (Aichi Univ. of Edu.)
 Steen Thorbjørnsen (Aarhus Univ.)

概要 We prove that the (classical) normal distributions are freely selfdecomposable. More generally it is established that the Askey–Wimp–Kerov distributions are freely self-decomposable. The main ingredient in the proof is a general characterization of the freely selfdecomposable distributions in terms of the derivative of their free cumulant transform.

- 5 中 田 寿 夫 (福岡教育大) The maxima for the generalized St. Petersburg game 10
 Toshio Nakata (Fukuoka Univ. of Edu.) The maxima for the generalized St. Petersburg game

概要 In this talk, we consider the maxima of payoffs for the generalized St. Petersburg game. The maxima for the original St. Petersburg game cannot be normalized to converge to a nondegenerate limit distribution. However, tuning the parameters appearing in the generalization, we show the normalized maxima converge to the Fréchet distribution.

- 6 鄭 容 武 (広 島 大 工) 平坦な臨界点を持つ単峰写像力学系の大偏差原理について 15
 高 橋 博 樹 (慶 大 理 工)
 Yong Moo Chung (Hiroshima Univ.) Large deviation principle for unimodal maps with flat critical point
 Hiroki Takahasi (Keio Univ.)

概要 We study a smooth unimodal map whose critical point is non-recurrent and flat. Assuming the critical order is either polynomial or logarithmic, we establish the large deviation principle and give a partial description of the zeroes of the corresponding rate function.

- 7 波 止 元 仁 (東京工高専) Relation between mixing properties and chaos in the sense of Devaney
 10
 Jin Hatamoto Relation between mixing properties and chaos in the sense of Devaney
 (Tokyo Nat. Coll. of Tech.)

概要 In this talk, we consider continuous maps of a metric space and introduce relations between mixing properties such as decay of correlations and topologically mixing and Devaney's chaos.

- 8 湯 浅 久 利 (大阪教育大教育)* 無限保測変換に対する厳格エルゴード的相対モデル定理 15
 Hisatoshi Yuasa (Osaka Kyoiku Univ.) A relative, strictly ergodic model theorem for infinite measure-preserving
 systems

概要 Every factor map between given ergodic, measure-preserving transformations on infinite Lebesgue spaces can be realized as a proper, factor map between strictly ergodic, locally compact Cantor systems. A locally compact Cantor system is a topological dynamical system of a homeomorphism on a locally compact (non-compact) metric space, whose one-point compactification is a Cantor set. Such a system, or a homeomorphism, is said to be strictly ergodic if the homeomorphism has a unique, up to scaling, invariant Radon measure and every orbit of it is dense in the metric space.

- 9 田中晴喜 (和歌山県立医大) On convergence of the Gibbs measures of perturbed graph iterated function systems with degeneration 15
 Haruyoshi Tanaka (Wakayama Med. Univ.) On convergence of the Gibbs measures of perturbed graph iterated function systems with degeneration

概要 We study a perturbation of graph iterated function systems (graph IFS). In this study, we consider a family of graph IFSs with small parameter $\epsilon > 0$ such that some functions which compose this IFS converge to constant values as $\epsilon \rightarrow 0$. Therefore, the limit graph IFS (unperturbed system) may have several Gibbs measures associated with the dimension of the limit set even if the parametrized graph IFS (perturbed system) possess a unique Gibbs measure $\mu(\epsilon, \cdot)$ for each $\epsilon > 0$. In this situation, we give a necessary and sufficient condition for convergence of $\mu(\epsilon, \cdot)$ in the case when the limit graph IFS has 2 or 3 Gibbs measures. In particular, this condition is composed of Perron eigenvalues of Ruelle operators.

- 10 平尾将刚 (愛知県立大情報) On p -frame potential of random point configurations on the sphere ... 15
 Masatake Hirao (Aichi Pref. Univ.) On p -frame potential of random point configurations on the sphere

概要 In this talk we deal with two types of random point configurations, spherical ensemble and the jittered sampling on the sphere. The former is a well-studied determinantal point process on the sphere, and the latter is one of the famous random sampling method. We compare these random point configurations with spherical designs, which are one of the non-random “good” point configurations on the sphere in the viewpoint of p -frame potential. We also discuss other random point configurations on the sphere if possible.

14:15~15:00

- 11 南就将 (慶大医) 減衰因子のあるホワイトノイズをポテンシャルとする1次元シュレーディンガー作用素について 15
 Nariyuki Minami (Keio Univ.) One-dimensional Schrödinger operator with decaying white noise potential

概要 We consider the Schrödinger operator whose potential consists of the white noise multiplied by a decaying factor. This model can be defined as a symmetric operator for any realization of white noise, and it is almost surely self-adjoint under some mild conditions. The nature of the positive part of this operator is similar to the model considered by Kotani and Ushiroya (CMP vol. 115 (1988)).

- 12 鈴木由紀 (慶大医) Diffusion processes with random potentials consisting of specially contracted self-similar processes 15
 Yuki Suzuki (Keio Univ.) Diffusion processes with random potentials consisting of specially contracted self-similar processes

概要 Limiting behaviors of one-dimensional diffusion processes with random potentials are studied. The potentials consist of specially contracted self-similar processes. The minimum processes and the maximum processes of the processes are also investigated.

- 13 星野浄生 (阪府大理) Wiener 汎関数の SFC からの広義の再構成式について 10
 Kiyoyuki Hoshino (Osaka Pref. Univ.) On the reconstruction formulas in the wide sense of Wiener functionals from the SFCs

概要 We discuss whether a random function (or a stochastic differential as an extension) is identified and how it is reconstructed from the stochastic Fourier coefficients (SFCs). In this talk, we focus on Skorokhod type SFC and give simple reconstruction formulas using Wiener chaos decomposition from SFCs of Wiener functionals on a space with Haar measure.

15:15~16:15 2017年度(第16回)日本数学会解析学賞受賞特別講演

竹田 雅好(東北大理) 緊密性をもつ対称マルコフ過程の性質

Masayoshi Takeda (Tohoku Univ.) Properties of symmetric Markov processes with tightness property

概要 We consider irreducible, strong Feller, symmetric Markov processes with tightness property, and call the class of symmetric Markov processes having such properties Class (T). If a symmetric Markov process in Class (T) is conservative, then it has a very strong recurrence property, and if not, then it explodes very fast. Using this fact, we can derive some spectral properties, for example, p -independence of the growth bound of its semi-group, compactness of semi-group, bounded continuity of eigenfunctions. As an application of these properties, we can prove the existence and uniqueness of quasi-stationary distribution.

By checking time-changed processes to be in Class (T), we give an analytic characterization of the criticality (or subcriticality) for Schroedinger forms. Moreover, we show that Green function for Schroedinger-type operator satisfies some principles in potential theory (eg. Ugaheri's maximum principle, the continuity principle) if the principal eigenvalue of the time-changed process is greater than one.

16:30~17:30 特別講演

高岡 浩一郎(一橋大商) 数理ファイナンスの第1基本定理—確率積分とマルチンゲール測度—

Koichiro Takaoka (Hitotsubashi Univ.) The first fundamental theorem of asset pricing: stochastic integrals and martingale measures

概要 The first fundamental theorem of asset pricing roughly states that a stochastic process satisfies a property concerning the Itô integrals with respect to it, called the no-arbitrage property, if and only if the process has an equivalent martingale measure. The theorem for discrete-time finite-dimensional processes on finite probability spaces, which goes back to Harrison and Kreps (1979), is a restatement of Stiemke's lemma of finite-dimensional linear algebra. The theorem in the general setting can thus be viewed as an infinite-dimensional extension of Stiemke's lemma. In the continuous-time setting, there is more than one way to formulate the "no-arbitrage property." In this presentation, an emphasis will be put on the L^0 space of random variables and the technique of numeraire change.

3月19日(月) 第VIII会場

9:30~11:30

14 鈴木 聡(島根大総理工) 逆準凸制約を持つ準凸計画問題について 15

Satoshi Suzuki (Shimane Univ.) Quasiconvex programming with a reverse quasiconvex constraint

概要 In this talk, we study quasiconvex programming with a reverse quasiconvex constraint. We introduce affine and quasilinear characterizations of a reverse quasiconvex constraint. By using these characterizations, we show necessary optimality conditions for the problem in terms of Greenberg-Pierskalla subdifferential. Additionally, we investigate surrogate duality for quasiconvex programming with a reverse quasiconvex constraint.

15 堀口 正之(神奈川大理) Adaptive approach in a multivariate Bayesian control chart 15

Masayuki Horiguchi (Kanagawa Univ.) Adaptive approach in a multivariate Bayesian control chart

概要 In this talk, we consider an adaptive control approach in Markov decision process in order to solve a problem of multivariate Bayesian control chart. We show that there exist an average optimal adaptive and asymptotically discounted optimal policies.

- 16 藤田 敏 治 (九 工 大 工) 合流型推移をもつ決定過程について —3つの再帰的アプローチ— 15
 Toshiharu Fujita Decision process with converging branch system —Three types of re-
 (Kyushu Inst. of Tech.) cursive equations—

概要 In this study, we consider a decision process model with a converging branch system which is one of the nonserial transition systems. The model is treated by three approaches. Thus we introduce three types of recursive equations by using dynamic programming technique.

- 17 間 野 修 平 (統 計 数 理 研) A 超幾何分布からの直接抽出とランダム Young 図形への応用 15
 Shuhei Mano (Inst. of Stat. Math.) A direct sampler for A-hypergeometric distributions and its application
 to random Young tableaux

概要 A distribution whose normalization constant is an A-hypergeometric polynomial is called an A-hypergeometric distribution. Such a distribution is in turn a generalization of the generalized hypergeometric distribution on the contingency tables with fixed marginal sums. For sampling from an A-hypergeometric distribution, the first choice may be use of Markov chain Monte Carlo (MCMC) with moves generated by a Markov basis. In this talk, as an alternative to MCMC methods, a direct sampling algorithm for general A-hypergeometric distribution will be presented. As an application of the exact sampler, sampling from random Young tableaux will be discussed. The Ferguson's Dirichlet process is an example of such random Young tableaux. A popular direct sampler, such as the Blackwell–MacQueen's urn scheme, does not work for random Young tableaux without infinite exchangeability. In contrast to the urn schemes, our direct sampler still works without exchangeability.

- 18 山 下 秀 康 (愛知学院大教養) Probabilistic loop path integral for spins 15
 Hideyasu Yamashita Probabilistic loop path integral for spins
 (Aichi Gakuin Univ.)

概要 Let $\text{Spin}(2\ell + 1)$ denote the spin group, represented as a subgroup of $\text{SU}(2^\ell)$ (spin representation). Let G be $\text{SU}(\ell)$ or $\text{Spin}(2\ell + 1)$, and V be \mathbb{C}^ℓ or \mathbb{C}^{2^ℓ} , respectively. Fix $T > 0$ and S^1 denote the loop, viewed as the interval $[0, 2T] \subset \mathbb{R}$ where the endpoints identified. Let H be a self-adjoint operator on V . For any operator A on V , let $A(t) := e^{-itH} A e^{itH}$, $t \in \mathbb{R}$. We show some formulas which give the value of

$$\text{Tr} A_n(t_n) \cdots A_0(t_0) B_1(t'_1) \cdots B_m(t'_m), \quad 0 = t_0 < \cdots < t_n = T, \quad 0 < t'_1 < \cdots < t'_m < T,$$

which can be interpreted as a quantum expectation value, by a limit of probability measures on $C^\infty(S^1, G)$.

- 19 小川重義 (立命館大理工) On the regularity of Gaussian processes indexed by Dirichlet spaces ··· 10
 G. Kerkyacharian
 (LPMA, Univ. Paris-Diderot)
 P. Petrushev
 (Univ. South Carolina)
 D. Picard (Univ. Paris-Diderot)
 Shigeyoshi Ogawa (Ritsumeikan Univ.) On the regularity of Gaussian processes indexed by Dirichlet spaces
 Gerard Kerkyacharian
 (LPMA, Univ. Paris-Diderot)
 Pencho Petrushev
 (Univ. South Carolina)
 Dominique Picard (Univ. Paris-Diderot)

概要 We are concerned with the regularity of centered Gaussian processes $(Z_x(\omega))_{x \in M}$ indexed by compact metric spaces (M, ρ) . We are to show as our main result that the almost everywhere Besov space regularity of such a process is (almost) equivalent to the Besov regularity of the covariance $K(x, y) = E(Z_x Z_y)$ under the assumption that (i) there is an underlying Dirichlet structure on M which determines the Besov space regularity, and (ii) the operator K with kernel $K(x, y)$ and the underlying operator A of the Dirichlet structure commute.

As an application of this result we investigate the case of compact homogeneous spaces and, in particular, the case where M is the sphere.

- 20 嶽村智子 (奈良女大理) Convergence of diffusion processes in a tube ··········· 15
 Tomoko Takemura Convergence of diffusion processes in a tube
 (Nara Women's Univ.)

概要 Diffusion processes in a tube are direct product diffusion processes \mathbb{Y} of one dimensional diffusion processes $X^{(1)}$ and skew product diffusions Ξ , or the time changed process \mathbb{X} which is based on a positive continuous additive functional $\Phi(t)$. The skew product Ξ are given by one dimensional diffusion processes R and a spherical Brownian motion Θ by means of positive continuous additive functional $\mathbf{f}(t)$. We show a limit theorem for a sequence of time changed process \mathbb{X}_n under some assumptions for R_n, ν_n (Revuz measure of $\mathbf{f}_n(t)$), and underlying measure.

11:30~12:00 統計数学分科会総会

3月20日(火) 第VIII会場

10:00~11:40

- 21 弓場 弘 (国際自然研) GA*-optimal balanced third-order designs of resolution $R^*({10,01})$
 兵頭 義史 with $N < \nu(m)$ for 3^m factorials 15
 (岡山理大総合情報研・国際自然研)
 榎田 正秀 (国際自然研)
 Hiromu Yumiba (Int. Inst. for Nat. Sci.) GA*-optimal balanced third-order designs of resolution $R^*({10,01})$
 Yoshifumi Hyodo with $N < \nu(m)$ for 3^m factorials
 (Okayama Univ. of Sci./Int. Inst. for Nat. Sci.)
 Masahide Kuwada
 (Int. Inst. for Nat. Sci.)

概要 We consider the third-order linear model for 3^m factorials. In previous talk (MSJ Autumn Meeting 2017), we gave the A^* -optimal 3^m -BTO designs of resolution $R^*({10,01})$ derived from SA's with the number of assemblies ($= N$) is less than the number of non-negligible factorial effects ($= \nu(m)$) and $m \geq 6$. Let T be a 3^m -BTO design of resolution $R^*({10,01})$ derived from an SA with N assemblies, and further let $\sigma^2 S_T(\alpha)$ ($\alpha = 0, 1, 2$) be the trace of the variance-covariance matrix of the estimators based on T . If $S_T(\alpha) \leq S_{T^*}(\alpha)$ for any T^* , then T is said to be GA_α^* -optimal, where T^* is a 3^m -BTO design of resolution $R^*({10,01})$ derived from an SA with N assemblies. In this talk, we present GA_α^* -optimal 3^m -BTO designs of resolution $R^*({10,01})$ derived from SA's for $m = 6, 7, 8$, where $N < \nu(m)$.

- 22 高見 光広 (東京理大理工) 正方分割表における局所対称モデルからの隔たりを測る調和平均型尺度 10
 三枝 祐輔 (横浜市大医)
 石井 晶 (東京理大理工)
 富澤 貞男 (東京理大理工)
 Mitsuhiro Takami (Tokyo Univ. of Sci.) Measure of departure from local symmetry for square contingency tables
 Yusuke Saigusa (Yokohama City Univ.)
 Aki Ishii (Tokyo Univ. of Sci.)
 Sadao Tomizawa (Tokyo Univ. of Sci.)

概要 In this talk, we propose a new measure for square contingency tables. We construct the measure on the basis of the weighted harmonic mean of the diversity index. We derive properties of the measure and introduce a new model, local symmetry model.

- 23 種市 信裕 (北教大札幌) 多次元分割表における完全独立性検定統計量の改良 15
 関谷 祐里 (北教大釧路)
 外山 淳 (数学利用研)
 Nobuhiro Taneichi Improvement of test of complete independence in multi-dimensional con-
 (Hokkaido Univ. of Edu.) tingency tables.
 Yuri Sekiya (Hokkaido Univ. of Edu.)
 Jun Toyama
 (Inst. for the Practical Application of Math.)

概要 We consider a test of complete independence in multi-dimensional contingency tables. We derive an expression for approximation of the null distribution of the test statistic based on asymptotic expansion. By using the continuous term of the expansion, we consider transformed statistics that increase the speed of convergence to a chi-square limiting distribution.

- 24 池澤友哉 (東京理大理工) 正方分割表における併合した 3×3 表を用いた点対称性からの隔たりを
 生亀清貴 (東京理大理工) 測る尺度 10
 山本紘司 (阪市大医)
 富澤貞男 (東京理大理工)
 Tomoya Ikezawa (Tokyo Univ. of Sci.) Measure of departure from point-symmetry for the collapsed square
 Kiyotaka Iki (Tokyo Univ. of Sci.) contingency tables
 Kouji Yamamoto (Osaka City Univ.)
 Sadao Tomizawa (Tokyo Univ. of Sci.)

概要 For square contingency tables with ordered categories, there may be some cases that one wants to analyze them by considering collapsed tables with some adjacent categories combined in the original table. This presentation considers the point-symmetry model for collapsed square contingency tables and proposes a measure to represent the degree of departure from point-symmetry. Also this presentation gives approximate confidence interval for proposed measure.

- 25 吉本拓矢 正方分割表における共分散対称モデルと対称モデルの分解 15
 (東京理大理工・中外製薬(株))
 田畑耕治 (東京理大理工)
 生亀清貴 (東京理大理工)
 富澤貞男 (東京理大理工)
 Takuya Yoshimoto Covariance symmetry model and decomposition of symmetry model for
 (Tokyo Univ. of Sci./Chugai Pharmaceutical Co., Ltd.) square contingency table
 Kouji Tahata (Tokyo Univ. of Sci.)
 Kiyotaka Iki (Tokyo Univ. of Sci.)
 Sadao Tomizawa (Tokyo Univ. of Sci.)

概要 For the analysis of square contingency table, Caussinus (1965) pointed out that the symmetry model holds if and only if both the quasi-symmetry model and the marginal homogeneity model hold. This presentation proposes the covariance symmetry model and the decomposition theorem of the symmetry model into the covariance symmetry model and the marginal homogeneity model, which are different from Causinnus's.

- 26 鈴木 讓 (阪大基礎工) 連続変量を含む条件付相互情報量の推定 15
 Jo Suzuki (Osaka Univ.) Estimation of conditional mutual information for discrete and continu-
 ous variables

概要 This paper considers to estimate conditional mutual information given three sequences each of which is either continuous or discrete. The estimation generates a sequence of quantizations, estimate conditional mutual information of quantized values, and choose the maximum estimation value. It estimates continuous and discrete variables alike in a seamless manner. In particular, we prove two important properties. First, with probability one as the sample size goes to infinity, the obtained estimation is zero if and only if they are conditionally independent. Secondly, the estimation asymptotically converges to the true value. The procedure has been implemented in the CRAN package BNSL developed by J. Suzuki and J. Kawahara.

14:15~15:05

- 27 矢田 和善 (筑波大数理物質) 高次元固有値推定におけるバイアス補正について 15
青嶋 誠 (筑波大数理物質)
Kazuyoshi Yata (Univ. of Tsukuba) Bias-corrected estimation of eigenvalues in high-dimensional settings
Makoto Aoshima (Univ. of Tsukuba)

概要 In this talk, we consider estimation of eigenvalues in high-dimensional settings. First, we show that the sample eigenvalue is not a consistent estimator of the true eigenvalue for high-dimensional settings. Yata and Aoshima (2012, JMVA) proposed a new PCA method called the noise reduction (NR) methodology. The estimation of the eigenvalue by the NR method has a first-order consistency. We investigate more deeply the asymptotic behavior of the NR method. We give a new eigenvalue estimation and show that it holds the second-order consistency.

- 28 石井 晶 (東京理大理工) ノイズ掃き出し法を用いた高次元共分散行列の同等性検定 15
矢田 和善 (筑波大数理物質)
青嶋 誠 (筑波大数理物質)
Aki Ishii (Tokyo Univ. of Sci.) Equality tests of high-dimensional covariance matrices by using the
Kazuyoshi Yata (Univ. of Tsukuba) noise-reduction methodology
Makoto Aoshima (Univ. of Tsukuba)

概要 In this talk, we consider the equality tests of covariance matrices for high-dimensional data. Aoshima and Yata (2017) proposed two eigenvalue models for high-dimensional data and constructed two-sample test procedures. One is called strongly spiked eigenvalue (SSE) model and the other one is called non-SSE (NSSE) model. Ishii et al. (2016) proposed an equality test of two covariance matrices under the SSE model. Li and Chen (2012) proposed a test procedure under the NSSE model. We evaluate the test statistic of Li and Chen (2012) under the SSE model and give new test procedures by using the noise-reduction method given by Yata and Aoshima (2012). We also compare our new test procedures with that given by Ishii et al. (2016).

- 29 前園 宜彦 (九大数理) Smoothed two-sample nonparametric tests and their asymptotic prop-
森山 卓 (九大数理) erties 10
Yoshihiko Maesono (Kyushu Univ.) Smoothed two-sample nonparametric tests and their asymptotic prop-
Taku Moriyama (Kyushu Univ.) erties

概要 In this paper we discuss smoothed rank statistics for testing a location shift parameter of the two-sample problem. They are based on the discrete test statistics—the median and Wilcoxon’s rank sum tests. For the one-sample problem, Maesono et al. (2017) reported that some nonparametric discrete tests have a problem with their p -values because of their discreteness. The p -values of the Wilcoxon’s test are frequently smaller than those of the median test in tail area. This causes an arbitrary choice of the median and Wilcoxon’s rank sum tests. In order to conquer this problem, we propose smoothed versions of those tests. The smoothed tests inherit good properties of the original tests, and asymptotically equivalent to the original test statistics. We study significance probabilities and local asymptotic powers of the proposed tests.

15:20~16:20 特別講演

松原 和樹 (中央学院大商) BIB デザインの組加法性と関連する組合せ構造

Kazuki Matsubara (ChuoGakuin Univ.) Pairwise additivity of BIB designs and related combinatorial structures

概要 A set of ℓ balanced incomplete block (BIB) designs with common parameters having pairwise additivity is called an ℓ -pairwise additive BIB design. The ℓ -pairwise additivity of BIB designs can be regarded as a decomposition of a BIB design and/or a composition of several BIB designs, and further yields BIB designs with ℓ distinct block sizes. In this talk, we discuss comprehensively the existence of ℓ -pairwise additive BIB designs through the relationship with other combinatorial structures. Especially, as related combinatorial structures, we focus on nested BIB designs, orthogonal arrays (mutually orthogonal latin squares), perpendicular arrays, difference matrices and finite geometries. Furthermore, we present some results on splitting-balanced block designs named newly. The pairwise additivity also relates to the splitting-balanced property. The splitting-balanced block design (also called a splitting BIB design in literature) was introduced for applications to authentication codes by Ogata et al. in 2004. Finally, the bound of the number of blocks and some existence of splitting-balanced block designs are provided.

16:40~17:40 特別講演

佃 康司 (東大総合文化) ヒルベルト空間における弱収束理論とその応用

Koji Tsukuda (Univ. of Tokyo) Studies on the theory of weak convergences in Hilbert spaces and its applications

概要 In this presentation, weak convergences of random processes taking values in a separable Hilbert space are discussed. Especially, by using a common approach based on the limit theory in L^2 spaces, we study partial sum processes of dependent random variables which appear in the following two topics: (i) statistical change point testing; (ii) functional central limit theorem for logarithmic combinatorial assemblies. (i) Statistical change point testing. When there exist chronologically obtained data, it is in interest to determine whether there are structural changes or not. In the domain of statistics, tests for such hypothesis is considered in the so-called change point testing. For parametric change point problems, several works have proposed test procedures based on score functions. Following these works, we propose a procedure based on the functional of a weighted random process whose weak convergence is discussed in $L^2(0, 1)$, which is a modification of test statistics in previous works. (ii) Functional central limit theorem for logarithmic combinatorial assemblies. Assemblies are a class of random combinatorial structures which includes random permutations, random mappings, random forests of labeled trees, and so on. The law of component counts of assemblies is provided from independent Poisson variables combined with the conditioning relation. In previous works, the weak convergence of the partial sum process of component counts of logarithmic assemblies, which are assemblies satisfying the logarithmic condition, has been considered in the Skorokhod space. On the other hand, we consider the partial sum process involving a weight function, which makes the limit different, and derive its weak convergence in $L^2(0, 1)$.

3月21日(水) 第VIII会場

10:00~12:00

- 30 吉田 朋 広 (東 大 数 理) Martingale expansion and power variation 15
 Nakahiro Yoshida (Univ. of Tokyo) Martingale expansion and power variation

概要 Inference for volatility under finite time horizon becomes non-ergodic statistics. The quasi-maximum likelihood estimator and the Bayesian type estimator of the volatility parameter are asymptotically mixed normal in general. Asymptotic expansion in non-ergodic systems is then indispensable to develop the higher-order inferential theory for volatility. We present asymptotic expansion of a martingale having a mixed normal limit. The expansion formula is expressed by the adjoint of a random symbol with coefficients described by the Malliavin calculus, differently from the standard invariance principle. We discuss its application to the power variation of a diffusion process. Identification of the random symbols is an issue.

- 31 吉田 朋 広 (東 大 数 理) Asymptotic expansion of Skorohod integrals 15
 Nakahiro Yoshida (Univ. of Tokyo) Asymptotic expansion of Skorohod integrals

概要 Asymptotic expansion of the distribution of the Skorohod integral jointly with a reference variable is derived. We introduce a second-order interpolation formula in frequency domain to expand a characteristic functional and combine it with the scheme developed in the martingale expansion. Random symbols are used for expressing the asymptotic expansion formula. Quasi tangent, quasi torsion and modified quasi torsion are introduced in this paper. This is a joint work with D. Nualart.

- 32 谷田 義 行 (早 大 理 工) 高次元時系列における Whittle 推定量の漸近理論とその数値例 10
 明石 郁 哉 (早 大 理 工)
 谷口 正 信 (早 大 理 工)
 Yoshiyuki Tanida (Waseda Univ.) Asymptotic theory and numerical studies of Whittle estimation for high-
 Fumiya Akashi (Waseda Univ.) dimensional time series
 Masanobu Taniguchi (Waseda Univ.)

概要 In this presentation, we develop the estimation theory for Whittle functional of high-dimensional non-Gaussian dependent processes. Using a sample version based on a thresholded periodogram matrix, we introduce a thresholded Whittle estimator of unknown parameter, and elucidate its asymptotics. It is shown that the thresholded Whittle estimator is a \sqrt{n} -consistent estimator of the unknown parameter, and that the standardized version has the asymptotic normality. Some numerical studies illuminate an interesting feature of the results. Concretely, for high-dimensional AR(2), we compared the difference of RMSE between the usual Whittle estimator $\hat{\theta}_w$ and the thresholded estimator $\hat{\theta}_{w,th}$, leading to a conclusion that $\hat{\theta}_{w,th}$ is better than $\hat{\theta}_w$.

- 33 Yujie Xue (早 大 理 工) LASSO estimators for high-dimensional time series with long-memory
 谷口 正 信 (早 大 理 工) disturbances 10
 Yujie Xue (Waseda Univ.) LASSO estimators for high-dimensional time series with long-memory
 Masanobu Taniguchi (Waseda Univ.) disturbances

概要 Consider a linear regression model: $Y_t = z_t' \beta + \varepsilon_t$ where $\{\varepsilon_t\}$ is a stationary process with mean zero and spectral density $f(\lambda)$, and z_t is a known nonrandom function vector of t . In this talk, it is desired to discuss the LASSO estimator of β when $\{\varepsilon_t\}$ is a long-memory strictly stationary process (i.e. $f(\lambda)$ is unbounded at the origin) all of whose moments exist and has the infinite moving average representation, and when the dimension of β defined as p increases with sample size n . An interesting property of the LASSO estimators is shown.

- 34 藤 森 洸 (早 大 理 工) Cox's proportional hazards model with a high-dimensional and sparse regression parameter 15
 Kou Fujimori (Waseda Univ.) Cox's proportional hazards model with a high-dimensional and sparse regression parameter

概要 This talk deals with the proportional hazards model proposed by D. R. Cox in a high-dimensional and sparse setting for a regression parameter. To estimate the regression parameter, the Dantzig selector is applied. The variable selection consistency of the Dantzig selector for the model will be proved. This property enables us to reduce the dimension of the parameter and to construct asymptotically normal estimators for the regression parameter and the cumulative baseline hazard function.

- 35 明 石 郁 哉 (早 大 理 工) 自己加重型 GEL 統計量の局所検出力及び加重関数選択手法 15
 Fumiya Akashi (Waseda Univ.) Local asymptotic power of self-weighted GEL method and choice of weighting function

概要 Recently, we often observe the heavy-tailed time series data in variety of fields, and it is unfeasible to apply the classical likelihood ratio-based method to such data directly. To overcome the difficulty, this talk constructs the self-weighted generalized empirical likelihood (SW-GEL) statistic for possibly infinite variance processes, and elucidates the local asymptotic power of the SW-GEL statistic. The self-weighting method proposed by Ling (2005, JRSS) enables us to control effects brought by the infinite variance of underlying time series models. By the self-weighting method, the proposed statistic converges to the non-central chi-square distribution under the local alternatives. This talk also introduces the selection procedure of tuning parameters in self-weights based on the local asymptotic power.

- 36 福 田 光 (阪 大 基 礎 工) 高頻度観測の下での安定過程の局所漸近正規性 15
 深 澤 正 彰 (阪 大 基 礎 工)
 Hikaru Fukuda (Osaka Univ.) Local asymptotic normality property for stable processes under high-frequency observations
 Masaaki Fukasawa (Osaka Univ.)

概要 It is well known that the property of local asymptotic normality (LAN) allows us to discuss the asymptotic efficiency of estimation via minimax theorems. We proved LAN property for symmetric stable processes and one-sided stable processes under high-frequency observations using non-diagonal rate matrices depending on the parameter to be estimated. In contrast to the classical LAN families in the literature, non-diagonal rate matrices are inevitable.

応 用 数 学

3月18日(日) 第VI会場

9:30~11:45

- 1 足立智子 (東邦大理) サイクリックになる二部グラフのラベル付けと RAID への応用 10
Tomoko Adachi (Toho Univ.) A labeling of a cyclic bipartite graph and its application to RAID

概要 A cluttered ordering is a kind of cyclic orderings, and can be used to minimize the number of disk operations in RAID system. Mueller et al. (2005) decomposed the complete bipartite graph into isomorphic copies of the special bipartite graph $H(h; t)$, where h and t are positive integers. The special bipartite graph $H(h; t)$ has each $h(t+1)$ vertices as upper vertex set and lower vertex set. In this talk, we define the special bipartite graph $H(h, k; t)$, where $h, k (h \neq k)$ and t are positive integers. The special bipartite graph $H(h, k; t)$ has $h(t+1)$ vertices as upper vertex set and $k(t+1)$ vertices as lower vertex set. We present some edge labeling of the infinite families of $H(1, 2; t)$.

- 2 M. Buchet Vietoris–Rips realization of indecomposable persistence modules of arbitrarily large dimension 15
(東北大 AIMR • TU Graz)
E. G. Escolar (東北大 AIMR)
Mickaël Buchet Vietoris–Rips realization of indecomposable persistence modules of arbitrarily large dimension
(Tohoku Univ./TU Graz)
Emerson Gaw Escolar (Tohoku Univ.)

概要 In this work we construct an infinite family, parametrized by dimension order d , of non-isomorphic indecomposable persistence modules over the commutative ladder of length 5. We provide a family of bifiltrations of topological spaces whose H_1 persistent homologies is the infinite family of persistence modules. Moreover, we provide Vietoris–Rips constructions of the family of bifiltrations. Our construction provides an elementary proof of the fact that the commutative ladders with length greater than or equal to 5 is representation-infinite. Furthermore, we aim by this example to illustrate that indecomposable persistence modules of high dimension as a representation may encode some interesting and easy to visualize phenomenon.

- 3 瀬戸道生 (防衛大) Application of the theory of quasi-orthogonal integrals to graph theory 10
須田庄 (愛知教育大)
Michio Seto Application of the theory of quasi-orthogonal integrals to graph theory
(Nat. Defense Acad. of Japan)
Sho Suda (Aichi Univ. of Edu.)

概要 In this talk, we give two inequalities concerning increasing families of finite graphs. Those inequalities are derived from the theory of quasi-orthogonal integrals developed by de Branges–Rovnyak and Vasyunin–Nikol’skii.

- 4 佐竹翔平 (神戸大システム情報) Remark on two typical properties of random graphs 15
Shohei Satake (Kobe Univ.) Remark on two typical properties of random graphs

概要 There are some well known properties which random graphs satisfy with probability 1. Here we deal with two such typical properties of random graphs, namely, the n -existentially closed (n -e.c.) property and the pseudo-random property. Cameron and Stark remarked that the pseudo-random property does not necessarily imply the n -e.c. property for large n . However there seems no results about the inverse relationship.

In this talk, we give the first construction of n -e.c. graphs without the pseudo-random property, which means that the n -e.c. property also does not necessarily imply the pseudo-random property for every n .

- 5 金澤 秀 (東 北 大 理) ランダム複体過程における生存時間和の漸近評価 15
 日野 正訓 (京 大 理)
 Shu Kanazawa (Tohoku Univ.) Asymptotic behavior of lifetime sums for random simplicial complex
 Masanori Hino (Kyoto Univ.) processes

概要 We study the homological properties of random simplicial complexes, which have received a lot of attention over the past several years. In particular, we obtain asymptotic behavior of lifetime sums of persistent homology for a class of increasing random simplicial complexes, which is a higher-dimensional counterpart of Frieze's zeta function theorem for the Erdős-Rényi graph process. Main results include solutions to the questions on the Linial-Meshulam complex process and the clique complex process that were posed in the preceding study by Hiraoka and Shirai. One of the key ingredients of the arguments is a new upper bound of Betti numbers of general simplicial complexes in terms of the number of small eigenvalues of Laplacians on links, which is regarded as a quantitative version of the cohomology vanishing theorem.

- 6 鈴木 航介 (広 島 大 理) 軸平行な直方体内にある Chebyshev-Frolov 格子点の列挙アルゴリズム
 芳木 武仁 (京 大 情 報) 15
 Kosuke Suzuki (Hiroshima Univ.) Enumeration of the Chebyshev-Frolov lattice points in axis-parallel
 Takehito Yoshiki (Kyoto Univ.) boxes

概要 For a positive integer d , the d -dimensional Chebyshev-Frolov lattice is the \mathbb{Z} -lattice in \mathbb{R}^d generated by the Vandermonde matrix associated to the roots of the d -dimensional Chebyshev polynomial. It is important to enumerate the points from the Chebyshev-Frolov lattices in axis-parallel boxes when $d = 2^n$ for a non-negative integer n , since the points are used for the nodes of Frolov's cubature formula, which achieves the optimal rate of convergence for many spaces of functions with bounded mixed derivatives and compact support. The existing enumeration algorithm for such points by Kacwin, Oettershagen and Ullrich is efficient up to dimension $d = 16$. In this paper we suggest a new enumeration algorithm of such points for $d = 2^n$, efficient up to $d = 32$.

- 7 阿川 真士 (九 大 数 理) 位相空間の圏と同型な関係 T 代数について 15
 溝口 佳寛 (九 大 I M I)
 Naoto Agawa (Kyushu Univ.) A relational T-algebra isomorphic to the category of the topological
 Yoshihiro Mizoguchi (Kyushu Univ.) spaces

概要 A continuous lattice is a semantic domain of a computation such as a lambda calculus. Our motivation comes from a development of a formal theory of semantic domains. We first talk about a theory of relational T-algebra, an extension of a T-algebra. Michael Barr proved the category of relational T-algebra defined by the ultra-filter monad is isomorphic to the category of topological spaces. We review their theory and reformulate it using our simple framework of a relational calculus.

- 8 須田 庄 (愛 知 教 育 大) The Smith normal form of skew-symmetric D-optimal designs of order
 G. Greaves (Nanyang Tech. Univ.) $n \equiv 2 \pmod{4}$ 10
 Sho Suda (Aichi Univ. of Edu.) The Smith normal form of skew-symmetric D-optimal designs of order
 Gary Greaves (Nanyang Tech. Univ.) $n \equiv 2 \pmod{4}$

概要 We show that the Smith normal form of a skew-symmetric D-optimal design of order $n \equiv 2 \pmod{4}$ is determined by its order. We apply our result to show that certain D-optimal designs of order $n \equiv 2 \pmod{4}$ are not equivalent to any skew-symmetric D-optimal design.

- 9 野崎 寛 (愛知教育大) Maximizing the order of a bipartite regular graph for given valency and second eigenvalue 15
 Hiroshi Nozaki (Aichi Univ. of Edu.) Maximizing the order of a bipartite regular graph for given valency and second eigenvalue

概要 Let $v(k, \lambda)$ be the maximum order of connected bipartite k -regular graphs whose second-largest eigenvalues are at most λ . We show an upper bound for $v(k, \lambda)$, which is based on the linear programming bound. If a graph attains the bound, then it is a distance-regular graph that satisfies $g \geq 2d - 2$, where g is the girth and d is the diameter of the graph. There are infinitely many bipartite distance-regular graphs that satisfy $g \geq 2d - 2$. We can prove the non-existence of bipartite distance-regular graphs with $g \geq 2d - 2$ for $d > 26$ by the manner of Fuglister (1987). This is a joint work with Sebastian Cioabă and Jack Koolen.

14:20~16:30

- 10 Yandong Bai Kernels by properly colored paths in arc-colored digraphs 10
 (Northwestern Polytechnical Univ.)
 藤田 慎也 (横浜市国際総合)
 Shenggui Zhang
 (Northwestern Polytechnical Univ.)
 Yandong Bai Kernels by properly colored paths in arc-colored digraphs
 (Northwestern Polytechnical Univ.)
 Shinya Fujita (Yokohama City Univ.)
 Shenggui Zhang
 (Northwestern Polytechnical Univ.)

概要 A subdigraph H of an arc-colored digraph D is *properly colored* if any two consecutive arcs of H receive distinct colors. A *kernel by properly colored paths* of an arc-colored digraph D is a set S of vertices of D such that (i) no two vertices of S are connected by a properly colored directed path in D , and (ii) every vertex outside S can reach S by a properly colored directed path in D . We conjecture that every arc-colored digraph with all cycles properly colored has such a kernel and verify the conjecture for unicyclic digraphs, semi-complete digraphs and bipartite tournaments, respectively.

- 11 藤沢 潤 (慶大商) Distance matching extension in cubic bipartite graphs 15
 R. E. L. Aldred (Univ. of Otago)
 斎藤 明 (日大文理)
 Jun Fujisawa (Keio Univ.) Distance matching extension in cubic bipartite graphs
 R. E. L. Aldred (Univ. of Otago)
 Akira Saito (Nihon Univ.)

概要 A matching M of a graph G is said to be extendable if M is a subset of a perfect matching of G , and M is said to be distance d matching if the edges of M lie pair-wise distance at least d . If every distance d matching of G is extendable, then we say that G is distance d matchable. In this talk we introduce the following results: 1) Let G be a 3-connected cubic bipartite graph. If there exist two cycles C_1, C_2 of length at most d such that $E(C_1) \cap E(C_2) = \{e\}$ for every $e \in E(G)$, then G is distance d matchable. 2) Let G be a 3-connected cubic bipartite planar graph. If there exist two cycles C_1, C_2 of length at most 6 such that $e \in E(C_i)$ ($i = 1, 2$) for every $e \in E(G)$, then G is distance 6 matchable.

- 12 大野由美子 (横浜国大環境情報) トーラス上の三角形分割の n -triad coloring 15
 Yumiko Ohno (Yokohama Nat. Univ.) n -Triad colorings of triangulations on the torus

概要 A *triangulation* on a closed surface is a graph embedded on the surface each of whose face is triangular. Let G be a triangulation on a closed surface and $n \geq 3$ be a natural number. A coloring $c : V(G) \rightarrow \mathbb{Z}_n$ is called an n -triad coloring if $\{c(u), c(v), c(w)\}$ belongs to $\{\{i, i+1, i+2\} \mid i \in \mathbb{Z}_n\}$ for any face uvw of G . We would like to determine the set of numbers n such that G has n -triad colorings. The set can be determined completely by the chromatic number of G if G is embedded on the sphere or the projective plane. In this talk, we shall focus on G which is embedded on the torus and investigate the above set of G .

- 13 朝山芳弘 (横浜国大環境情報) 3-dynamic coloring for triangulations on surfaces 15
 川崎雄貴 (横浜国大環境情報)
 Seog-Jin Kim (Konkuk Univ.)
 中本敦浩 (横浜国大教育人間)
 小関健太 (横浜国大環境情報)
 Yoshihiro Asayama 3-dynamic coloring for triangulations on surfaces
 (Yokohama Nat. Univ.)
 Yuki Kawasaki (Yokohama Nat. Univ.)
 Seog-Jin Kim (Konkuk Univ.)
 Atsuhiko Nakamoto
 (Yokohama Nat. Univ.)
 Kenta Ozeki (Yokohama Nat. Univ.)

概要 An r -dynamic k -coloring of a graph G is a proper k -coloring such that any vertex v has at least $\min\{r, \deg_G(v)\}$ distinct colors in $N_G(v)$. The r -dynamic chromatic number $\chi_r^d(G)$ of a graph G is the least k such that there exists an r -dynamic k -coloring of G . Loeb and et al, proved that $\chi_3^d \leq 10$ if G is a planar graph, however this result is not considered not to be sharp. Thus finding an optimal upper bound on $\chi_3^d(G)$ for a planar graph G is a natural interesting problem. We will show some upper bounds on $\chi_3^d(G)$ for triangulations on the plane, the projective plane and the torus.

- 14 永並健吾 (横浜国大環境情報) 球面以外の閉曲面に埋め込まれた 3-連結 3-正則平面的グラフ 15
 Kengo Enami (Yokohama Nat. Univ.) 3-connected 3-regular planar graphs embedded on non-spherical surfaces

概要 It is well-known that every 3-connected planar graph is uniquely embeddable on the sphere but it is not uniquely embeddable on any surface other than the sphere. We shall focus on a 3-connected 3-regular planar graph and classify structures of its embeddings on the torus, the projective plane and the Klein bottle.

- 15 小関健太 (横浜国大環境情報) 射影平面上の正則グラフの符号 15
 阿部敏生 (横浜国大環境情報)
 Kenta Ozeki (Yokohama Nat. Univ.) The signature of edge colorings on the projective plane
 Toshiki Abe (Yokohama Nat. Univ.)

概要 Let G be a k -regular k -edge colorable graph. Moreover, let φ be a k -edge coloring of G . Let v be a vertex of $V(G)$ and $E(v)$ denotes the set of all edges incident with v . Let ρ_v be a bijective map $E(v) \rightarrow \{1, 2, \dots, k\}$ and we call the set $\rho = \{\rho_v : v \in V(G)\}$ basis. Now we obtain the correspondence $\pi_v = \varphi \circ \rho_v^{-1}$ for every v . The signature of φ is defined as $\prod_{v \in V(G)} \text{sign}(\pi_v)$. In this talk, we obtain the signatures of k -edge-colorings in k -regular graphs on the projective plane for a certain basis ρ . Moreover, this has an application to the list coloring conjecture for certain graph class by using former result.

- 16 野口 健太 (東京電機大情報) 偶三角形分割における辺数の大きな二部的全域部分グラフ 15
 Kenta Noguchi (Tokyo Denki Univ.) Spanning bipartite subgraphs having large size of even triangulations

概要 We consider a spanning bipartite subgraph of an even (i.e. Eulerian) triangulation G on a surface. If G has a spanning bipartite quadrangulation, then its size is two-thirds of $E(G)$. In this talk, we discuss the condition of G having a spanning bipartite quadrangulation Q . We also discuss the maximum size of a spanning bipartite subgraph of G when G does not have Q .

- 17 鈴木 有祐 (新潟大理) 例外的 3-染色的三角形分割について 15
 村井 聡 (阪大情報)
 Yusuke Suzuki (Niigata Univ.) Exceptional balanced triangulations on closed surfaces
 Satoshi Murai (Osaka Univ.)

概要 In our recent research, we proved that two balanced (or 3-colorable) triangulations of a closed surface are not necessarily connected by a sequence of balanced stellar subdivisions and welds. This answers a question posed by Izmistiev, Klee and Novik. In this talk, we especially discuss two local operations called a pentagon contraction and a pentagon splitting, which are also defined for balanced triangulations of closed surfaces. We show that most two balanced triangulations of a closed surface are transformed into each other by a sequence of the above two operations. Furthermore, we introduce such exceptional balanced triangulations of closed surfaces with low genera.

16:45~17:45 特別講演

- 篠原 雅史 (滋賀大教育) 距離集合における分類問題と極値問題
 Masashi Shinohara (Shiga Univ.) Classification problems and extremal problems on distance sets

概要 We consider good configurations from the point of view of the theory of distance sets. A subset X of a Euclidean space is called a k -distance set if there exists exactly k values of distances between two distinct points in X . The study of distance sets was initiated by Erdős (1946). One of the major problems in the theory of k -distance sets in the d -dimensional Euclidean space is to determine the largest possible cardinality $g_d(k)$ of k -distance sets and classify the distance sets X satisfying $|X| = g_d(k)$. Such k -distance sets are said to be optimal. Furthermore, we are also interested in characterization of k -distance sets with large points relative to k . In this talk, we introduce some results for optimal k -distance sets and extremal problems on distance sets.

3月19日(月) 第VI会場

9:30~11:45

- 18 佐藤 巖 (小山工高専) A weighted generalized Bartholdi zeta function of a digraph 15
 三橋 秀生 (法政大理工)
 森田 英章 (室蘭工大工)
 Iwao Sato (Oyama Nat. Coll. of Tech.) A weighted generalized Bartholdi zeta function of a digraph
 Hideo Mitsuhashi (Hosei Univ.)
 Hideaki Morita (Muroran Inst. of Tech.)

概要 We define a weighted generalized Bartholdi zeta function and a weighted generalized Bartholdi L -function of a digraph, and present their determinant expressions. Furthermore, we give express the weighted generalized Bartholdi zeta function of a group covering of a digraph by a product of its weighted generalized Bartholdi L -functions.

- 19 井手 勇介 (神奈川県大工) 空間的に非一様な 1 次元 2 状態の量子ウォーク 10
 今野 紀雄 (横浜国大工)
 中山 大地 (横浜国大工)
 Yusuke Ide (Kanagawa Univ.) Two-state space-inhomogeneous coined quantum walk
 Norio Konno (Yokohama Nat. Univ.)
 Daichi Nakayama
 (Yokohama Nat. Univ.)

概要 We consider the two-state space-inhomogeneous coined quantum walk (QW) in one dimension. We obtain the uniform measure as the stationary measure by solving the eigenvalue problem. This approach is based on the method giving by Kawai, Komatsu and Konno. (2017).

- 20 小松 堯 (横浜国大理工) 2次元トーラス上の量子ウォーク 10
 今野 紀雄 (横浜国大工)
 成松 明廣 (横浜国大工)
 Takashi Komatsu The Quantum walk on the 2-dimensional torus
 (Yokohama Nat. Univ.)
 Norio Konno (Yokohama Nat. Univ.)
 Akihiro Narimatsu
 (Yokohama Nat. Univ.)

概要 We analyze two types of the Quantum walk on the 2-dimensional torus, the Grover walk and the Fourier walk, and obtain provability amplitude and the search algorithm using the Fourier walk.

- 21 船川 大樹 (北大理工) 2次元スプリットステップ量子ウォークの固有値解析 15
 布田 徹 (北大理工)
 笹山 智司 (北大理工)
 鈴木 章斗 (信州大工)
 Daiju Funakawa (Hokkaido Univ.) Eigenvalue analysis of a 2-dimensional quantum walk
 Toru Fuda (Hokkaido Univ.)
 Satoshi Sasayama (Hokkaido Univ.)
 Akito Suzuki (Shinshu Univ.)

概要 We consider the 2-dimensional 4-states quantum walk. This quantum walk is an extension of the 1-dimensional split-step quantum walk. By the earlier study, it is known that if a particular function f has zero points, then time evolution operator U has eigenvalues. In such a case, localization occurs. In this talk, we introduce the necessary and sufficient condition of f has zero points. This research is a joint work with T. Fuda, S. Sasayama and A. Suzuki.

- 22 寺田 知幸 (金沢工大) 1次元系上のものぐさフーリエウォーク 10
 小松 堯 (横浜国大理工)
 今野 紀雄 (横浜国大工)
 齋藤 溪 (横浜国大工)
 Tomoyuki Terada Study on the one-dimensional lazy Fourier walk
 (Kanazawa Inst. of Tech.)
 Takashi Komatsu
 (Yokohama Nat. Univ.)
 Norio Konno (Yokohama Nat. Univ.)
 Kei Saito (Yokohama Nat. Univ.)

概要 We consider the property of lazy Fourier walk in one dimension. For example, stationary measure, time-averaged limit measure, and periodicity.

- 23 大野 博道 (信州大工) 1次元量子ウォークのユニタリ同値類 15
 Hiromichi Ohno (Shinshu Univ.) Unitary equivalence classes of one-dimensional quantum walks

概要 This study investigates the unitary equivalence classes of one-dimensional quantum walks. We determine the unitary equivalence classes of one-dimensional quantum walks, two-phase quantum walks with one defect, complete two-phase quantum walks, one-dimensional quantum walks with one defect and translation-invariant quantum walks.

- 24 小松 堯 (横浜国大理工) 高次元整数格子上の量子ウォークの定常測度 15
 今野 紀雄 (横浜国大工)
Takashi Komatsu Stationary measures of quantum walks on the higher-dimensional integer lattice
 (Yokohama Nat. Univ.)
 Norio Konno (Yokohama Nat. Univ.)

概要 Stationary measures of quantum walks on the one-dimensional integer lattice are well studied. However, the stationary measure for the higher dimensional case has not been clarified. In this talk, we give the stationary amplitude for quantum walks on the higher-dimensional integer lattice with a finite support by solving the corresponding eigenvalue problem. As a corollary, we can obtain the stationary measures of the Grover walks.

- 25 小松 堯 (横浜国大理工) 空間非一様な量子ウォークの定常測度 10
 川合 光 (横浜国大工)
 今野 紀雄 (横浜国大工)
Takashi Komatsu Stationary measures of space-inhomogeneous quantum walks
 (Yokohama Nat. Univ.)
 Hikari Kawai (Yokohama Nat. Univ.)
 Norio Konno (Yokohama Nat. Univ.)

概要 We consider the two-state space-inhomogeneous coined quantum walk in one dimension. For a general setting, we obtain the stationary measures of the quantum walks by solving the eigenvalue problem. As a corollary, stationary measures of the multi-defect model and space-homogeneous quantum walks are derived.

- 26 吉江 佑介 (東北大情報) グラフのジョイントとグローヴァーウォークの周期性 15
 Yusuke Yoshie (Tohoku Univ.) Periodicity of the Grover walk on joint of graphs

概要 The Grover walk is a kind of quantum walks on graphs, and it is applied to various study fields. The Grover walk is determined by a unitary time evolution operator given by the underlying graph. So we can say that the Grover walk is induced by the graph. We focus on characterizations of graphs inducing periodic Grover walks, that is, there exists a integer k such that k -th iteration of the time evolution operator becomes identity operator. In previous work, such graphs have been found, e.g. cycle graphs, path graphs, complete bipartite graphs. In this talk, we construct new graphs with these graphs, and consider the periodicity of the Grover walk on these graphs.

14:00~14:20 2017年度日本数学会応用数学研究奨励賞授賞式

3月20日(火) 第VI会場

9:15~11:50

- 27 草野元紀(東北大理) Persistence weighted Gaussian kernel によるブートストラップ法 15
Genki Kusano (Tohoku Univ.) The bootstrap method by the Persistence weighted Gaussian kernel

概要 In topological data analysis, the topological structures in data are transformed into a persistence diagram, and its statistical method is proposed by the Persistence weighted Gaussian kernel (PWGK). Here, let us consider the expectation of persistence diagrams by the PWGK. While it is difficult to calculate the expectation in general, the confidence set which contains the true expectation can be constructed by the bootstrap method. In this talk, we will discuss the bootstrap method by the PWGK and its asymptotic consistency.

- 28 竹内博志(東北大理) 対応のパーシステントホモロジー: 箆の表現論の視点から 15
平岡裕章(東北大AIMR)
Hiroshi Takeuchi (Tohoku Univ.) The persistent homology of a correspondence: A viewpoint from quiver
Yasuaki Hiraoka (Tohoku Univ.) representations

概要 The theory of homology induced maps of correspondences proposed by Shaun Harker et al. is a powerful tool which allows the retrieval of underlying homological information from sampling data with noise or defects. In this study, we redefine induced maps of correspondences within the framework of quiver representations, and provide more concise proofs of the main theorems in the original paper. With this point of view, we easily extend these ideas to filtration analysis, which provides a new method for analyzing dynamical systems.

- 29 大林一平(東北大AIMR) Volume optimal cycles for persistent homology 15
Ipppei Obayashi (Tohoku Univ.) Volume optimal cycles for persistent homology

概要 Persistent homology (PH) is a significant tool for topological data analysis, which analyzes shape of data efficiently and quantitatively. A persistence diagram (PD) is a visualization tool of PH, which is a multiset on $\mathbb{R} \times (\mathbb{R} \cup \infty)$. Each point on a PD (called a birth-death pair) corresponds a homological structure such as a ring, cavity, etc. appearing in the data. For a practical application of PH, we want to identify such a homological structure for a selected birth-death pair. The idea of a volume optimal cycle and the computation algorithm are proposed in this talk.

- 30 寺本敬(旭川医科大医) The case studies of using computational homology in medical science
灌山晃弘 10
(北海道文教大人間科学)
Takashi Teramoto The case studies of using computational homology in medical science
(Asahikawa Medical Univ.)
Akihiro Takiyama
(Hokkaido Bunkyo Univ.)

概要 We present a methodology using computation homology for quantitative measurements in medical science, in collaboration with diagnostic doctors. Our talk will consist of two folds: cubical homology index for bone morphometry in three-dimension, and immunohistochemical scoring based on persistent homology.

- 31 岡 宏 枝 (龍谷大理工) The Conley index over base 15
 神山 蒼 太 (龍谷大理工)
 Hiroe Oka (Ryukoku Univ.) The Conley index over base
 Sota Koyama (Ryukoku Univ.)

概要 The Conley index over a base [Mrozek Reineck, Szrednicki 1997] is a generalization of the Conley index for flows [Conley 1972]. In the case when the base is the circle, it is naturally related to the discrete type of Conley index of a Poincare section of the flow. We will consider the homology version of the Conley index over a base, and discuss the relation between them.

- 32 堀 口 俊 二 拡張複素ニュートン法の根の収束と分布について 15
 Shunji Horiguchi On convergences and distributions of roots of extended complex Newton's method

概要 We extend the complex Newton's method. We give the followings for the extended complex Newton's method. Relationship between extended complex Newton's method and Riemann surface. Conditional expression of initial values for convergence of extended complex Newton's method. Speeds of convergences of extended complex Newton's method. The distributions of roots of extended complex Newton's Method.

- 33 坂 口 文 則 (福 井 大 工) 微分方程式の整数型解法における余剰解の整数数列の「解読」..... 15
 Fuminori Sakaguchi (Univ. of Fukui) 'Decipherment' of integer sequences corresponding to extra solutions in an integer-type algorithm for ODEs

概要 An integer-type algorithm for solving ODEs was proposed by the author and M. Hayashi. This algorithm is based on the expansion of solution functions by rational-function-type basis functions, and it is based on the 'exact' kernel vectors of non-square matrices. In this algorithm, we can read and 'decipher' integer coefficient sequences directly, and hence we can analyze behavior of numerical solutions exactly and pure-mathematically. In this study, the author gives further numerical examples where it is clearly shown that many hyperfunction components are contained in extra solutions mixed in numerical solutions obtained by this algorithm, by a direct 'decipherment' of integer coefficient sequences contained in numerical results by this algorithm, than the examples already presented.

- 34 湯 浅 智 意 (立命館大理工) Second order unbiased simulation method for reflected stochastic differ-
 秋 山 樹 生 (立命館大理工) ential equations 15
 A. Kohatsu-Higa (立命館大理工)
 Tomooki Yuasa (Ritsumeikan Univ.) Second order unbiased simulation method for reflected stochastic differ-
 Tatsuki Akiyama (Ritsumeikan Univ.) ential equations
 Arturo Kohatsu-Higa
 (Ritsumeikan Univ.)

概要 In this talk, we consider unbiased simulation methods functionals of solutions of one-dimensional reflected stochastic differential equations. Alfonsi-Hayashi-Kohatsu propose an unbiased simulation method for the present problem based on the parametrix method. However, the variance of this method is not finite in general unless one uses an importance sampling method. We propose a different way of obtaining what we call a second order parametrix method which leads to an alternative unbiased random variable with finite moments. We call this method "Second order unbiased simulation method for reflected stochastic differential equations".

- 35 木下 武彦 楕円型作用素に対する精度保証付き可逆性検証 .. 15
 渡部 善隆
 (九大情報基盤研究開発センター)
 中尾 充宏 (早大理工)
 Takehiko Kinoshita An invertibility verifications for linear elliptic operators mapping to dual
 Yoshitaka Watanabe (Kyushu Univ.) space
 Mitsuhiro T. Nakao (Waseda Univ.)

概要 This talk presents an computer-assisted procedure for verifying the invertibility of second-order linear elliptic operators and for computing a bound on the norm of its inverse. This approach is an improvement of a theorem (Nakao, et al. 2015, Jpn. J. Ind. Appl. Math. 32, 19–32) that uses projection and constructive a priori error estimates. Several examples which confirm the actual effectiveness of the procedure are reported on.

- 36 宮路 智行 (明大 MIMS) Proudman–Johnson 方程式の unimodal な定常解に対する精度保証付き
 岡本 久 (学習院大理) 数値計算 15
 Tomoyuki Miyaji (Meiji Univ.) Computer-assisted proof of the existence of a unimodal solution to the
 Hisahi Okamoto (Gakushuin Univ.) Proudman–Johnson equation

概要 We study the existence of unimodal stationary solutions to the Proudman–Johnson equation with an external force. In particular, we are interested in the case of a high Reynolds number. In order to prove the existence and unimodality of a solution, we resort to interval arithmetic. We formulate the stationary problem for the Proudman–Johnson equation as a system of first order ordinary differential equations, and we apply the shooting method and the interval Newton method for proving the existence of a solution. As the shooting method is numerically unstable, we encounter some difficulties especially when the Reynolds number is high. For solving this problem, we apply the multiple shooting method and the multiple-precision floating-point arithmetic.

14:15~16:40

- 37 上田 祐暉 (東大数理) 放物型問題の Nitsche’s method に対する inf-sup 条件及び誤差評価につ
 齊藤 宣一 (東大数理) いて 15
 Yuuki Ueda (Univ. of Tokyo) The inf-sup condition and error estimate of the Nitsche’s method for
 Norikazu Saito (Univ. of Tokyo) the parabolic problems

概要 We study the application of the Nitsche’s method to the parabolic problems. Under some assumptions, the parabolic initial-boundary value problem has a unique weak solution. The problem is discretized in space by the Galerkin method and the Dirichlet boundary condition is enforced weakly by the Nitsche’s method. It is well known that the bilinear form satisfies the Galerkin orthogonality. In this presentation, we will prove that the bilinear form also satisfies the inf-sup condition. This condition implies that the resulting semi-discretized problem has a unique solution. Moreover, the error estimate follows directly from the inf-sup condition and the Galerkin orthogonality.

- 38 小山 大介 (電通大)^b 重調和問題に対するある混合型非適合有限要素法への内部ペナルティ法
 の適用 15
 Daisuke Koyama An application of the interior penalty method to a mixed nonconforming
 (Univ. of Electro-Comm.) finite element method for biharmonic problems

概要 We introduce a numerical method for biharmonic problems which is obtained by applying the interior penalty method to a mixed nonconforming finite element method which is called the Hermann–Johnson (HJ) method. We show that a priori error estimates of our method can be the same as those of the HJ method by appropriately choosing a penalty parameter in our method.

- 39 劉 雪 峰 (新潟大 自然) 3次元領域における Stokes 方程式の有限要素解の事前誤差評価 15
 中 尾 充 宏 (早大理工・九大*)
 遊 春 光
 (China Acad. of Eng. Phys.)
 大 石 進 一 (早 大 理 工)
 Xuefeng Liu (Niigata Univ.) A priori error estimation for the finite element solution to Stokes equation in 3D domain
 Mitsuhiro Nakao
 (Waseda Univ./Kyushu Univ.*)
 Chun'guang You
 (China Acad. of Eng. Phys.)
 Shin'ichi Oishi (Waseda Univ.)

概要 For the Stokes equation defined in 3D domain with a general shape, the Scott–Vogelius finite elements are used to obtain strictly divergence-free approximation solution. Then, by using the hypercircle equation method, a quantitative a priori error estimation is obtained for the FEM solution. Such an a priori error estimation can be used in solution existence verification of nonlinear Navier–Stokes equation defined in 3D domain with general shapes. The convergence rate is confirmed by numerical results.

- 40 齊 藤 宣 一 (東 大 数 理) 不連続 Galerkin 時間離散化法の変分法的な解析 15
 Norikazu Saito (Univ. of Tokyo) Variational analysis of the discontinuous Galerkin time-stepping method

概要 The discontinuous Galerkin (DG) time-stepping method applied to abstract evolution equation of parabolic type is studied using a variational approach. We establish the inf-sup condition for the DG bilinear form. Then, the optimal order error estimates under appropriate regularity assumption on the solution are derived as direct applications of the standard interpolation error estimates. Our method of analysis is new. It differs from previous works on the DG time-stepping method by which the method is formulated as the one-step method.

- 41 剣 持 智 哉 (東 大 数 理) DG time-stepping 法に対する解析半群論的なアプローチ 15
 Tomoya Kemmochi (Univ. of Tokyo) An analytic semigroup approach for the DG time-stepping method

概要 The discontinuous Galerkin time-stepping method (DG time-stepping method) is a time-discretization method based on the discontinuous Galerkin finite element method. In contrast to one-step methods, the approximated solution is well-defined at each time in the DG time-stepping method. Therefore, it gives an efficient numerical algorithm with space-time methods for moving boundary problems such as fluid structure interaction. However, there are few studies on theoretical analysis for the behavior of approximated solutions at each time. In this talk, we address the DG time-stepping method for parabolic problems in the framework of analytic semigroup theory. We present optimal order error estimates for the homogeneous heat equation. The key point is rigorous estimates for rational functions that express the approximated solutions.

- 42 東 森 信 就 (京 大 CPIER) 熱方程式の初期値問題 (横向きを含む) に対する不安定差分スキームの数値計算例 15
 Nobuyuki Higashimori (Kyoto Univ.) Numerical examples of unstable finite difference schemes for the initial value problem of the heat equation including sideways case

概要 We previously reported examples of convergence of unstable finite difference schemes applied to quasilinear partial differential equations of the normal form with analytic initial data. This time we report that convergence also holds for initial value problems for the heat equation with initial data in a certain class of analytic functions and also for the sideways heat equation with initial data in a Gevrey class.

- 43 土屋 拓也 (早大理工) De Sitter 時空における半線形 Klein–Gordon 方程式の数値安定性について
中村 誠 (山形大理) 15
Takuya Tsuchiya (Waseda Univ.) On the numerical stability of semi-linear Klein–Gordon equations in de
Makoto Nakamura (Yamagata Univ.) Sitter spacetime

概要 We investigate the numerical stability for semi-linear Klein–Gordon equations in de Sitter spacetime. We show the differences of the numerical stability between some positive nonlinear terms and the negative one. In addition, the numerical stability in the four dimension and more is shown.

- 44 石川 歩 惟 (神戸大システム情報) 変分原理に基づくエネルギー保存数値解法の Lie 群上への拡張 15
谷口 隆 晴 (神戸大システム情報)
Ai Ishikawa (Kobe Univ.) The extension of the energy-preserving method based on the variational
Takaharu Yaguchi (Kobe Univ.) principle to the Lie group

概要 The energy-preserving method based on the variational principle and the discrete gradient method are methods for designing a scheme for the Hamilton equations that preserves the energy conservation law exactly. In the discrete gradient method, the various extensions, e.g. the extension to the equations on a manifold have been proposed. Although the method based on the variational principle has an advantage in the view of the computational cost, this method has not been applied to the equations on the manifold yet. In this talk, we extend the energy-preserving method based on the variational principle to equations on the Lie group and show a numerical test for the heavy top problem.

- 45 奥村 真善美 (阪大情報) Nonlinear and linear DVDM scheme for the conservative non-local Allen–
Cahn equation 15
Makoto Okumura (Osaka Univ.) Nonlinear and linear DVDM scheme for the conservative non-local Allen–
Cahn equation

概要 We propose nonlinear and linear finite difference scheme for the conservative non-local Allen–Cahn equation. Both proposed schemes inherit characteristic properties, the conservation of mass and the decrease of the global energy from the equation. We show that the schemes are stable in the sense that the numerical solution is bounded concerning max-norm, and have a unique solution. Since the nonlinear scheme is the system of equations concerning the new time step, it takes time to compute. Numerical examples demonstrate the effectiveness of the proposed scheme and that the computational time of the linear scheme is shorter than one of the nonlinear scheme. In this talk, we mainly introduce the linear scheme.

16:50～17:50 特別講演

- 吉川 周二 (大分大理工) 構造保存型差分法のエネルギー法
Shuji Yoshikawa (Oita Univ.) Energy method for structure-preserving finite difference schemes

概要 We introduce the energy method for structure-preserving finite difference schemes which inherit the physical structures such as energy conservation or dissipative laws. Another aim is to give some useful properties for difference quotient which is compatible with the structure-preserving finite difference schemes. The method and properties enable us not only to take the problem with more general nonlinearity but also to improve proofs of error estimate between the numerical and exact solutions. In this talk, after explaining our procedure by using a simple example, several our recent results are introduced.

3月21日(水) 第VI会場

9:15~11:55

- 46 渡辺雅二 (岡山大環境) Inverse analysis for microbial population in biodegradation process of
河合富佐子 xenobiotic polymer 15
 (京大工繊大ナノ材料・デバイス研究センター)
Masaji Watanabe (Okayama Univ.) Inverse analysis for microbial population in biodegradation process of
Fusako Kawai (Kyoto Inst. Tech.) xenobiotic polymer

概要 This study shows that monomer consumption and viability loss are the primary factors for microbial growth in exogenous type depolymerization processes. A mathematical model for the weight distribution and microbial population is described. Inverse problems for a time factor and a molecular factor of degradation rate are illustrated.

- 47 江夏洋一 (東京理大理) 自由境界をもつある感染症モデルについて 15
石渡恵美子 (東京理大理)
牛島健夫 (東京理大理工)
Yoichi Enatsu (Tokyo Univ. of Sci.) On an SIR model with free boundary
Emiko Ishiwata (Tokyo Univ. of Sci.)
Takeo Ushijima (Tokyo Univ. of Sci.)

概要 Free boundary problems are recently used to model phenomena of biological invasion for species such as migration into a new habitat (e.g., Du & Lin (2010) and references therein). These ideas are also applied to epidemic models. In this talk, we extend the result in Kaellen (2017) to the simple diffusive epidemic model with free boundary, namely we prove the existence of a semi wave solution. We numerically observe the semi wave and the front motion of this model with free boundary.

- 48 岡本 守 (北大理) 対流の効果を含む界面活性剤系自走粒子の自律運動に関する数理モデル
長山雅晴 (北大電子研) 15
Mamoru Okamoto (Hokkaido Univ.) Mathematical model including fluid's effect of camphor disk's self-motion
Masaharu Nagayama (Hokkaido Univ.)

概要 Many researchers have studied the self-driven particles. In one example, there is camphor atop water channel. It is now said that the motion of camphor is caused by differences in surface tension. The gradient of surface tension is induced by a camphor molecular layer development atop the surface. Mathematical models for the camphor motion have been constructed used the above mechanisms, and the models reproduce many characteristic motion. Although Marangoni Convection seems to influence the self-motion of the camphor, there are only a few reports discussing mathematical models that include convection explicitly. We have constructed a mathematical model for the self-motion of camphor including influence of convection, and now are calculating some cases to compare with some experiments.

- 49 後藤田剛 (北大電子研) 表皮構造の数値モデルにおける顆粒層の安定化とタイトジャンクション
 上坂正晃 (北大電子研) 形成への取り組み 15
 安ヶ平裕介 (北大理)
 小林康明 (お茶の水女大理)
 北畑裕之 (千葉大理)
 傳田光洋 ((株)資生堂)
 長山雅晴 (北大電子研)
- Takeshi Gotoda (Hokkaido Univ.) Mathematical modeling for stable formation of the granular layer and
Masaaki Uesaka (Hokkaido Univ.) tight junctions in the epidermis
Yusuke Yasugahira (Hokkaido Univ.)
Yasuaki Kobayashi (Ochanomizu Univ.)
Hiroyuki Kitahata (Chiba Univ.)
Mitsuhiro Denda
 (Shiseido Company, Ltd.)
Masaharu Nagayama (Hokkaido Univ.)

概要 We propose a mechanism of stable formation of the granular layer in the epidermis. In our mathematical model of the epidermis, we assume that a stimulant, which promotes the differentiation process of epidermal cells, is released when cells undergo cornification. We demonstrate that our model forms the granular layer and confirm that its layer structure is maintained stably by using some cost functions. We are also working on formation of tight junctions, which exist in the granular layer and play an important role in skin barrier function, and we introduce our trial models for them.

- 50 須志田隆道 (北大電子研) 基底膜に覆われた細胞群の回転運動を表現するための数値モデル 15
 森 瞳 (北大理)
 石田すみれ (北大生命)
 古澤和也 (北大先端生命)
 芳賀 永 (北大先端生命)
 秋山正和 (北大電子研)
- Takamichi Sushida (Hokkaido Univ.) A mathematical model for representing collective rotational migrations
Hitomi Mori (Hokkaido Univ.) of cell groups covered by basement membrane
Sumire Ishida (Hokkaido Univ.)
Kazuya Furusawa (Hokkaido Univ.)
Hisashi Haga (Hokkaido Univ.)
Masakazu Akiyama (Hokkaido Univ.)

概要 Recently, it has been attended the relation between the elongation phenomena and rotational migration of cell group since they are observed in three-dimensional morphogenesis such as fruiting body formation of *Dictyostelium Discoideum* and somite formation of zebra fish. In particular, although it is known that the somite is covered by basement membrane, it is not clear the reason why rotational migration occurs. In order to understand cellular mechanism for rotational migration of cell groups covered by basement membrane, we propose a mathematical model which consists of a self-propelled particle model representing cellular migration and a phase-field model representing basement membrane. Moreover, we will show phase diagram of parameters for migration modes and give a theoretical suggestion for biological experiments.

- 51 佐藤 峻 (東大情報理工) 混合微分を含む発展方程式の再定式化について 15
 Shun Sato (Univ. of Tokyo) Reformulation of evolutionary differential equations with a mixed derivative

概要 In this talk, we deal with the initial value problem of evolutionary differential equations with a mixed derivative on the periodic domain. Here, “mixed derivative” indicates the case where a spatial differential operator is operating on the time derivative, obscuring the vector field describing the flow. Therefore, some reformulation to reveal it is the first step of PDE-theoretical and numerical studies. However, it is nontrivial because the spatial differential operator is not invertible and cannot be easily eliminated. Though this issue was already settled for linear cases, general theory has been undeveloped. In this talk, we propose a novel procedure for wider class of equations. Moreover, as an application, we establish the global well-posedness of the sine-Gordon equation in characteristic coordinates.

- 52 清水 雄貴 (京大理) 対称曲面上の Green 関数 15
 Yuuki Shimizu (Kyoto Univ.) Green’s function on surfaces with symmetry

概要 Uniform flow is one of fundamental steady solutions of Euler equation on a plane. A generalization of the flow on curved surfaces is a Killing vector field, which is also a steady solutions of Euler equation on Riemannian manifolds. In this talk, we introduce how surfaces has a no-normal regular Killing vector field and as its application, construction of an analytic formula of Green’s function on the surface.

- 53 坂上 貴之 (京大理) 非粘性保存量の乱流カスケードを実現する 1 次元流体方程式について .. 15
 松本 剛 (京大理)
 Takashi Sakajo (Kyoto Univ.) One dimensional hydrodynamic PDE generating turbulent cascade of
 Takeshi Matsumoto (Kyoto Univ.) inviscid invariant

概要 We propose a one-dimensional hydrodynamic partial differential equation. This model is based on a Constantin–Lax–Majda–De Gregorio model generalized by Okamoto, Sakajo and Wunsch. The equation admits an inviscid invariant quantity. In the presence of the viscosity and a large-scale random forcing, the solution gives rise to a turbulent state with cascade of the inviscid invariant. We will give how those phenomenon is understood from the view point dynamical system.

- 54 坂上 貴之 (京大理) Kasper 翼まわりの定常点渦の安定化を実現する線型フィードバック制御
 R. Nelson 15
 (Imperial Coll. London)
 B. Protas (McMaster Univ.)
 Takashi Sakajo (Kyoto Univ.) Linear feedback control stabilizing point vortex equilibria near a Kasper
 Rhodri Nelson (Imperial Coll. London) Wing
 Bartosz Protas (McMaster Univ.)

概要 Based on the Linear-Quasi-Gaussian compensator, we design a linear feedback system stabilizing point vortex equilibria near an aerodynamic wing with two auxiliary flaps known as a Kasper Wing in the presence of a uniform flow. This is modeled by a two-dimensional incompressible an inviscid flow. The actuation mechanism is blowing and suction localized on the main plate represented as a sink- source singularity, whereas we measure pressure across the plate as system output. We show that the linearised system around these equilibria are both controllable and observable for almost all actuator and sensor locations. Numerical computations illustrate that Kasper Wing configurations are in general not only more controllable than their single plate counterparts, but also acquire larger basins of attraction owing to the feedback control.

- 55 大塚 厚二 (広島国際学院大情報文化) 一般 J 積分による固有値の形状感度解析 15
 Kohji Ohtsuka (Hiroshima Kokusai Gakuin Univ.) Shape sensitivity analysis of eigenvalue by generalized J-integral in boundary value problems

概要 Generalized J-integral is the tool which is effective to study the shape optimization of singular points (containing boundary) with respect to various cost functions, energy, mean compliance, least square errors, in boundary value problems for partial differential equations. I will talk an application of Generalized J-integral method to shape sensitivity of eigenvalue problems.

14:15~16:40

- 56 上田 肇一 (富山大理) 自発的なパラメータ制御による集団振動子の同期現象 15
 Keiichi Ueda (Univ. of Toyama) Synchronization of two interacting populations of oscillators by autonomous parameter control

概要 We study synchronization of two interacting populations of oscillators. We assume that the sign and the strength of the interactions are taken as system variables, and that the variables are determined by a function of the oscillator variables. Under an appropriate choice of the function, we can observe spontaneous intra- and inter-group phase synchronization.

- 57 渡辺 毅 (東大工) 振動テールパルスと非一様媒質の相互作用を支配する大域的分岐構造 .. 15
 Zhijun Gao (東北大 AIMR) 西浦 廉政 (東北大 AIMR)
 Takeshi Watanabe (Univ. of Tokyo) Zhijun Gao (Tohoku Univ.) Yasumasa Nishiura (Tohoku Univ.)
 Global bifurcation structure governing interaction between bump heterogeneity and pulse with oscillatory tail

概要 Three-component FitzHugh–Nagumo equation is investigated. This equation has a parameter region in which traveling pulse with oscillatory tail appears. When such moving pulse interacts with heterogeneity of the media, the interaction between tail and heterogeneity is important. We consider a bump-type heterogeneity and investigate the asymptotic behavior of the pulse motion when a pulse collides with the bump. When the width of the bump is fixed and the height of that is changed, three different asymptotic behaviors arise sequentially: Oscillatory pinning (OSC), stationary pinning (STA), and rebound (REB). In these asymptotic states, it is conjectured that all asymptotic states are contained in the set of bifurcation branches which is generated from the trivial branch.

- 58 西 慧 (京都産大理) 円環水路上の 2 個の樟脳ろ紙がみせる運動の分岐解析 15
 長山 雅晴 (北大電子研) 中田 聡 (広島大理)
 Kei Nishi (Kyoto Sangyo Univ.) Masaharu Nagayama (Hokkaido Univ.) Satoshi Nakata (Hiroshima Univ.)
 Bifurcation analysis of the motion of two self-propelled camphor disks on an annular field

概要 The dynamics of two self-propelled camphor disks on an annular field is explored both numerically and analytically. In our previous study, it was already found by the direct numerical simulation of a model equation that the two camphor disks exhibited a variety of behavior, and underwent transition between the behavior as the length of the annular field was varied. In order to analytically elucidate the mechanism for the transition in behavior, we reduced the model equation which consisted of two ODEs and one PDE into three ODEs for the motion of the two camphor disks. In this talk, the bifurcation structure that causes the transition will be revealed, based on the reduced ODEs.

- 59 山中祥五 (京大情報) 2自由度ハミルトン系における横断的なヘテロクリニック軌道の存在と非
矢ヶ崎一幸 (京大情報) 可積分性 15
Shogo Yamanaka (Kyoto Univ.) Existence of transverse heteroclinic orbits and nonintegrability in two-
Kazuyuki Yagasaki (Kyoto Univ.) degree-of-freedom Hamiltonian systems with saddle-centers

概要 We consider a class of two-degree-of-freedom Hamiltonian systems with saddle-centers connected by heteroclinic orbits. We show that if the sufficient conditions for real-meromorphic nonintegrability hold, then the stable and unstable manifolds of the periodic orbits intersect transversely, are quadratically tangent or do not intersect in general, and they do not intersect when the Hessian matrix of the Hamiltonian has a different number of positive eigenvalues at the associated saddle-centers. Our theory is illustrated for a system with quartic single-well potential.

- 60 中野直人 導関数座標埋め込みによって得られる曲面の性質と函数の時間発展の予
(京大国際高等教育院・JSTさきがけ) 測可能性について 15
Naoto Nakano Characteristics of derivative embedded surfaces and predictability of
(Kyoto Univ./JST PRESTO) timeseries

概要 Here, we consider the derivative embedding of a scalar function. Since the embedded surface can hold some characteristics of the original time-series, one can derive some predictability result from characteristics of the embedded surface.

- 61 佐々木多希子 (明大理工) 1次元非線形波動方程式の爆発曲線に関する数値・数学解析 15
石渡哲哉
(芝浦工大システム理工)
Takiko Sasaki (Meiji Univ.) Numerical and mathematical analysis for the blow-up curve of solutions
Tetsuya Ishiwata to 1-dimensional nonlinear wave equations
(Shibaura Inst. of Tech.)

概要 We study a blow-up curve for the one dimensional wave equation $\partial_t^2 u - \partial_x^2 u = 2^p |\partial_t u|^p$ with the Dirichlet boundary condition. The purpose of this talk is to show that the blow-up curve T satisfies that $T'(x) \rightarrow -1$ as $x \rightarrow 0+0$ (1) under the suitable initial conditions. To prove the result, we convert the equation into a first order system, and then present some numerical investigations of the blow-up curves. From the numerical results, we were able to confirm (1) holds numerically. Moreover, under some assumptions, we were also able to confirm (1) holds mathematically.

- 62 穴田浩一 (早大高等学院) ある準線形放物型偏微分方程式の後方自己相似解に関する一考察 15
石渡哲哉
(芝浦工大システム理工)
牛島健夫 (東京理大理工)
Koichi Anada A study for backward self similar solutions of a quasi-linear parabolic
(Waseda Univ. Senior High School) equation
Tetsuya Ishiwata
(Shibaura Inst. of Tech.)
Takeo Ushijima (Tokyo Univ. of Sci.)

概要 In this talk, we consider properties of backward self similar solutions for a quasi-linear parabolic equation $v_t = v^\delta (v_{xx} + v)$. Their properties are very important to investigate asymptotic behavior of solutions to this parabolic equation, especially, the blow-up sets and rates.

- 63 松江 要 無限遠ダイナミクスが導く解の爆発レート 15
 (九大IMI・九大I2CNER)
 Kaname Matsue Blow-up rates of blow-up solutions determined by dynamics at infinity
 (Kyushu Univ./Kyushu Univ.)

概要 Geometric treatments of blow-up solutions for autonomous ordinary differential equations and their blow-up rates are concerned. Our approach focuses on the type of invariant sets at infinity via compactifications of phase spaces, and dynamics on their center-stable manifolds. In particular, we show that dynamics on center-stable manifolds of invariant sets at infinity with appropriate time-scale desingularizations as well as blowing-up of singularities characterize dynamics of blow-up solutions as well as their rigorous blow-up rates not only of so-called “type-I” but also other types.

- 64 松江 要 速いレートで振る舞う振動爆発解と, 振動発散解 15
 (九大IMI・九大I2CNER)
 Kaname Matsue Oscillatory blow-up solutions with fast blow-up rates and oscillatory
 (Kyushu Univ./Kyushu Univ.) grow-up solutions

概要 Geometric treatments of oscillatory blow-up solutions for autonomous ordinary differential equations and their blow-up rates are concerned. As in the preceding talk, we apply compactification of phase spaces and time-scale desingularization to characterization of blow-up solutions. In particular, when divergent solutions are characterized by trajectories on center manifolds of non-hyperbolic periodic orbits on the horizon for desingularized vector fields, they blow up in finite time with infinitely fast oscillation and faster blow-up rate than type-I rates in typical cases, while they can also grow up in infinite time in some cases. We see such behavior in a certain system known as Liénard equation.

16:50~17:50 特別講演

- 高安亮紀 (筑波大システム情報) 半群理論を利用する発展方程式に対する解の精度保証付き数値計算
 Akitoshi Takayasu (Univ. of Tsukuba) Verified computations for solutions of evolution equations with semi-
 group theory

概要 In this talk, we provide a methodology of verified computing for solutions to evolution equations (nonlinear heat equations, 1-dimensional advection equations with variable coefficients, and the complex Ginzburg–Landau equations). Our methodology is based on semigroup theory, which is widely used in analytical studies and originated from pioneering works by Hille and Yosida. The main contribution of this study is to combine a “classical analysis” with “computer-assisted methods” to provide a numerical method of enclosing a solution for evolution equations. The combination of quantitative estimates arising from verified numerical computations and qualitative results obtained by classical analysis is expected to open the access to many unsolved problems by purely analytical means.

トポロジー

3月18日(日) 第II会場

9:40~12:00

- 1 岩山 瑛子 n 次元単体の構成とオイラーポアンカレ標数の拡張と組合せの数 5
Eiko Iwayama The simplex of n -dim. and New Euler–Poincaré Expanded characteristic and the relation of combination numbers

概要 I have constructed the simplex MS_n of n -dim. The alternative sum of numbers of k -dim-sub-simplexes MS_n^k of n -simplex MS_n are New Euler–Poincaré Expanded characteristic “NEPE”. I have had $NEPE = 1$ for all dim. n . I have obtained the table like “Pascal’s triangle” for M_n^k ($k = 0, 1, \dots, n$) and we also have $M_n^k = {}_{n+1}C_{k+1}$ (${}_{n+1}C_{k+1}$: combinatorial numbers).

- 2 岩山 瑛子 n 次元直方体の三角形とオイラーポアンカレ標数の拡張 5
Eiko Iwayama The triangle of the cuboids of n -dimension and New Euler–Poincaré Expanded characteristic

概要 I have constructed the cuboid M_n of n -dim. The alternative sum of numbers of k -dim. sub-cuboids M_n^k are “New Euler–Poincaré Expanded characteristic NEPE”. I have had $NEPE = 1$ for all dim. n . We can connect 4 dim. cuboid with hyper planes. If the complex of cubes has 2-holes, then $NEPE = -1$.

- 3 小笠原 義仁 (早大複雑系高等研) トポロジーが持つ応用可能性について 10
Yoshihito Ogasawara (Waseda Univ.) On an applicability of topology

概要 New possibility of topological application is explored. In this study, topology is not only regarded as a system of morphological concepts, but also tried to be interpreted as the discussion of conceptual morphology. This means that our concepts themselves are tried to be described by the topology which is a system of our concepts, as mathematical system itself is described by mathematics in the field of foundations of mathematics.

Then, new picture of the world is tried to be drawn by such a sense of topology. The world is not regarded as a priori existence, but tried to be described as the emergence from our recognition.

- 4 直江 央寛 (東北大理) Infinitely many corks having large shadow-complexities 15
Hironobu Naoe (Tohoku Univ.) Infinitely many corks having large shadow-complexities

概要 Any exotic pair of simply connected closed 4-manifolds are related by a cork twist. Every 4-manifold can be represented by a simple polyhedron with a coloring on each region, called a shadow. Using shadows of 4-manifolds, Costantino defined a complexity of a 4-manifold, which is the minimum number of true vertices of its shadow. We have known many examples of corks having low complexities. In this talk, we will show that there also exist infinitely many corks with large complexity.

- 5 宮村 旭 (東工大 理) 種数0のLefschetzファイバー空間の符号数について 10
Akira Miyamura (Tokyo Tech) A note on signature of Lefschetz fibrations with planar fiber

概要 In this talk, we present a signature formula for allowable Lefschetz fibrations over D^2 with planar fiber by computing Maslov index appearing in Wall’s non-additivity formula.

- 6 稲垣 友介 (阪大 理) パンツの Hitchin 成分の Fuchs 跡について 15
 Yusuke Inagaki (Osaka Univ.) On Fuchsian loci of Hitchin components of a pair of pants

概要 Hitchin components are the connected components of character varieties of surface groups containing Teichmüller spaces, and the subsets of Hitchin components which correspond to Teichmüller spaces are called Fuchsian loci. Recently Bonahon–Dreyer constructed a parameterization of $\mathrm{PSL}_n(\mathbb{R})$ -Hitchin components by using the Anosov property of elements of $\mathrm{PSL}_n(\mathbb{R})$ -Hitchin components and invariants of flags introduced by Fock–Goncharov, which is a parameterization by Euclidian convex polytopes. In this talk, we give an explicit description of Fuchsian loci of a pair of pants by using the Bonahon–Dreyer parameterization.

- 7 田所 勇樹 (木更津工高専) Pointed harmonic volume and its relation to extended Johnson homomorphism 15
 Yuuki Tadokoro Pointed harmonic volume and its relation to extended Johnson homomorphism
 (Nat. Inst. of Tech., Kisarazu Coll.)

概要 As a natural extension of the period, the pointed harmonic volume for a compact Riemann surface is defined using Chen’s iterated integrals. It captures more detailed information of the complex structure. It is also one of a few explicitly computable examples of complex analytic invariants. We obtain its new value for a certain pointed hyperelliptic curve. An application of the pointed harmonic volume is presented. We explain the relationship between the pointed harmonic volume and first extended Johnson homomorphism on the mapping class group of a pointed oriented closed surface.

- 8 山本 卓宏 (東京学大教育) 境界付き多様体上のモース関数の同境界群 15
 Takahiro Yamamoto Cobordism groups of Morse functions on manifolds with boundary
 (Tokyo Gakugei Univ.)

概要 In this talk, we discuss whether cobordism groups of Morse functions on manifolds with boundary are trivial or not.

- 9 土田 旭 (北大 理) 接分布と同階数ベクトル束の間の束準同型の特異点 15
 佐治 健太郎 (神戸大 理)
 Asahi Tsuchida (Hokkaido Univ.) Singularities of bundle homomorphism between a distribution and a
 Kentaro Saji (Kobe Univ.) vector bundle

概要 We consider singularities of bundle homomorphisms from a tangent distribution and a vector bundle of the same rank. Generic classification of the singularities for low dimensional cases are studied. We also consider a bundle homomorphism which is induced from a Morin map. In the case a distribution is a contact structure, we give a characterization of singularities of the bundle homomorphisms by using the contact Hamiltonian vector field.

- 10 佐治 健太郎 (神戸大 理) スワローテイルの $SO(3)$ -標準形と幾何的模様 10
 Kentaro Saji (Kobe Univ.) $SO(3)$ -normal form of swallowtail and geometric patterns

概要 We construct a form of swallowtail singularity in \mathbf{R}^3 which uses coordinate transformations on the source and isometries on the target. As an application, we classify configurations of asymptotic curves and characteristic curves near swallowtail.

14:15~15:15 特別講演

岸本大祐 (京大理) ポリヘドラル・プロダクトのホモトピー論
Daisuke Kishimoto (Kyoto Univ.) Homotopy theory of polyhedral products

概要 A polyhedral product is a space constructed combinatorially from a given abstract simplicial complex. Its homotopy invariants like cohomology give important combinatorially defined algebras such as Stanley–Reisner rings, and it is also important in toric topology. So there have been considerable efforts to develop the homotopy theory of polyhedral products in view of both combinatorics and topology. But there is no general technique to develop the homotopy theory of polyhedral products until the fat wedge filtration has been introduced. I will present a survey of the recent development of the homotopy theory of polyhedral products based on the fat wedge filtration and its applications. This talk is based on the joint work of Kouyemon Iriye (Osaka Prefecture University).

15:30~18:00

- 11 山崎正之 (岡大理) 非球面的曲面における正則閉曲線の回転数について 15
Masayuki Yamasaki On rotation numbers of regular closed curves on aspherical surfaces
(Okayama Univ. of Sci.)

概要 I define a rotation number of a regular closed curve on a complete euclidean/hyperbolic surface, which, together with the free homotopy class, determines a regular homotopy class. I also give a Whitney-type formula for this rotation number.

- 12 栗林勝彦 (信大理) 階層体の圏と Serre–Swan の定理 15
青木稔樹
Katsuhiko Kuribayashi (Shinshu Univ.) On the category of stratifolds and the Serre–Swan theorem
Toshiki Aoki

概要 Stratifolds are considered from a categorical point of view. We show among others that the category of stratifolds fully faithfully embeds into the category of \mathbb{R} -algebras as does the category of smooth manifolds. We prove that a variant of the Serre–Swan theorem holds for stratifolds. In particular, the category of vector bundles over a stratifold is shown to be equivalent to the category of vector bundles over an associated affine scheme although the latter is in general larger than the stratifold itself.

- 13 松岡拓男 Higher theories of algebraic structures 15
Takuo Matsuoka Higher theories of algebraic structures

概要 Discovery or recognition of the right kind of *algebraic structure* is often important in the development of mathematical subjects. In situations where various complex kinds of algebraic structure can arise, special technology for systematically finding and treating algebraic structures would be desirable. In particular, such technology would be necessary for broad application of *higher category theory*, since algebraic structures of high categorical dimension are varied and can be complicated. We shall describe how concrete understanding of higher categorical *coherence* leads to a systematic view on some (quite general) kinds of algebraic structure. A consequently found new phenomenon concerning *topological field theories* is interesting in its contrast to the *cobordism hypothesis*.

- 14 林晋 (MathAM-OIL) バルクエッジ対応と指数のコボルディズム不変性 15
Shin Hayashi (MathAM-OIL) Bulk-edge correspondence and the cobordism invariance of the index

概要 We show that the bulk-edge correspondence for two-dimensional type A topological insulators follows directly from the cobordism invariance of the index.

- 15 林 晋 (MathAM-OIL) バルクエッジコーナー対応 15
 Shin Hayashi (MathAM-OIL) Bulk-edge and corner correspondence

概要 We consider a translation invariant bounded linear self-adjoint operator (model of a Hamiltonian) on a three-dimensional lattice (bulk) and its restrictions onto two subsemigroups (edges) and their intersection (corner). We first show that, if our bulk and edges Hamiltonians have a common spectral gap, we can define a topological invariant for the gapped bulk and edges. We next show a relation between this invariant and another invariant defined for the corner.

- 16 森谷 駿二 (阪 府 大) The space of knots in a manifold and the right operadic module of configuration spaces 15
 Syunji Moriya (Osaka Pref. Univ.) The space of knots in a manifold and the right operadic module of configuration spaces

概要 For closed simply connected manifold M of dimension ≥ 4 , we introduce a new spectral sequence converging to the space of knots in M .

- 17 森谷 駿二 (阪 府 大) Non-formality of the odd dimensional framed little disks operads 15
 Syunji Moriya (Osaka Pref. Univ.) Non-formality of the odd dimensional framed little disks operads

概要 We prove the odd dimensional framed little disks operads is not formal as a non-symmetric operad.

- 18 鎌田 英也 (近畿大工高専) n -cube の頂点カット上のトーリック多様体の分類 15
 蓮井 翔 (阪府大高等教育)
 柘田 幹也 (阪市大理)
 朴 善正 (阪市大理)
 Hideya Kuwata (Kindai Univ. Tech. Coll.) Classification of toric manifolds over an n -cube with one vertex cut
 Sho Hasui (Osaka Pref. Univ.)
 Mikiya Masuda (Osaka City Univ.)
 Seonjeong Park (Osaka City Univ.)

概要 A complete nonsingular toric variety (called a toric manifold) is over P if its quotient by the compact torus is homeomorphic to P as a manifold with corners. Bott manifolds are toric manifolds over an n -cube I^n and blowing them up at a fixed point produces toric manifolds over $\text{vc}(I^n)$ an n -cube with one vertex cut. They are all projective. On the other hand, Oda's 3-fold, the simplest non-projective toric manifold, is over $\text{vc}(I^3)$. In this paper, we classify toric manifolds over $\text{vc}(I^n)$ ($n \geq 3$) as varieties and as smooth manifolds. It consequently turns out that there are many non-projective toric manifolds over $\text{vc}(I^n)$ but they are all diffeomorphic, and toric manifolds over $\text{vc}(I^n)$ in some class are determined by their cohomology rings as varieties.

- 19 阿部 拓 (阪市大数学研) 正則半単純なヘッセンバーグ多様体のコホモロジー環について 10
堀口 達也
(阪大情報・阪市大数学研)
栞田 幹也 (阪市大理)
Hiraku Abe (Osaka City Univ.) On the cohomology rings of regular semisimple Hessenberg varieties
Tatsuya Horiguchi
(Osaka Univ./Osaka City Univ.)
Mikiya Masuda (Osaka City Univ.)

概要 We investigate the cohomology rings of regular semisimple Hessenberg varieties whose Hessenberg functions are of the form $h = (h(1), n \cdots, n)$ in Lie type A_{n-1} . The main result gives an explicit presentation of the cohomology rings in terms of generators and their relations. Our presentation naturally specializes to Borel's presentation of the cohomology ring of the flag variety, and it is compatible with the representation of the symmetric group on the cohomology constructed by J. Tymoczko.

3月19日(月) 第II会場

9:30~10:30

- 20 森 淳 秀 (大阪歯大歯) t分布のシンプレクティック幾何・接触幾何 15
Atsuhide Mori (Osaka Dental Univ.) Symplectic/contact geometry of t-distributions

概要 In the previous work, the speaker studied the positive and negative symplectic structures on the space of the pairs of normal distributions and found a Lagrangian submanifold with nice properties. In this talk, we extend this result to the space of the pairs of t-distributions and propose an application concerning the smoothness of the movement of a parameter.

- 21 片山 拓 弥 (広島大理) Embeddability of the right-angled Artin groups on the complement graphs of linear forests 15
Takuya Katayama (Hiroshima Univ.) Embeddability of the right-angled Artin groups on the complement graphs of linear forests

概要 In 2011, Sang-hyun Kim and Thomas Koberda proved that, for any finite graphs Λ and Γ , a full embedding of Λ into the extension graph Γ^e of Γ gives rise to an embedding between the corresponding right-angled Artin groups, $A(\Lambda) \hookrightarrow A(\Gamma)$. Then the following natural question arises: for which graphs Λ and Γ , can we reduce an embedding $A(\Lambda) \hookrightarrow A(\Gamma)$ into a full embedding $\Lambda \rightarrow \Gamma^e$? Recently, several authors proved that the reduction is impossible for some Λ and Γ . In this talk, we give a positive answer when Λ is the complement graph of a linear forest. In addition, we can further reduce an embedding $A(\Lambda) \hookrightarrow A(\Gamma)$ into a full embedding between the defining graphs, $\Lambda \rightarrow \Gamma$, if Λ is the complement graph of a linear forest.

- 22 加藤 本 子 (東大数理) 符号付きトンプソン群の同型問題 10
Motoko Kato (Univ. of Tokyo) On the isomorphism problem of signed Thompson groups

概要 Higman–Thompson groups are groups of homeomorphisms of the Cantor space which are locally orientation preserving. They are examples of finitely presented virtually simple groups. Generalizing these groups, Funar and Neretin defined signed Higman–Thompson groups. Signed Higman–Thompson groups are groups of homeomorphisms of the Cantor space which are locally orientation preserving or orientation reversing. In this talk, we give a necessary and sufficient condition for a signed Higman–Thompson group to be isomorphic to one of Higman–Thompson groups. This is based on a joint work with Javier Aramayona and Julio Aroca (Autonomous University of Madrid).

- 23 小 沢 登 高 (京大数理研) Kazhdan's property (T) and semidefinite programming 15
 Narutaka Ozawa (Kyoto Univ.) Kazhdan's property (T) and semidefinite programming

概要 It has been known that Kazhdan's property (T) is semi-decidable and an algorithm to detect property (T) has been proposed. In this talk, I will describe an improved algorithm that exploits the symmetry on the given test group G . The improved algorithm makes computer verification of property (T) for certain groups possible otherwise impossible. I will report the result of a large-scale calculation. This talk is based on a joint work with M. Kaluba and P. Nowak.

10:45~11:45 特別講演

浅 岡 正 幸 (京 大 理) 滑らかな力学系の周期点の個数の増大度

Masayuki Asaoka (Kyoto Univ.) Growth rate of the number of periodic points for smooth dynamical systems

概要 The exponential rate of the growth rate of the number of periodic points is an important invariant of a dynamical systems. For example, it determines the convergence radius of the dynamical zeta function of the system and, for hyperbolic dynamics, it determines the topological entropy. It is natural to ask whether 'most' of smooth dynamical systems exhibit at most exponential growth of the number of periodic points or not. Some classical results showed that systems in a dense subset of the set of smooth maps in general dimensions and all real-analytic one-dimensional systems exhibit such tame growth. However, in 2000, Kaloshin proved that super-exponential growth is 'abundant' in smooth dynamics, and recently, the author found 'abundant' examples in real-analytic dynamics. In the first part of the talk, we survey classical results on at most exponential growth for tame cases, including hyperbolic systems. In the second part, we discuss contemporary results for wild cases.

13:00~14:20

- 24 川 口 徳 昭 (東大数理)^b On the shadowing and limit shadowing properties 15
 Noriaki Kawaguchi (Univ. of Tokyo) On the shadowing and limit shadowing properties

概要 In this talk, we consider the relation between the shadowing property and the limit shadowing property of topological dynamical systems. We show that for any continuous self-map f of a compact metric space, if f has the limit shadowing property, then the restriction of f to the non-wandering set satisfies the shadowing property. As an application, we prove the equivalence of the two shadowing properties for equicontinuous maps.

- 25 橋 本 忍 (首都大東京理工) 3次接触をもつ2次元微分同相写像の moduli 15
 Shinobu Hashimoto Moduli of two-dimensional diffeomorphisms with cubic tangencies
 (Tokyo Metro. Univ.)

概要 We study conjugacy invariants for 2-dimensional diffeomorphisms with homoclinic cubic tangencies (two-sided tangencies of the lowest order) under certain open conditions. Ordinary arguments used in past studies of conjugacy invariants associated with one-sided tangencies do not work in the two-sided case. In this talk, we will present a new method which is applicable to the two-sided case.

- 26 平 出 耕 一 (愛媛大理)^b 位相的エントロピーとリアプノフ指数の関係 15
 松 岡 千 博 (阪市大工)

Koichi Hiraide (Ehime Univ.) Relationship between topological entropy and Lyapunov exponents
 Chihiro Matsuoka (Osaka City Univ.)

概要 In this talk we discuss the relationship between topological entropy and Lyapunov exponents, provided that the Pesin entropy formula holds, and mention some results by experimental mathematics.

- 27 山崎 薫里 (高崎経大経済) Questions on monotone operators for vector-valued maps 15
 Kaori Yamazaki Questions on monotone operators for vector-valued maps
 (Takasaki City Univ. of Econ.)

概要 Answering a question of Yang, we show that, for an ordered topological vector space Y with positive interior points, if each non-zero positive element is an order unit, then Y is isomorphic to the real line. We also provide a technique which reduces some vector-valued results to the original real-valued ones by using some Minkowski functionals.

- 28 矢島 幸信 (神奈川大工) Three embeddings and their implications in products of generalized met-
 平田 康史 (神奈川大工) ric spaces 15
 Yukinobu Yajima (Kanagawa Univ.) Three embeddings and their implications in products of generalized met-
 Yasushi Hirata (Kanagawa Univ.) ric spaces

概要 We discuss when C^* -embedding or C -embedding implies P -embedding in products of generalized metric spaces, such as M -spaces, Σ -spaces and semi-stratifiable spaces.

3月20日(火) 第II会場

9:20~12:00

- 29 鄭 仁大 (近畿大理工) Achiral 1-cusped hyperbolic 3-manifolds not coming from amphicheiral
 市原 一裕 (日大文理) null-homologous knot complements 15
 谷山 公規 (早大教育)
 In Dae Jong (Kindai Univ.) Achiral 1-cusped hyperbolic 3-manifolds not coming from amphicheiral
 Kazuhiro Ichihara (Nihon Univ.) null-homologous knot complements
 Kouki Taniyama (Waseda Univ.)

概要 It is experimentally known that achiral hyperbolic 3-manifolds are quite sporadic at least among those with small volume, while we can find plenty of them as amphicheiral knot complements in the 3-sphere. In this talk, we show that there exist infinitely many achiral 1-cusped hyperbolic 3-manifolds not homeomorphic to any amphicheiral null-homologous knot complement in any closed achiral 3-manifold.

- 30 市原 一裕 (日大文理) 対掌矯飾の手術とキャッソン不変量 15
 伊藤 哲也 (阪大理)
 斎藤 敏夫 (上越教育大)
 Kazuhiro Ichihara (Nihon Univ.) Chirally cosmetic surgeries and Casson invariants
 Tetsuya Itoh (Osaka Univ.)
 Toshio Saito (Joetsu Univ. of Edu.)

概要 I will report on our recent study of chirally cosmetic surgery, that is, a pair of Dehn surgeries on a knot producing homeomorphic 3-manifolds with opposite orientations. Several constraints on knots and surgery slopes to admit such surgeries are given. Our main ingredients are the original and the $SL(2, C)$ version of Casson invariants. As an application, we give a complete classification of chirally cosmetic surgeries on two bridge knots of genus one.

- 31 阪田直樹 (広島大理) 双曲的ファイバー二橋絡み目補空間の標準的分割の veering 構造 10
 Naoki Sakata (Hiroshima Univ.) Veering structures of the canonical decompositions of hyperbolic fibered two-bridge link complements

概要 Epstein–Penner has proved that each finite-volume cusped complete hyperbolic manifold admits a canonical decomposition into ideal polyhedra. I proved that the canonical decompositions of hyperbolic fibered two-bridge link complements are layered with respect to the fiber structures. On the other hand, Agol has shown that every pseudo-Anosov mapping torus of a surface, punctured along the singular points of the stable and unstable foliations, admits a canonical “veering” layered triangulation. In this talk, we completely determine, for each hyperbolic fibered two-bridge link, whether the canonical decomposition of its complement is veering with respect to the fiber structure.

- 32 吉田建一 (京大理) タングルの貼り合わせにおける双曲性 15
 Ken'ichi Yoshida (Kyoto Univ.) Hyperbolicity on tangle gluings

概要 Let us glue a 3-manifold and the complement of a trivial tangle along bounding 4-punctured spheres. We propose that this construction can be regarded as an analogue of a Dehn filling. We will show an analogous result for Thurston’s hyperbolic Dehn surgery theorem.

- 33 R. Blair Height, trunk and representativity of knots 15
 (California State Univ.)
 小沢 誠 (駒澤大総合)
 Ryan Blair (California State Univ.) Height, trunk and representativity of knots
 Makoto Ozawa (Komazawa Univ.)

概要 In this talk, we investigate three geometrical invariants of knots, the height, the trunk and the representativity. First, we give a counterexample for the conjecture which states that the height is additive under connected sum of knots. Next, we show that the representativity is bounded above by a half of the trunk. We also define the trunk of a tangle and show that if a knot has an essential tangle decomposition, then the representativity is bounded above by half of the trunk of either of the two tangles. Finally, we remark on the difference among Gabai’s thin position, ordered thin position and minimal critical position. We also give an example of a knot which bounds an essential non-orientable spanning surface, but has arbitrarily large representativity.

- 34 R. Blair The incompatibility of crossing number and bridge number for knot diagrams 15
 (California State Univ.)
 A. A. Kjachukova
 (Univ. of Pennsylvania)
 小沢 誠 (駒澤大総合)
 Ryan Blair (California State Univ.) The incompatibility of crossing number and bridge number for knot diagrams
 Alexandra A. Kjachukova
 (Univ. of Pennsylvania)
 Makoto Ozawa (Komazawa Univ.)

概要 We define and compare several natural ways to compute the bridge number of a knot diagram. We study bridge numbers of crossing number minimizing diagrams, as well as the behavior of diagrammatic bridge numbers under the connected sum operation. For each notion of diagrammatic bridge number considered, we find crossing number minimizing knot diagrams which fail to minimize bridge number. Furthermore, we construct a family of minimal crossing diagrams for which the difference between diagrammatic bridge number and the actual bridge number of the knot grows to infinity.

- 35 Jie Chen (東北大情報) A pair of Seifert matrices that cannot have algebraic Gordian distance one 15
- Jie Chen (Tohoku Univ.) A pair of Seifert matrices that cannot have algebraic Gordian distance one

概要 Murakami introduced the Gordian distance as the least crossing-changes to transform one knot into another. Based on a matrix operation analogous to the crossing-change, he also introduced the algebraic Gordian distance between Seifert matrices. We consider the restrictions when the algebraic Gordian distance is one and improve a result of Kawauchi that if two matrices have algebraic Gordian distance one, then their Alexander polynomials have a certain relation. We give new answers to a question of Jong, showing that some Alexander polynomials cannot be realized by distance one matrices if a corresponding quadratic equation does not have an integer solution.

- 36 北野晃朗 (創価大理工) Brieskorn ホモロジー 3 球面の $SL(2; \mathbb{C})$ -既約表現の Reidemeister torsion について 10
- Teruaki Kitano (Soka Univ.) Reidemeister torsion of a Brieskorn homology 3-sphere for $SL(2; \mathbb{C})$ -irreducible representations

概要 Let $M = \Sigma(a_1, a_2, a_3)$ be a Brieskorn homology 3-sphere. Here $2 \leq a_1 < a_2 < a_3$ are pairwise coprime integers. Further we suppose that $a_1 = 2$, or a_1, a_2, a_3 are odd integers. We write $\tau_\rho(M)$ to Reidemeister torsion of M for an irreducible representation $\rho : \pi_1(M) \rightarrow SL(2; \mathbb{C})$. Now we consider the set $RT(M) = \{\tau_\rho(M)\} \subset \mathbb{R}$ of all values, which is a finite set of real numbers.

In this talk, we would like to discuss the problem that how strong $RT(M)$ is as an invariant for Brieskorn homology 3-spheres and show $RT(M)$ determines M .

- 37 坂井駿介 (広島大理) 立方複体を用いた交代絡み目外部空間の特徴付け 15
- Shunsuke Sakai (Hiroshima Univ.) A characterization of alternating link exteriors in terms of cubed complexes

概要 Recently, J. Greene and J. Howie gave intrinsic characterizations of alternating links in terms of a pair of definite spanning surfaces. These answer the Fox problem which asked what non-diagrammatic properties characterize alternating links. In this talk, we give a characterization of alternating link exteriors in terms of cubed complexes.

- 38 松田能文 (青学大理工) セパタクロー絡み目について 15
- Yoshifumi Matsuda (Aoyama Gakuin Univ.) On the sepaktakraw link

概要 We call the link obtained from a ball of sepaktakraw by replacing each annulus piece with a circle the sepaktakraw link. The sepaktakraw link is an alternating link with rich symmetry. In this talk, we introduce several properties of the sepaktakraw link.

14:20~15:20 特別講演

野坂 武史 (東工大) 3次元トポロジーの冪零的研究
Takefumi Nosaka (Tokyo Tech) Nilpotent studies in 3-dimensional topology

概要 The purpose of this talk is to explain nilpotent studies in low-dimensional topology, and to introduce my recent results on this topic. In particular, we focus on the topic of Milnor–Orr invariants, higher Massey products, and tree part of the Kontsevich invariant of links. The main result is that I gave diagrammatic computation of these invariants (of appropriate degree), and computed some examples. In the nilpotent work, the nilpotent quotient of the free group plays key role. So, in this talk, I start by reviewing properties and homology of the quotient group. After that, I briefly explain the above invariants with properties, and introduce the diagrammatic computations. Here we consider a comparison with known results concerning the mapping class group. Finally, I roughly show a future plan.

15:35~17:45

39 伊藤 昇 (東大数理) On equivalence classes of spherical curves by deformations of types RI
橋爪 恵 (明大研究・知財) and RIII 15
Noboru Ito (Univ. of Tokyo) On equivalence classes of spherical curves by deformations of types RI
Megumi Hashizume (Meiji Univ.) and RIII

概要 Let P, P' be spherical curves. Suppose that P, P' are reduced spherical curves. Then the following conditions are pairwise equivalent.

(A) P' is obtained from P by applying a sequence of deformations of RI, RIII and ambient isotopy.

(B) P' is obtained from P by applying a sequence deformations of RIII, α, β and ambient isotopy.

40 伊藤 昇 (東大数理) Finite type invariants and n -similarity of virtual knots by forbidden
櫻井みぎ和 (茨城工高専) moves 10
Noboru Ito (Univ. of Tokyo) Finite type invariants and n -similarity of virtual knots by forbidden
Migiwa Sakurai moves
(Ibaraki Nat. Coll. of Tech.)

概要 Vassiliev introduced filtered invariants of knots using crossing changes (1990), called finite type invariants. For the finite type invariants, Ohyaama introduced a notion of n -triviality (1990) and Taniyama generalized it to obtain a notion of n -similarity (1992). Goussarov, Polyak, and Viro introduced universal finite type invariants of virtual knots using virtualization (2000). We mimicked their ideas, and defined finite type invariants of virtual knots and introduced a notion that corresponds to n -similarity, using forbidden moves (J. Math. Soc. Japan). In this talk, we give infinitely many examples of n -similar pairs of virtual knots by forbidden moves and show that every invariant of Goussarov, Polyak, and Viro is an invariant of us.

41 小鳥居祐香 (理化学研) C_n -move on long virtual knot and Goussarov–Polyak–Viro’s finite type
invariant 10
Yuka Kotorii (RIKEN) C_n -move on long virtual knot and Goussarov–Polyak–Viro’s finite type
invariant

概要 A C_n -move is a family of local moves on knots and links, which gives a topological characterization of finite type invariants of knots. We extend the C_n -move to (long) virtual knots by using the lower central series of the pure virtual braid, and call it a virtual C_n -move. We then prove that for long virtual knots a virtual C_n -equivalence generated by virtual C_n -moves is equal to n -equivalence, which is an equivalence relation on (long) virtual knots defined by Goussarov–Polyak–Viro. Moreover we directly prove that two long virtual knots are not distinguished by any finite type invariants of degree $n - 1$ if they are virtual C_n -equivalent, for any positive integer n .

- 42 水澤 篤彦 (Atsuhiko Mizusawa) 3成分ハンドル体絡み目の HL-homotopy 類について 15
On HL-homotopy classes for 3-component handlebody-links

概要 Two handlebody-links are HL-homotopic if they are transformed from one to the other by a sequence of self-crossing changes of their handles. For 3-component handlebody-links, we construct a bijection between the set of the HL-homotopy classes and the set of the equivalence classes of the set of tuples of one 3-dimensional matrix and three matrices with respect to some relations.

- 43 木村 直記 (早大理工) (Naoki Kimura (Waseda Univ.)) Dijkgraaf–Witten invariants of cusped hyperbolic 3-manifolds 10
Dijkgraaf–Witten invariants of cusped hyperbolic 3-manifolds

概要 The Dijkgraaf–Witten invariant is a topological invariant for compact oriented 3-manifolds in terms of a finite group and its 3-cocycle. The invariant is a state sum invariant constructed by using a triangulation, likewise the Turaev–Viro invariant. In this talk, we discuss an extension of the Dijkgraaf–Witten invariants to cusped 3-manifolds. We show that the Dijkgraaf–Witten invariants distinguish some pairs of orientable cusped hyperbolic 3-manifolds with the same hyperbolic volumes and Turaev–Viro invariants. We also give an example of a pair of cusped hyperbolic 3-manifolds with the same hyperbolic volumes and homology groups, meanwhile with the distinct Dijkgraaf–Witten invariants.

- 44 湯浅 亘 (東工大) (Wataru Yuasa (Tokyo Tech)) A q -series identity via the \mathfrak{sl}_3 colored Jones polynomials for the $(2, 2m)$ -torus link 10
A q -series identity via the \mathfrak{sl}_3 colored Jones polynomials for the $(2, 2m)$ -torus link

概要 The colored Jones polynomial is a q -polynomial invariant of links colored by irreducible representations of a simple Lie algebra. A q -series called the tail of a knot K is obtained as the limit of the \mathfrak{sl}_2 colored Jones polynomials $\{J_n(K; q)\}_n$ ($n \rightarrow \infty$). We give two explicit formulae of the tail of the \mathfrak{sl}_3 colored Jones polynomials colored by $(n, 0)$ for the $(2, 2m)$ -torus link. These two expressions of the tail derive Andrews–Gordon identities for the \mathfrak{sl}_3 false theta function.

- 45 湯浅 亘 (東工大) (Wataru Yuasa (Tokyo Tech)) The \mathfrak{sl}_3 colored Jones polynomials for 2-bridge links 10
The \mathfrak{sl}_3 colored Jones polynomials for 2-bridge links

概要 Kuperberg introduced web spaces for some Lie algebras which are generalizations of the Kauffman bracket skein module on a disk. We derive some formulas for A_1 and A_2 clasped web spaces by graphical calculus using skein theory. These formulae are colored version of the skein relation, a twist formula, and a bubble skein expansion formula. We calculate the \mathfrak{sl}_2 and \mathfrak{sl}_3 colored Jones polynomials of 2-bridge knots and links explicitly using the twist formula.

- 46 村尾 智 (筑波大数理物質) (Tomo Murao (Univ. of Tsukuba)) 多重共役カンドル彩色数と多重共役バイカンドル彩色数の関係 10
A relationship between MCQ coloring numbers and MCB coloring numbers

概要 We define a functor \mathcal{Q} from the category of multiple conjugation biquandles to that of multiple conjugation quandles. We show that for any multiple conjugation biquandle X , there is a one-to-one correspondence between the set of X -colorings and that of $\mathcal{Q}(X)$ -colorings diagrammatically for any handlebody-link and spatial trivalent graph.

- 47 松土 恵理 (日大総合基礎) (Eri Matsudo (Nihon Univ.)) \mathbb{Z} -彩色可能絡み目の最小彩色数 10
Minimal coloring number of \mathbb{Z} -colorable links

概要 The minimal coloring number of a \mathbb{Z} -colorable link is the minimal number of colors for non-trivial \mathbb{Z} -colorings on diagrams of the link. We determine the minimal coloring number for any \mathbb{Z} -colorable links.

- 48 阿 蘇 愛 理 (首都大東京理工) $(-2, 3, 2n + 1)$ -プレッツェル結び目のねじれアレキサンダー多項式 ... 15
Airi Aso (Tokyo Metro. Univ.) Twisted Alexander polynomials of $(-2, 3, 2n + 1)$ -pretzel knots

概要 We calculate the twisted Alexander polynomials of $(-2, 3, 2n + 1)$ -pretzel knots associated to their holonomy representations.

無限可積分系

3月20日(火) 第V会場

10:00~11:30

- 1 小嶋健太郎 (中大理工) q ホイン方程式の多項式解について 15
 佐藤 司 (中大理工)
 竹村剛一 (中大理工)
 Kentaro Kojima (Chuo Univ.) Polynomial solutions of q -Heun equation
 Tsukasa Sato (Chuo Univ.)
 Kouichi Takemura (Chuo Univ.)

概要 We study polynomial solutions of q -Heun equation. In particular we investigate the condition for the accessory parameter E of q -Heun equation which admits a non-zero polynomial solution.

- 2 Zhijie Chen (Yau Math. Sci. Center) Real-root property of the spectral polynomial of the Treibich–Verdier potential and related problems 15
 Ting-Jung Kuo (Nat. Taiwan Normal Univ.)
 Chang-Shou Lin (Nat. Taiwan Univ.)
 竹村剛一 (中大理工)
 Zhijie Chen (Yau Math. Sci. Center) Real-root property of the spectral polynomial of the Treibich–Verdier potential and related problems
 Ting-Jung Kuo (Nat. Taiwan Normal Univ.)
 Chang-Shou Lin (Nat. Taiwan Univ.)
 Kouichi Takemura (Chuo Univ.)

概要 We study the spectral polynomial of the Treibich–Verdier potential. Such spectral polynomial, which is a generalization of the classical Lamé polynomial, plays fundamental roles in both the finite-gap theory and the ODE theory of Heun’s equation. In this talk, we prove that all the roots of such spectral polynomial are real and distinct under some assumptions. The proof uses the classical concept of Sturm sequence and isomonodromic theories. We also prove an analogous result for a polynomial associated with a generalized Lamé equation. Differently, our new approach is based on the viewpoint of the monodromy data.

- 3 星野 歩 (広島工大工) 一列型 C, D 型 Macdonald 多項式の明示公式 15
 白石潤一 (東大数理)
 Ayumu Hoshino (Hiroshima Inst. of Tech.) Explicit formulas for one-column Macdonald polynomials of types C_n and D_n
 Jun’ichi Shiraishi (Univ. of Tokyo)

概要 We present explicit formulas for the Macdonald polynomials of types C_n and D_n in the one-column case.

- 4 伊藤雅彦 (琉球大理) G_2 型 Weyl 群不変な q 超幾何積分の行列式公式 15
 宮永愛子 (神戸大理)
 野海正俊 (神戸大理)
 Masahiko Ito (Univ. of Ryukyus) Determinant formulas for the q -hypergeometric integrals associated with the root system of type G_2
 Aiko Miyanaga (Kobe Univ.)
 Masatoshi Noumi (Kobe Univ.)

概要 We present some determinant formulas for the q -hypergeometric integrals associated with the root system of type G_2 , which generalize Macdonald’s constant term formula. We introduce a method of deriving the q -difference equation satisfied by the determinant and finding its special value.

- 5 伊藤公毅 (豊橋技科大) q サイクルのホモロジー 15
 Kôki Itô (Toyohashi Univ. of Tech.) Homology of q -cycles

概要 We introduce homology to be dual of the q -de Rham cohomology. Conventional q -cycles correspond to q -analogues of (noncompact) locally finite chains. We need regularize Jackson integrals over such q -cycles. Essentially, a regularization of such a q -cycle has been introduced. Nevertheless, such a regularization has not been understood as a compact chain. Thus, we introduce q -cycles including compact ones in the case of dimension 1.

- 6 朴佳南 (神戸大理) q 超幾何関数の一般化と、それを特殊解に持つモノドロミー保存変形 ... 15
 Kanam Park (Kobe Univ.) An extension of q -hypergeometric series and a monodromy preserving deformation

概要 Tsuda obtained a monodromy preserving deformation which has a special solution represented by a generalization of Gauss hypergeometric function. Our purpose is to obtain its q -analog. We define a series $\mathcal{F}_{M,N}$ as an extension of a q -hypergeometric series. In this talk, we give such a monodromy preserving deformation when $N = 1$.

14:00~15:30

- 7 大山陽介 (徳島大理工) q -超幾何関数 ${}_r\phi_{r-1}(\mathbf{0}; \mathbf{b}; q, x)$ の接続問題 15
 Yousuke Ohyama (Tokushima Univ.) A connection problem for basic hypergeometric series ${}_r\phi_{r-1}(\mathbf{0}; \mathbf{b}; q, x)$

概要 We show a connection formula of a linear q -differential equation satisfied by ${}_r\phi_{r-1}(\mathbf{0}; \mathbf{b}; q, x)$. We use a q -Laplace transformation to obtain an integral representation of solutions of the q -differential equation.

- 8 大山陽介 (徳島大理工) q -超幾何関数 ${}_3\phi_2(a_1, a_2, a_3; b_1, 0; q, x)$ の満たす差分方程式の q -Stokes 係数 15
 Yousuke Ohyama (Tokushima Univ.) q -Stokes coefficients of a difference equation satisfied by basic Hypergeometric Series ${}_3\phi_2(a_1, a_2, a_3; b_1, 0; q, x)$

概要 We study a resummation of a divergent solution of a q -difference equation satisfied by ${}_3\phi_2(a_1, a_2, a_3; b_1, 0; q, x)$. For the divergent series which is not hypergeometric type, we determine the q -Stokes coefficients.

- 9 神原北斗
 竹田悠人
 上野喜三雄 (早大理工) 15
 Hokuto Kanbara
 Yuto Takeda
 Kimio Ueno (Waseda Univ.) Expansion of solutions for KZ-theoretical monodromy preserving deformation in terms of multiple polylogarithms

概要 We introduce a system of nonlinear differential equations which is the integrable condition of deformation of the KZ equation of two variables (z, w) . We denote this system by 1DE which is equations in the variable w . We consider solutions holomorphic at the origin $w = 0$ of 1DE. In this talk, we will show that these solutions are expanded in terms of multiple polylogarithms.

- 10 上野喜三雄 (早大理工) モノドロミー保存変形への KZ 理論的アプローチと Schlesinger 方程式との関係 10
 Kimio Ueno (Waseda Univ.) KZ-theoretical approach to monodromy preserving deformation and its relation to Schlesinger equations

概要 We consider the relation between 1DE and the Schlesinger equation of one variable, 1SE. Particularly, we show that from constant solutions to 1DE, one can construct solutions to 1SE. Moreover an example related to Appell $F_1(\alpha, \beta, \beta', \gamma; z, zw)$ are discussed.

- 11 尾角正人 (阪市大理) アフィン非例外型のパスと臙装配位の全単射 15
 A. Schilling (UC Davis)
 T. Scrimshaw (Univ. of Queensland)
 Masato Okado (Osaka City Univ.) Bijection between paths and rigged configurations of nonexceptional
 Anne Schilling (UC Davis) affine types
 Travis Scrimshaw (Univ. of Queensland)

概要 We establish a bijection between rigged configurations and highest weight elements of a tensor product of Kirillov–Reshetikhin crystals for all nonexceptional types. A key idea for the proof is to embed both objects into bigger sets for simply-laced types $A_n^{(1)}$ or $D_n^{(1)}$, whose bijections have already been established. As a consequence we settle the $X = M$ conjecture in full generality for nonexceptional types.

- 12 高崎金久 (近畿大理工) 位相的頂点と Volterra 型可積分階層 15
 Kanehisa Takasaki (Kindai Univ.) Topological vertex and integrable hierarchies of Volterra type

概要 A generating function of the two-partition topological vertex $W_{\lambda\mu}(r)$ with positive integral parameter r is known to become a tau function of the 2D Toda hierarchy. We reconsider this special solution of the 2D Toda hierarchy in the Lax formalism, and show that this solution is related to an integrable hierarchy of the Volterra type. The case where $r = 1$ amounts to the ordinary Volterra hierarchy. This observation seems to show an interpretation of recent work of Dubrovin, Liu, Yang and Zhang on integrable structures of cubic Hodge integrals.

15:45~16:45 特別講演

- 木村太郎 臙ゲージ理論と臙 W 代数
 (慶大自然科学研究教育センター)
 Taro Kimura (Keio Univ.) Quiver gauge theory and quiver W-algebra

概要 Quiver W-algebra is a gauge theory construction of (q-deformed) W-algebra associated with a quiver. In this formalism, the generating current of the W-algebra is obtained through double quantization of Seiberg–Witten geometry, describing the moduli space of supersymmetric vacua, and the gauge theory partition function, known as the Nekrasov function, is explicitly given by a correlator of the screening charge. The formalism of quiver W-algebra naturally reproduces the construction of $W_{q,t}(g)$ by Frenkel–Reshetikhin, and also gives rise to several generalized situations for W-algebra and gauge theory: (1) affine quiver W-algebra (2) elliptic deformation of W-algebra (3) non-simply-laced (fractional) quiver variety.

3月21日(水) 第V会場

9:30~10:30

- 13 茂木康平 (東京海洋大海洋工)^b 楯円 Felderhof 模型と楯円 Schur 関数 15
 Kohei Motegi Elliptic Felderhof model and elliptic Schur functions
 (Tokyo Univ. of Marine Sci. and Tech.)

概要 We apply the recently developed Izergin–Korepin analysis on the wavefunctions of integrable lattice models to the elliptic Felderhof model. We prove that the wavefunctions are expressed as the product of a deformed elliptic Vandermonde determinant and elliptic Schur functions. As an application of the correspondence between the wavefunctions and the elliptic Schur functions, we derive dual Cauchy formula for the elliptic Schur functions.

- 14 山根 宏之 (富山大理工) Bruhat order of Weyl groupoids 15
 I. Angiono
 (Nat. Univ. of Córdoba)
Hiroyuki Yamane (Univ. of Toyama) Bruhat order of Weyl groupoids
Iván Angiono (Nat. Univ. of Córdoba)

概要 We introduce Bruhat order of Weyl groupoids. We use nil-Hecke algebras of Weyl groupoids.

- 15 橋本 義武 (東京都市大知識工) Screening operators and \mathfrak{sl}_2 action on the lattice vertex operator algebras of type A_1 15
松本 拓也 (名大多元数理)
土屋 昭博 (Kavli IPMU)
 Yoshitake Hashimoto Screening operators and \mathfrak{sl}_2 action on the lattice vertex operator algebras of type A_1
 (Tokyo City Univ.)
 Takuya Matsumoto (Nagoya Univ.)
 Akihiro Tsuchiya (Kavli IPMU)

概要 In this talk, we shall consider the marginal deformations of the Belavin–Polyakov–Zamolodchikov (BPZ) minimal models, which are the fundamental models of the two-dimensional conformal field theory. These deformations preserve the Virasoro symmetries and parametrized by the formal deformation parameter ϵ . In particular, by formulating the deformed theories over the pair $(\mathcal{K}, \mathcal{O})$ of the ring of formal power series $\mathcal{O} = \mathbb{C}[[\epsilon]]$ and the quotient field $\mathcal{K} = \mathbb{C}((\epsilon))$, we discuss the characteristic features of the BPZ minimal models and their extensions.

- 16 佐藤 僚 (東大数理) Modular transformation properties and the Verlinde formula 15
 Ryo Sato (Univ. of Tokyo) Modular transformation properties and the Verlinde formula

概要 The classification of simple modules over the $\mathcal{N} = 2$ vertex operator superalgebra (VOSA) of central charge $3(1 - \frac{2p'}{p})$ is obtained by D. Adamović via the Kazama–Suzuki coset construction. When $p' = 1$, the simple modules coincide with the $\mathcal{N} = 2$ unitary minimal series. On the other hand, when $p' > 1$, there are uncountably many simple modules and they are non-unitary. In this talk we give the modular transformation law of the characters of the simple non-unitary modules. As an application, we propose a conjectural Verlinde formula for the non-unitary $\mathcal{N} = 2$ VOSA. Note that this result is an analogue to the conjectural Verlinde formula for the admissible affine $\mathfrak{sl}(2)$ VOA proposed by T. Creutzig and D. Ridout.

10:45~11:45 特別講演

- 成瀬 弘 (山梨大教育) シューベルト・カルキュラスの視点からの Hall–Littlewood 函数の一般化・母函数表示と応用
 Hiroshi Naruse (Univ. of Yamanashi) Generalization of Hall–Littlewood function from the view point of Schubert calculus, generating function and application

概要 We generalize Hall–Littlewood function in the framework of generalized cohomology theory. We get a generating function expression for the generalized Hall–Littlewood functions. For the case of connective K-theory we recover determinantal or Pfaffian formula for K-theoretic Schur or Schur Q-function.