

 日本数学会

2017年度秋季総合分科会

英文サマリ集

2017年9月

於 山形大学

2017 日本数学会

秋季総合分科会プログラム

期 日 2017年9月11日(月)～9月14日(木)

会 場 山形大学小白川キャンパス
〒990-8560 山形市小白川町 1-4-12

連絡先 山形大学理学部
〒990-8560 山形市小白川町 1-4-12
E-mail yamagata17sept@mathsoc.jp

(会期中) Tel 090-1791-3483

一般社団法人 日本数学会
Tel 03-3835-3483

	第I会場 基盤教育1号館 112教室	第II会場 基盤教育1号館 121教室	第III会場 基盤教育1号館 122教室	第IV会場 基盤教育2号館 211教室	第V会場 基盤教育2号館 212教室	第VI会場 基盤教育2号館 213教室	第VII会場 基盤教育2号館 214教室	第VIII会場 基盤教育2号館 221教室	第IX会場 基盤教育2号館 222教室
11日 (月)	統計数学 9:15～12:00 14:15～15:00	幾何学 9:30～11:40 14:20～15:45	トポロジー 10:00～12:00 15:30～17:30	応用数学 10:00～12:00 14:15～15:45	無限可積分系 9:50～12:00 14:20～16:30	函数解析学 14:15～16:45	函数論 9:00～12:00 14:15～15:00	函数方程式論 9:00～12:00 14:15～16:15	代数学 9:15～11:45 15:30～16:50
	企画特別講演 13:00～14:00								
	特別講演 15:15～16:15 16:30～17:30	特別講演 16:00～17:00	特別講演 14:15～15:15	特別講演 16:00～17:00	特別講演 16:40～17:40	特別講演 17:00～18:00	特別講演 15:20～16:20 16:30～17:30	特別講演 16:30～17:30	特別講演 14:15～15:15
12日 (火)	統計数学 9:20～11:30 13:15～14:15		幾何学・ トポロジー 特別講演 10:30～11:30 13:00～14:00	応用数学 9:30～12:00	無限可積分系 9:50～12:00 特別講演 13:00～14:00	函数解析学 9:45～12:00 特別講演 13:00～14:00	函数論 9:00～12:00 特別講演 13:00～14:00	函数方程式論 9:15～12:00 特別講演 13:00～14:00	代数学 9:15～12:00 特別講演 13:00～14:00
	日本数学会賞授賞式(やまぎんホール)……………(14:45～15:20) 総合講演() 日本数学会賞秋季賞受賞者……………(15:30～16:30) 堤 誉 志 雄(京 大 理)……………(16:40～17:40) 懇 親 会(山形グランドホテル)……………(18:00～20:00)								
13日 (水)	統計数学 9:40～12:10	幾何学 10:00～11:40 14:20～15:45	トポロジー 10:00～12:00 15:30～17:30	応用数学 9:30～11:40 14:15～16:15	数学基礎論 および歴史 9:45～11:20 14:15～16:15	函数解析学 9:00～12:00 14:15～15:30	実函数論 9:30～12:00 14:15～15:45	函数方程式論 9:15～12:00 14:15～16:15	代数学 9:15～11:15 14:15～16:45
	企画特別講演 13:00～14:00								
	特別講演 14:15～15:15 15:30～16:30	特別講演 16:00～17:00	特別講演 14:15～15:15	特別講演 16:30～17:30	特別講演 16:30～17:30	特別講演 15:45～16:45	特別講演 16:00～17:00	特別講演 16:30～17:30	特別講演 17:00～18:00
14日 (木)			トポロジー 10:00～11:30	応用数学 9:30～10:30	数学基礎論 および歴史 9:30～11:20 14:15～16:00		実函数論 9:30～12:00 14:15～16:00	函数方程式論 9:00～12:00 14:15～16:15	代数学 9:15～12:00 14:15～16:15
	企画特別講演 13:00～14:00								
				特別講演 10:45～11:45	特別講演 16:15～17:15		特別講演 16:15～17:15	特別講演 16:30～17:30	特別講演 16:30～17:30

目次

総合講演	1
企画特別講演	2
数学基礎論および歴史	5
9月13日(水)	5
9月14日(木)	8
代数学	11
9月11日(月)	11
9月12日(火)	15
9月13日(水)	17
9月14日(木)	22
幾何学	27
9月11日(月)	27
9月12日(火)	30
9月13日(水)	30
函数論	34
9月11日(月)	34
9月12日(火)	37
函数方程式論	40
9月11日(月)	40
9月12日(火)	45
9月13日(水)	49
9月14日(木)	55
実函数論	62
9月13日(水)	62
9月14日(木)	65
函数解析学	70
9月11日(月)	70
9月12日(火)	72
9月13日(水)	74
統計数学	79
9月11日(月)	79
9月12日(火)	82
9月13日(水)	85
応用数学	89

9月11日(月)	89
9月12日(火)	92
9月13日(水)	95
9月14日(木)	99
トポロジー	102
9月11日(月)	102
9月12日(火)	106
9月13日(水)	106
9月14日(木)	110
無限可積分系	112
9月11日(月)	112
9月12日(火)	115

総 合 講 演

9月12日(火)

日本数学会賞秋季賞受賞者 (15:30~16:30)

Autumn Prize Winner

堤 誉 志 雄 (京 大 理) Well-posedness and smoothing effect for nonlinear dispersive equations (16:40~17:40)

Yoshio Tsutsumi (Kyoto Univ.) Well-posedness and smoothing effect for nonlinear dispersive equations

概要 We consider the well-posedness and the ill-posedness in the Sobolev space of the Cauchy problem for the third order nonlinear Schrödinger equation (3NLS) on the one dimensional torus. Especially, we focus on what role the smoothing type estimates of the cubic nonlinearity play in the well-posedness issue. First, I talk about the time local well-posedness in the negative Sobolev space and the nonuniqueness of solutions without auxiliary spaces for (3NLS). Second, I talk about the ill-posedness in the Sobolev space for (3NLS) with Raman scattering term. In the latter case, I also present the result of the Cauchy-Kowalevsky type on the local unique solvability in the analytic function space. These topics show that the nonlinear interaction often yields the smoothing effect.

企 画 特 別 講 演

9月11日(月)

第I会場

- 三 上 敏 夫 (津田塾大学芸) 確率最適輸送問題 —力学における E. Schrödinger の確率論的
問題のある一般化— (13:00~14:00)
- Toshio Mikami (Tsuda Coll.) Stochastic optimal transportation problem —a generaliza-
tion of probabilistic problem in E. Schrödinger’s mechanics—

概要 E. Schrödinger proposed the following problem in his 1932’s paper. Suppose that there exist N particles in a subset A of 3-dimensional Euclidean space and each particle moves independently, with a given transition probability, to a different subset B of 3-dimensional Euclidean space. He tried to find the maximal probability of such events, provided the number of particles in each point in A and B are fixed. Though he did not succeed in finding the maximal probability, he obtained Euler’s equation, for the variational problem above, which is called Schrödinger’s functional equation. After S. Bernstein’s talk in ICM 1932, E. Schrödinger’s problem has been developed as the study of Bernstein process (or reciprocal process) and that of Doob’s h -path process. It is also known that the problem is closely related to E. Nelson’s stochastic mechanics. In this talk, we focus on our research about Schrödinger’s problem and its generalization as stochastic optimal transportation problem, its application to Nelson’s stochastic mechanics, Monge’s problem as the zero-noise limit of stochastic optimal transportation problem and relation to mean field PDEs.

第VII会場

- 齋 藤 恭 司 (東大IPMU) 多変数特殊函数論をめざして (13:00~14:00)
- Kyoji Saito (Univ. of Tokyo) Special global functions in several complex variables

概要 In the first half of 20th century, general frameworks for the study of functions of several complex variables have been developed by Behnke, Thullen, Cartan, Oka, Grauert and others. They clarified the conditions for a domain to be holomorphically complete and showed sufficiently many existences of global holomorphic functions on such domains. The results were crystallized to the theory of Stein manifolds, and were sheaf theoretically formulated (Theorem A and B) by Cartan and Serre.

On the other hand, new interactions in the last few decades between mathematics and physics (e.g. string theory, gauge theory, quantum field theory, etc.) gave new impetus to mathematicians to understand global analytic functions. For instance, mirror symmetry between complex geometry and symplectic geometry was first observed by physicists, and was later formulated by Kontsevich in language of categories. Inspired by Douglas’ stability condition in physics, Tom Bridgeland introduced the space of stability conditions for such a triangulated category.

These spaces (of stability conditions) are complex manifolds of interest in mathematics and in physics. In examples, they are Stein manifolds, but do not seem to be classical symmetric domains. New problems appear: develop and understand global analytic functions on such spaces. The problem seems to be mirror symmetric to the construction of certain higher dimensional ”automorphic forms” on the period domains which I shall describe in the lecture. Our understanding of global holomorphic functions for such domains is still quite limited, compared with the rich theory of the one variable case, where there exist strong means like Fourier analysis and Eisenstein series.

In the lecture, I will describe some aspects of these problems from the perspective of the period map theory, with which I have been engaged.

第VIII会場

特別招待講演 (大韓数学会)

Hi Jun Choe (Yonsei Univ.) Regularity condition to incompressible Navier–Stokes equations (13:00~14:00)

Hi Jun Choe (Yonsei Univ.) Regularity condition to incompressible Navier–Stokes equations

概要 We discuss local regularity properties of a weak solution to the Cauchy problem of the incompressible Navier–Stokes equations. We present a new regularity criterion for the weak solution u satisfying the condition that it belongs to a critical weak Lebesgue space without any smallness assumption on that scale. As an application, we conclude that there are at most a finite number of blowup points at any singular time t . The condition that the weak Lebesgue space norm of the velocity field is bounded in time is encompassing type I singularity and significantly weaker than the end point case of the so-called Ladyzhenskaya–Prodi–Serrin condition proved by Escauriaza–Sergin–Šverák. In addition we like to roam around historical ground of the fascinating mathematical fluid questions to regularity.

9月13日(水)

第II会場

太田啓史 (名大多元数理) 探検 深谷圏 (13:00~14:00)

Hiroshi Ohta (Nagoya Univ.) Wander in Fukaya category

概要 Around the early 1990's, Kenji Fukaya introduced the notion of an A_∞ category, roughly speaking, whose objects are Lagrangian submanifolds in a symplectic manifold and the morphism spaces are the Floer chain complexes equipped with A_∞ structures defined by moduli spaces of holomorphic maps from a 2-dimensional disc to the symplectic manifold with Lagrangian boundary conditions. Now it is called Fukaya category. M. Kontsevich used this category to formulate his homological mirror symmetry conjecture. In this talk I will try to give a brief introduction and discuss some aspects of the Fukaya category with emphasis on mirror symmetry. Based on my joint works with K. Fukaya, Y.-G. Oh, K. Ono, and also with M. Abouzaid.

第VI会場

泉正己 (京大理) Dadarlat–Pennig による Dixmier–Douady 理論 (twisted K-theory) の一般化について (13:00~14:00)

Masaki Izumi (Kyoto Univ.) A generalization of the Dixmier–Douady theory (twisted K-theory) after Dadarlat–Pennig

概要 The classical Dixmier–Douady theory describes the structure of continuous trace C^* -algebras in terms of the third cohomology of its spectrum. In 1989, Rosenberg formulated twisted K-theory in full generality as the K-theory of a continuous trace C^* -algebra with its spectrum homeomorphic to a prescribed space and with a prescribed third cohomology class. Since then twisted K-theory has been extensively studied, partly because its relationship with string theory was revealed in the late '90s.

On the other hand, in the Elliott program of the classification of amenable C^* -algebras, the importance of a certain class of C^* -algebra with very simple structure had been recognized among the specialists long before the formal definition as strongly self-absorbing C^* -algebras was introduced in 2007. Recently, a surprising and unexpected application of them was found by Dadarlat–Pennig, who showed that the Dixmier–Douady theory can be generalized to every strongly self-absorbing C^* -algebra in that the classical Dixmier–Douady theory is for the trivial C^* -algebra, the complex numbers. Moreover, a generalized cohomology theory arises from every strongly self-absorbing C^* -algebra, whose characteristic classes have higher terms beyond the third cohomology. In this talk, I will give an account of this theory for non-specialists.

9月14日(木)

第VIII会場

柳田 英二 (東工大理) 非線形拡散系のダイナミクス —漸近安定性とバタフライ効果— (13:00~14:00)

Eiji Yanagida (Tokyo Tech) Dynamics of nonlinear diffusive systems —Asymptotic stability and the butterfly effect—

概要 It is well known that in some nonlinear diffusive systems, we can observe various interesting phenomena such as spatial pattern formation, traveling waves and complex spatio-temporal dynamics. Recently, it was revealed that some simple nonlinear diffusive systems can exhibit irregular behavior of solutions by a sort of the butterfly effect. In this talk, I will present a few examples and explain the mechanism of stabilization and destabilization.

第IX会場

中村 健太郎 (佐賀大理工) p -進局所ラングランズ対応と岩澤主予想 (13:00~14:00)

Kentaro Nakamura (Saga Univ.) p -adic local Langlands correspondence and Iwasawa main conjecture

概要 As a p -adic version of local Langlands correspondence, p -adic local Langlands correspondence for $GL_2(\mathbb{Q}_p)$, which is a correspondence between two dimensional irreducible p -adic representations of $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ and irreducible p -adic Banach representations of $GL_2(\mathbb{Q}_p)$, was recently established by Breuil, Berger–Breuil, Colmez, Kisin, Paskunas. As this correspondence also encodes information about p -adic variations of the both sides, it is expected to have many applications to some important problems in number theory concerning relationships between Galois side and automorphic (or analytic) side. For example, Kisin and Emerton independently applied it to prove Fontaine–Mazur conjecture on the modularity of two dimensional geometric odd p -adic representation of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$.

In our talk, I'd like to explain these topics and the recent developments. In particular, I'd like to explain another application of p -adic local Langlands correspondence, precisely, its application to the rank two case of global and local epsilon conjectures on the functional equation of Kato's Euler systems associated to Hecke eigencuspforms.

数 学 基 礎 論 お よ び 歴 史

9月13日(水) 第V会場

9:45~11:20

- 1 増田 茂 (京大数理研) The solubilities in the mathematical physics by poisson 15
Shigeru Masuda (Kyoto Univ.) The solubilities in the mathematical physics by poisson

概要 We discuss the solubilities in the Study of Mechanics of Poisson 1833, which Poisson issued again in about 20 years after the first publications in 1811, in which he discusses statics, dynamics, the hydrostatics and the hydrodynamics, relating topics, which compose of mechanics. Poisson introduces the methods of solving mathematically the problems in mechanics, in which we have a strong interest. We discuss, in introducing these methods, how he did handle his big scope, since the year 1811, standing on the basis composed of capillary action, mechanics and heat theory, and so on, including his last conclusions of problems reserving before.

- 2 増田 茂 (京大数理研) The hydrostatics and the hydrodynamics in a study of mechanics by poisson 15
Shigeru Masuda (Kyoto Univ.) The hydrostatics and the hydrodynamics in a study of mechanics by poisson

概要 We discuss two points on the Study of Mechanics of Poisson 1833, which Poisson issued in his last period of his life of learning, in which he discusses the hydrostatics and the hydrodynamics. Previously, he discuss in the precedings (1829). We are considering this as the origin of the equations of the Navier–Stokes owing to Stokes’ referring in 1859. The other point is his conjecture on the defect of the preceding proofs of exact differential. We aim to discuss his process of theoretical convergence in this arena.

- 3 田村 誠 『九章算術』 方程術における「算」の解釈について 15
(大阪産大全学教育機構)
Makoto Tamura (Osaka Sangyo Univ.) On the term “Suan,” the amount of calculations, in the chapter “Fangcheng” of “Nine Chapters”

概要 A technical term “Suan” is used as the amount of calculations in the annotation for the problem 18 of the chapter “Fangcheng” of the “Nine Chapters on the mathematical art.” However, the way of counting it has been misunderstood, so we will correct it in this talk.

- 4 脇 克志 (山形大理) 和算の画像データベースにおける幾何要素認識と自動タグ付け 15
土橋 拓馬 (明大先端数理)
Katsushi Waki (Yamagata Univ.) Towards an automatic tagging system for image databases of Japanese
Takuma Tsuchihashi (Meiji Univ.) mathematics (wasan)

概要 In recent years, construction of digital image databases from historical documents of Japanese mathematics (wasan) is progressing. Wasan researchers widely utilize these databases, for example, they search images from titles, authors and years. On the other hand, existing databases don’t have enough functions such as searching from shapes or geometric elements. In this study, the authors propose a system that helps us to tag on wasan images automatically. This system can recognize geometric elements such as triangles, squares and circles from images of geometric problems in wasan documents and it can analyze relationship such as tangency of circles and number of elements from these images. Finally, it can tag each wasan image with these information automatically.

- 5 齋藤 憲 (阪府大人間社会) 『原論』第 XII 巻の図版とその校訂 15
 Ken Saito (Osaka Pref. Univ.) Editing diagrams of Book XII of Euclid's *Elements*

概要 The Bologna manuscript (Archiginnasio Library) of Euclid's *Elements* (Heiberg's codex b) represents a tradition of Greek text totally different from other manuscripts. Though this tradition has turned out to be closer to the original thanks to W. R. Knorr's study in 1996, very little research has been done since then. There are quite a few errors of copyist both in text and diagram. Tentative edition of diagrams (with labels added or corrected, lines suppressed or supplied) will be shown.

11:30~12:00 歴史部門懇談会

14:15~16:15

- 6 依岡輝幸 (静岡大理) Aronszajn 木の組合せ論とその他の Ramsey 理論的な組合せ論の関係 15
 Teruyuki Yorioka (Shizuoka Univ.) A combinatorial relationship between Aronszajn trees and other Ramsey theoretic structures

概要 To approach a longstanding open problem on Todorčević's fragments of Martin's Axiom, we present the consistent assertions with the existence of a non-special Aronszajn tree. In this talk, we explain the assertions from not only Martin's Axiom but also Proper Forcing Axiom.

- 7 依岡輝幸 (静岡大理) Galvin の non-productive ccc posets と Suslin tree 15
 Teruyuki Yorioka (Shizuoka Univ.) Galvin's non-productive ccc posets and Suslin tree

概要 Galvin proved that, under the Continuum Hypothesis, there are two ccc posets P_0 and P_1 such that $P_0 \times P_1$ is not ccc. Roitman proved that Cohen forcing and random forcing adds such Galvin's example respectively. It is proved that it is consistent that (coherent) Suslin tree adds Galvin's example. In this talk, I will explain the motivation of this research and idea of the proof.

- 8 酒井拓史 (神戸大システム情報) Indescribable cardinals and reflection of indescribable sets 10
 Hiroshi Sakai (Kobe Univ.) Indescribable cardinals and reflection of indescribable sets

概要 We discuss relationships between Π_n^1 -indescribable cardinals and the reflection of Π_n^1 -indescribable sets. Among other things, we generalize the classical result of Jensen to show that in the constructible universe L , a Π_n^1 -indescribable cardinal κ is Π_{n+1}^1 -indescribable if and only if every Π_n^1 -indescribable subset of κ reflects.

- 9 菊池 誠 (神戸大システム情報) ZFC の可算モデル上の包含関係について 15
 J. D. Hamkins (CUNY)
 Makoto Kikuchi (Kobe Univ.) On the inclusion relations defined on countable models of ZFC
 Joel David Hamkins (CUNY)

概要 We show that the inclusion reduct of any countable model of ZFC is countable saturated. It follows that the structures arising as the inclusion relation of a countable model of ZFC are all isomorphic, and that they are exactly the countable saturated models of the theory of set-theoretic mereology: an unbounded atomic relatively complemented distributive lattice.

- 10 桔 梗 宏 孝 (神戸大情報) クラス \mathbf{K}_f について 15
 Hirotaka Kikyo (Kobe Univ.) On class \mathbf{K}_f

概要 Let A be a graph. Put $\delta(A) = |A| - \alpha e(A)$ where $|A|$ is the number of vertices in A and $e(A)$ the number of edges in A . Let f be an unbounded concave function on the set of all non-negative real numbers. \mathbf{K}_f is the class of all graphs A such that $B \subseteq A$ implies $\delta(B) \geq f(|B|)$. There are some conditions on f that imply the free amalgamation property of \mathbf{K}_f . We discuss relations between those conditions, and see how some constructions work out in \mathbf{K}_f under those conditions.

- 11 坪 井 明 人 (筑波大数理物質) Graphs and automorphism groups 10
 Akito Tsuboi (Univ. of Tsukuba) Graphs and automorphism groups

概要 K. Eda raised a question concerning a graph structure and its automorphism groups. We don't know the exact answer to this question. But we give an affirmative answer to it under an additional model theoretic assumption.

- 12 池田宏一郎 (法政大経営) Ehrenfeucht 理論に関する注意 15
 Koichiro Ikeda (Hosei Univ.) A remark on Ehrenfeucht theories

概要 A type $p \in S(T)$ is said to be special, if there are $a, b \models p$ such that $\text{tp}(b/a)$ is isolated and non-algebraic, and $\text{tp}(a/b)$ is non-isolated. I proved that a stable Ehrenfeucht theory had a special type. In this talk, I will introduce a result, which says that every Ehrenfeucht theory has a special type.

16:30~17:30 特別講演

- 竹 内 耕 太 (筑波大数理物質) Recent interaction between model theory and finite combinatorics
 Kota Takeuchi (Univ. of Tsukuba) Recent interaction between model theory and finite combinatorics

概要 The n -dependent property is one of model theoretic dividing lines between first order theories. Recent studies of the property add evidence that model theory has a deep connection with finite combinatorics, such as hyper graph, structural Ramsey theory and Vapnik–Chervonenkis theory. In this talk it will be explained how they interact with each other.

9月14日(木) 第V会場

9:30~11:20

- 13 大 藪 卓 ^b $ZF;: S \text{ =====} \Rightarrow \mathbb{R}$, 他 5 件 5
Takashi Oyabu $ZF;: S \text{ =====} \Rightarrow \mathbb{R}$, and other 5 talks

概要 (1) $ZF;: S \text{ =====} \Rightarrow \mathbb{R}$

We discuss the possibility of the construction of math. From the standpoint of the axiomatic set theory: introduction of the numbers and the construction of geometry and math.

(2) Poincaré conjectures

$G \text{ == } G' :: \text{homotopy equivalent} \text{ ==}\Rightarrow G \text{ == } G' : \text{group isomorphic};$; compact Lie groups:: $M = G/K$: homotopy equivalent $\text{ =====} \Rightarrow M \text{ == } G/K : \text{homeomorphic}$:

(3) Taniyama–Simura conjecture

Elliptic curves are modular and automorphic functions field $\text{ == modular functions field} :: \text{===== Taniyama Simura conjecture}$:

(4) Lie group, Group-manifold

$G \text{ == } G' : \text{homotopy equivalent} \text{ ==}\rightarrow G \text{ == } G' : \text{group isomorphic}$: $\text{ == } G \text{ == } G' : \text{not group isomorphic}$ $\text{ =====}\rightarrow G \text{ == } G' : \text{not homotopy equivalence} :/:$ This shows that Co-equivalence $\text{ ==}\rightarrow$ C1-equivalence- \rightarrow Cr-equivalence $::$ Hilbert 5th problem:

(5) Algebraic curve

Riemann surface $::$ algebraic curve: abelian differential $::$ abelian integral construction of Jacobi variety; Abel–Jacobi map::

(6) Construction

Construction from axiomatic set theory is discussed:: Numbers::geometry::math. Constructions:

- 14 齋 藤 三 郎 (群馬大*・再生核研)* History, logic, results and impacts of the division by zero $1/0=0$ 15
Saburou Saitoh History, logic, results and impacts of the division by zero $1/0=0$
(Gunma Univ.*/Inst. of Reproducing Kernels)

概要 In this talk, we would like to present about the simple history, basic logical background, main results and impacts to mathematics and human beings of the division by zero.

- 15 齋 藤 三 郎 (群馬大*・再生核研)^b Relations of 0 and ∞ 15
奥 村 博 (大和大教育)
松 浦 勉 (群馬大工)
Saburou Saitoh Relations of 0 and ∞
(Gunma Univ.*/Inst. of Reproducing Kernels)
Okumura Hiroshi (Yamato Univ.)
Tsutomu Matsuura (Gunma Univ.)

概要 In this talk, as the representation of the point at infinity on the Riemann sphere by the zero $z = 0$, we will show some delicate geometric relations between 0 and infinity which show a strong discontinuity at the point of infinity on the Riemann sphere:

- 16 関 隆 宏 (新潟大経営戦略本部) 適切論理の semilattice 意味論再考 15
Takahiro Seki (Niigata Univ.) A semilattice semantics in relevant logics

概要 Semilattice semantics has been considered in relevant logics. In this semantics, completeness for implicational fragment of familiar relevant logics **R** and **E** can be proved but cannot for full **R** and **E**. In this talk, we consider semantics in which completeness of weaker relevant logics (and their neighbours) can be proved.

- 17 横溝恭平 (日大理工) 全順序な Kripke frame のクラスが特徴付ける hypersequent calculus の断片について 15
 Kyohei Yokomizo (Nihon Univ.) Reducts of a hypersequent calculus characterized by the class of all totally ordered Kripke frames

概要 It is known that there are two hypersequent calculi, **GLCW** and **GLC**, characterized by the class of all totally ordered Kripke frames (equivalent to Dummett's **LC**). However, it is also known that **GLCW** is strictly weaker than **GLC** if part of the propositional logical symbols, \rightarrow , \wedge , \vee and \neg , are restricted. We correct a mistake of the proof of cut-elimination theorem of the hypersequent calculi obtained by restricting the logical symbols of **GLCW** claimed by Avron. Also, we generalize **GLCW** for all $n \geq 1$ by Jankov's characteristic formula.

- 18 藤田憲悦 (群馬大工) A formal system of reduction paths 15
 Kenetsu Fujita (Gunma Univ.) A formal system of reduction paths

概要 We introduce a formal system of reduction paths, based on which paths can be generated from a quiver by means of three operators. Next, we define reduction rules on paths and then show that the rules on paths are terminating and confluent, so that we can obtain normal paths. Following this, linearly ordered quivers Q' and Q'' can be generated by the path operators from a quiver Q called the Dynkin diagram of type A_n , such that the constructed Q' and Q'' provide witness on behalf of the Church–Rosser property for Q .

- 19 鹿島亮 (東工大情報理工) 二階の関数記号を持つ一階述語論理について 15
 中村知己 (東工大情報理工)
 Ryo Kashima (Tokyo Tech) On first order predicate logic with second order function symbols
 Kazuki Nakamura (Tokyo Tech)

概要 We introduce second order function symbols into first order predicate logic. Syntax and semantics are naturally defined. The equality axioms are extended, and the completeness theorem is proved.

11:30~12:00 数学基礎論および歴史分科会総会

14:15~16:00

- 20 倉橋太志 (木更津工高専) 第一不完全性定理の拡張と一般化について 15
 Taishi Kurahashi On extensions and generalizations of the first incompleteness theorem
 (Kisarazu Nat. Coll. of Tech.)

概要 We prove that for each $n \geq 0$, if the set of all theorems $\text{Th}(T)$ of a consistent theory T is Π_{n+1} -definable, then there exists a true Π_n sentence which is not provable in T . This improves Jeroslow's extension and Hájek's generalization of the first incompleteness theorem.

- 21 倉橋太志 (木更津工高専) Sacchetti の論理に対する算術的健全性と完全性 15
 Taishi Kurahashi Arithmetical soundness and completeness for Sacchetti's logics
 (Kisarazu Nat. Coll. of Tech.)

概要 We prove that for every recursively axiomatized consistent extension T of Peano Arithmetic and $n \geq 1$, there exists a Σ_2 numeration $\tau(v)$ of T such that whose provability logic is exactly Sacchetti's logic $K + \Box(\Box^n p \rightarrow p) \rightarrow \Box p$. This settles Sacchetti's problem.

- 22 黒田 覚 (群馬県立女大文) 限定算術と強制法 15
 Satoru Kuroda Bounded arithmetic and forcing
 (Gunma Pref. Women's Univ.)

概要 We will reformulate the forcing construction by Takeuti and Yasumoto by two-sort bounded arithmetic. As a result, we can construct models for various complexity classes below PTIME. We will also give an alternative proofs of some theorems in Takeuti–Yasumoto’s paper. Finally, we will discuss some open problems.

- 23 田中 一之 (東北大理) 無限ゲームとプッシュダウン ω 言語 15
 李文娟 (東北大理)
 Kazuyuki Tanaka (Tohoku Univ.) Infinite games and pushdown ω -languages
 Wenjuan Li (Tohoku Univ.)

概要 This research is part of a project to answer “how hard is it to show that an infinite game is determined?” In terms of the foundational program “Reverse Mathematics”, the strength of determinacy is measured by the complexity of a winning strategy required by the determinacy of a given game. In this talk, we will discuss infinite games whose winning sets are defined by deterministic 2-stack visibly pushdown automata (2DVPA), nondeterministic pushdown automata (NPDA) and some others with various acceptance.

- 24 高橋 勇人 (Random Data Lab.) Bayesian definition of random sequences with respect to conditional probabilities 15
 Hayato Takahashi (Random Data Lab.) Bayesian definition of random sequences with respect to conditional probabilities

概要 We review the recent progress on the definition of random set with respect to conditional probabilities and a generalization of van Lambalgen theorem (Takahashi 2006, 2008, 2009, 2011). In addition we generalize Kjos Hanssen theorem (2010) when the consistency of the posterior distributions holds. We propose a definition of random sequences with respect to conditional probabilities as the section of the Martin–Löf random set at the random parameters and argue the validity of the definition from the Bayesian statistical point of view.

- 25 只木 孝太郎 (中部大工) 典型性原理の量子情報処理への適用: BB84 量子暗号プロトコルの精密化 15
 Kohtaro Tadaki (Chubu Univ.) An application of the principle of typicality to quantum information processing: A refinement of the BB84 QKD protocol

概要 The notion of probability plays a crucial role in quantum mechanics. It appears as the Born rule. In modern mathematics which describes quantum mechanics, however, probability theory means nothing other than measure theory, and therefore any operational characterization of the notion of probability is still missing in quantum mechanics. In our former works, based on the toolkit of algorithmic randomness, we presented an alternative rule to the Born rule for specifying the property of results of measurements in an operational way. In this talk, we make an application of our framework to the BB84 quantum key distribution protocol in order to demonstrate how properly our framework works in practical problems in quantum mechanics.

16:15~17:15 特別講演

- 佐藤 洋祐 (東京理大理) 限量子記号消去アルゴリズムとその計算の現状について
 Yosuke Sato (Tokyo Univ. of Sci.) On quantifier elimination algorithm and current situation of its computation

代 数 学

9月11日(月) 第IX会場

9:15~11:45

- 1 和山裕嗣(東北大理) 荒川-金子型ゼータ関数の補間について 10
 大野泰生(東北大理)
 Hirotsugu Wayama (Tohoku Univ.) Interpolation of Arakawa-Kaneko-type multiple zeta functions
 Yasuo Ohno (Tohoku Univ.)

概要 We introduce a kind of generalization of multiple zeta function to interpolate Arakawa-Kaneko and Kaneko-Tsumura multiple zeta functions. We show that the function is closely related to polynomials called t -MZVs, which interpolates multiple zeta and zeta-star values, and that its values at non-positive integers can be written as polynomials whose coefficients are linear combinations of multi-poly-Bernoulli numbers.

- 2 和山裕嗣(東北大理) t 多重ゼータ値の関係式について 10
 Hirotsugu Wayama (Tohoku Univ.) On a family of relations among t -MZVs

概要 The families of Le-Murakami relations of multiple zeta values and Aoki-Ohno relations of multiple zeta-star values are not equivalent with each other. In this talk, we present new \mathbb{Q} -linear relations among t -MZVs which interpolate Le-Murakami and Aoki-Ohno relations of height 1. In our proof, two different expressions of the value of the interpolated function between Arakawa-Kaneko and Kaneko-Tsumura multiple zeta functions, introduced by Ohno and the speaker, are used.

- 3 梅澤瞭太(名大多元数理) 荒川-金子ゼータ関数の類似の関数とある種の多重ゼータ値の関係式について 10
 Ryota Umezawa (Nagoya Univ.) On an analog of Arakawa-Kaneko zeta function and a certain relation of related multiple zeta values

概要 In this talk, we discuss the function defined by Ito and prove some relations of multiple series, which can be regarded as a generalization of Mordell-Tornheim multiple zeta values.

- 4 遠藤健太(名大多元数理) Hurwitz ゼータ関数の区間 $(0, 1)$ における実零点 10
 鈴木雄太(名大多元数理)
 Kenta Endo (Nagoya Univ.) Real zeros of Hurwitz zeta-functions in the interval $(0, 1)$
 Yuta Suzuki (Nagoya Univ.)

概要 Let $0 < a \leq 1$, $s \in \mathbb{C}$, and $\zeta(s, a)$ be the Hurwitz zeta-function. Recently, T. Nakamura showed that $\zeta(\sigma, a)$ does not vanish for any $0 < \sigma < 1$ if and only if $1/2 \leq a \leq 1$. In this talk, we show that $\zeta(\sigma, a)$ has precisely one zero in the interval $(0, 1)$ if $0 < a < 1/2$. Moreover, we reveal the asymptotic behavior of this unique zero with respect to a .

- 5 川村悟史(東北大理) 高次 Mahler 測度とゼータ Mahler 測度の解析的性質 10
 Satoshi Kawamura (Tohoku Univ.) Analytic properties of higher Mahler measures and zeta Mahler measures

概要 Higher Mahler measures (HMM) and zeta Mahler measures (ZMM) are two kinds of generalization of classical Mahler measures, which were introduced by N. Kurokawa, M. Lalín and H. Ochiai and by H. Akatsuka, respectively. In this talk, we present a formula for limiting values of HMM and an analytic continuation of ZMM.

- 6 立谷 洋平 (弘前大理工) テータ関数値の代数的独立性について 10
 Yohei Tachiya (Hirosaki Univ.) Algebraic independence results for the values of the theta function

概要 The theta function is given by the series $\theta_3(\tau) := \sum_{n=-\infty}^{\infty} e^{i\pi n^2 \tau}$, which converges for τ in the complex upper half-plane \mathbb{H} . In this talk we give algebraic independence results for the values of $\theta_3(\tau)$. For example, the three values $\theta_3(\tau)$, $\theta_3(n\tau)$, and $D\theta_3(\tau)$ are algebraically independent over \mathbb{Q} for any $\tau \in \mathbb{H}$ such that $q = e^{i\pi\tau}$ is an algebraic number, where $n \geq 2$ is an integer and $D := (\pi i)^{-1} d/d\tau$ is a differential operator. This is a joint work with Carsten Elsner.

- 7 木村 巖 (富山大理工) Fricke 群上のモジュラー形式の零点について 10
 Iwao Kimura (Univ. of Toyama) Note of zeros of modular forms on Fricke groups

概要 We give a sufficient condition for zeros of certain modular forms on Fricke groups of level 2 or 3.

- 8 境 優一 (九大多重ゼータ研究センター) Modular forms of certain fractional weights and modular linear differential equations 10
 永友 清和 (阪大情報)
 Yuichi Sakai (Kyushu Univ.) Modular forms of certain fractional weights and modular linear differential equations
 Kiyokazu Nagatomo (Osaka Univ.)

概要 A modular linear differential equation is one of tools to see the relation between elliptic modular forms and characters of specific vertex operator algebras (VOAs). In this talk, we give the relation between modular forms of certain fractional weights and characters of minimal models (the simple Virasoro VOA). Furthermore, we also give the order of a modular linear differential equation which has such modular forms as solutions.

- 9 兒玉 浩尚 (工学院大学学習支援センター) On certain vector valued Siegel modular forms of type $(k, 2)$ over $\mathbb{Z}_{(p)}$ 10
 Hirotaka Kodama (Kogakuin Univ.) On certain vector valued Siegel modular forms of type $(k, 2)$ over $\mathbb{Z}_{(p)}$

概要 We will give the generators of the $M_*^{even}(\Gamma_2)$ -module of vector-valued Siegel modular forms of type $(k, 2)$ over $\mathbb{Z}_{(p)}$. This gives an example of the positive solution to more general problem whether the module of vector-valued modular forms of arbitrary degree is finitely generated over the ring of modular forms for $\mathbb{Z}_{(p)}$.

- 10 広中由美子 (早大教育) Quaternion hermitian forms の空間の球関数 10
 Yumiko Hironaka (Waseda Univ.) Spherical functions on the space of quaternion hermitian forms

概要 We define typical spherical functions $\omega(x; s)$ on the space X_n of quaternion hermitian forms over a p -adic field of size n , by Poisson transform of certain relative invariants on X_n . They can be regarded as generating functions of local densities of integral representations of quaternion hermitian forms. We study functional equations of $\omega(x; s)$ with respect to S_n acting on $s \in \mathbb{C}^n$, and give an explicit formula of $\omega(x; s)$ by the method that the author gave in a general context. The situation is similar to the space of sesquilinear forms, but the main term of the explicit formula is written by a series of symmetric Laurent polynomial of different type from usual Hall–Littlewood polynomials.

- 11 長谷川 武博 (滋賀大教育) Explicit formula of a supersingular polynomial for rank-2 Drinfeld modules 10
 Takehiro Hasegawa (Shiga Univ.) Explicit formula of a supersingular polynomial for rank-2 Drinfeld modules

概要 Rank-2 Drinfeld modules are a function-field analogue of elliptic curves. It is natural to investigate similarities and differences between rank-2 Drinfeld modules and elliptic curves. An explicit formula of a supersingular polynomial for elliptic curves was given by Max Deuring. We show an explicit formula of a supersingular polynomial for rank-2 Drinfeld modules.

- 12 矢 城 束 (東京電機大先端科学) 数列の隣接 2 項から得られる点列の離散トモグラフィー 10
 Tabane Yashiro (Tokyo Denki Univ.) Discrete tomography for the point sequence generated by the adjacent terms of a sequence

概要 Tomography is the field that reconstruct a three-dimensional object from its two-dimensional cuts. Let f be a function on \mathbb{Z}^n , and w be a finite subset of \mathbb{Z}^n . Discrete tomography reconstructs the function f from the data $f_{w+p} = \sum_{x \in w+p} f(x), p \in \mathbb{Z}^n$. This problem is proved by F. Hazama to be described completely by the zero locus of a certain polynomial in n variable associated with w . The purpose of this talk is to apply his result to the zero-sum arrays when the window w has the form $w = (s_0, s_1), (s_1, s_2), \dots, (s_{n-2}, s_{n-1}), (s_{n-1}, s_0)$, where $s_i \in \mathbb{Z} (0 \leq i \leq n-1)$. Furthermore we describe the way how one can find the rational zero-sum arrays for w .

- 13 飯 高 茂 (学習院大*) 劣完全数の基本性質 10
 Shigeru Iitaka (Gakushuin Univ.*) On subperfect numbers

概要 Given an integer m and an odd prime, if the following equality $\overline{P}\sigma(a) = Pa - m$ is satisfied then any natural number a is said to be subperfect number with base P , translation parameter m .

14:15~15:15 特別講演

- 金子 元 (筑波大数理物質) 整数及び小数の digit 展開における一様分布論の最近の展開
 Hajime Kaneko (Univ. of Tsukuba) Recent developments in uniform distribution theory: digital expansion of integers and real numbers

概要 In this talk, we discuss the uniformity property of the digits of special sequences: base- b expansion of smooth numbers and beta expansion of real numbers, where b is an integer greater than 1. In particular, we investigate asymptotic behavior of the number of nonzero digits as partial results of uniformity property. For a real number $x \geq 2$, we call a positive integer n x -smooth if every prime factor of n is at most x . It is believed that the digits 0, 1, 2 in the ternary expansion of 2-smooth numbers, namely, integers of the form $2^m (m = 0, 1, 2, \dots)$, are uniform. However, this conjecture is unsolved. We introduce recent results for the uniformity property of the digits of general smooth numbers. In particular, we consider a problem on the number of nonzero digits suggested by Bugeaud.

Next, we consider the uniformity property of the digits in the beta expansion of real numbers. Beta expansion of real numbers is a generalization of base- b expansion of real numbers, which plays an important role in dynamical systems. A real number having uniform digits in its beta expansion is called a normal number. It is difficult to show the normality of a given real number. In this talk, we introduce recent results on the normality of the beta expansion of algebraic numbers. In particular, we consider the number of nonzero digits as partial results of normality.

15:30~16:50

- 14 武田 渉 (京大 理) The subconvexity problem for relatively r -prime lattice points 10
Wataru Takeda (Kyoto Univ.) The subconvexity problem for relatively r -prime lattice points

概要 Let K be a number field and let \mathcal{O}_K be its ring of integers. We regard an m -tuple of ideals of \mathcal{O}_K as a lattice point in K^m . We say that a lattice point $(\mathfrak{a}_1, \mathfrak{a}_2, \dots, \mathfrak{a}_m)$ is relatively r -prime, if there exists no prime ideal \mathfrak{p} such that $\mathfrak{a}_1, \mathfrak{a}_2, \dots, \mathfrak{a}_m \subset \mathfrak{p}^r$.

We study the distribution of relatively r -prime lattice points in K^m with their components having norm less than x . We show some results for abelian extensions or extensions with small degree by using the subconvexity bounds of Dedekind zeta functions on the critical line.

- 15 渋川 元樹 (阪大 情報) Cubic Pell's equations associated with the simplest cubic fields 10
Genki Shibukawa (Osaka Univ.) Cubic Pell's equations associated with the simplest cubic fields

概要 We introduce a cubic analogue of the Pell's equations associated with the simplest cubic fields and write down all integer solutions of this system explicitly by using special values of complete symmetric polynomials.

- 16 島倉 雅光 (東京理大理) Ramification in Kummer extensions arising from algebraic tori 10
Masamitsu Shimakura Ramification in Kummer extensions arising from algebraic tori
(Tokyo Univ. of Sci.)

概要 We describe the ramification in cyclic extensions arising from the Kummer theory of the Weil restriction of the multiplicative group. This generalizes the classical theorem by Hecke describing the ramification of Kummer extensions.

- 17 小松 亨 (東京理大理工) 類数の整除性をみたく虚 2 次体の組の無限族について 10
Toru Komatsu (Tokyo Univ. of Sci.) On an infinite family of pairs of imaginary quadratic fields with divisibility of class numbers

概要 Let n and m be natural numbers greater than one. In this talk we construct an infinite family of imaginary quadratic fields $\mathbb{Q}(\sqrt{D})$ such that both $\mathbb{Q}(\sqrt{D})$ and $\mathbb{Q}(\sqrt{mD})$ have ideal classes of order n .

- 18 伊東 杏希子 (東京情報大総合情報)* 実二次体の類数の非可除性と岩澤 λ 不変量について (II) 10
Akiko Ito On the indivisibility of the class numbers and the Iwasawa λ -invariants
(Tokyo Univ. of Information Sci.) of real quadratic fields (II)

概要 In this talk, concerning Greenberg's conjecture, we will show the existence of certain infinite families of real quadratic fields whose Iwasawa λ -invariant of the cyclotomic \mathbb{Z}_p -extension is equal to 0.

- 19 小塚 和人 (都城工高専)* ある rational polytopes の整数点に関する Dirichlet 指標付き冪乗和と多重 Dedekind 和 10
Kazuhiro Kozuka On power sums attached to Dirichlet characters for the integer points
(Miyakonojo Nat. Coll. of Tech.) in certain rational polytopes and multiple Dedekind sums

概要 In this talk, for certain rational polytopes, we consider power sums attached to Dirichlet characters for the integer points. The main result is expressed by making use of generalized multiple Dedekind sums attached to Dirichlet characters.

- 20 南出 真 (山口大理)* ゼータ関数の微分に関連した約数問題について 10
 古屋 淳 (浜松医大)
 谷川 好男

Makoto Minamide (Yamaguchi Univ.) On representations for error terms related to the derivatives for some
 Jun Furuya Dirichlet series
 (Hamamatsu Univ. School of Medicine)
 Yoshio Tanigawa

概要 We study an error term in a certain divisor problem related to the derivatives of the Riemann zeta-function. In particular, we obtain an analogue of Chowla–Walum formula in an error term of an asymptotic formula for $\sum_{n \leq x} \sum_{d|n} d^a (\log n)^k (\log n/d)^l$. Moreover we get an upper bound for the error term.

9月12日(火) 第IX会場

9:15~12:00

- 21 宮崎 充弘 (京都教育大)* On the almost Gorenstein property of Hibi rings 15
 Mitsuhiro Miyazaki On the almost Gorenstein property of Hibi rings
 (Kyoto Univ. of Edu.)

概要 We state a criterion of when a Hibi ring is non-Gorenstein and almost Gorenstein in terms of the combinatorial structure of the poset which defines the Hibi ring.

- 22 松田 一徳 (阪大情報)* 順序凸多面体の対の正規性について 15
 Kazunori Matsuda (Osaka Univ.) Normality of twinned order polytopes

概要 I proved that the twinned order polytope associated with partially ordered sets P and Q is normal if P and Q have a common linear extension (joint work with Takayuki Hibi (Osaka University)). In this talk, I generalize this theorem.

- 23 土谷 昭善 (阪大情報) 与えられた δ 多項式を持つ Gorenstein 単体 15
 日比 孝之 (阪大情報)
 吉田 恒太郎 (阪大情報)

Akiyoshi Tsuchiya (Osaka Univ.) Gorenstein simplices with a given δ -polynomial
 Takayuki Hibi (Osaka Univ.)
 Koutarou Yoshida (Osaka Univ.)

概要 It is fashionable among the study on convex polytopes to classify the lattice polytopes with a given δ -polynomial. As a basic challenges toward the classification problem, we achieve the study on classifying lattice simplices with a given δ -polynomial of the form $1 + t^{k+1} + \dots + t^{(v-1)(k+1)}$, where $k \geq 0$ and $v > 0$ are integers. The lattice polytope with the above δ -polynomial is necessarily Gorenstein. A complete classification is already known, when v is prime. In this talk, we will give a complete classification when v is either p^2 or pq , where p and q are prime integers with $p \neq q$.

- 24 東平光生 (明大研究・知財) Sequentially generalized Cohen–Macaulay となる単体的複体と二部グラフについて 15

Hiroataka Higashidaira (Meiji Univ.) On sequentially generalized Cohen–Macaulay simplicial complexes and bipartite graphs

概要 Let K be a field, and let S be the polynomial ring in n variables over K . Let Δ be a simplicial complex with n vertices and H be a bipartite graph with n vertices. We denote by I_Δ the Stanley–Reisner ideal of Δ and by $I(H)$ the edge ideal of H . In 2003, N. T. Cuong and L. T. Nhan introduced the notion of sequentially generalized Cohen–Macaulay (seq. gen. CM for short) rings. In this talk, we investigate conditions of Δ when S/I_Δ is a seq. gen. CM ring. We see that S/I_Δ is a seq. gen. CM ring if all pure skeletons of Δ are generalized Cohen–Macaulay. As an application, we give examples of a bipartite graph H such that $S/I(H)$ is a seq. gen. CM ring.

- 25 神戸祐太 (埼玉大理工) 被約グレブナー基底のモジュライ空間の構成 15

Yuta Kambe (Saitama Univ.) Construction of the moduli space of reduced Gröbner bases

概要 A reduced Gröbner basis of an ideal I gives a flat deformation from I to its initial ideal. Therefore, when we give a monomial ideal J and its minimal generator \mathcal{C} , we are interested in the moduli space of reduced Gröbner bases whose set of leading monomials is \mathcal{C} . In this seminar, we show that there exists such a moduli space as a scheme or an ind-scheme, and we give its relation with the Hilbert scheme and a characterization of its singularity.

- 26 吉田健一 (日大文理) Strong Rees property for powers of the maximal ideal 15
渡辺敬一 (日大文理)

Ken-ichi Yoshida (Nihon Univ.) Strong Rees property for powers of the maximal ideal
Kei-ichi Watanabe (Nihon Univ.)

概要 We introduce the notion of the strong Rees property (SRP) for \mathfrak{m} -primary ideals of a Noetherian local ring and prove that any power of the maximal ideal \mathfrak{m} has its property if the associated graded ring G of \mathfrak{m} satisfies $\text{depth } G$ is greater than 1. As its application, we characterize two-dimensional excellent normal local domains so that \mathfrak{m} is a p_g -ideal, which is related to Takahashi–Dao’s question.

- 27 中村力 (岡山大自然) 可換ネーター環の導来圏の局所化と余局所化 15
吉野雄二 (岡山大自然)

Tsutomu Nakamura (Okayama Univ.) Localization and colocalization in derived categories of commutative
Yuji Yoshino (Okayama Univ.) Noetherian rings

概要 Let R be a commutative Noetherian ring. The notion of localization (resp. colocalization) functors in the derived category $D(R)$ is a natural generalization of left (resp. right) derived functors of completion (resp. section) functors. In this talk, we report several results about localization and colocalization functors. As an application, we can show that Grothendieck type vanishing theorem holds for colocalization functors. Moreover, by using localization functors, it is possible to give a simple proof of a classical theorem due to Raynaud and Gruson, which states that the projective dimension of a flat R -module is at most the Krull dimension of R .

- 28 松井 紘 樹 (名大多元数理) Some topological structures of the Balmer spectra of right bounded derived categories of commutative noetherian rings 15
- Hiroki Matsui (Nagoya Univ.) Some topological structures of the Balmer spectra of right bounded derived categories of commutative noetherian rings

概要 By virtue of Balmer's celebrated theorem, the classification of thick tensor ideals of a tensor triangulated category \mathcal{T} is equivalent to the topological structure of its Balmer spectrum $\mathrm{Spc}\mathcal{T}$. Motivated by this theorem, we discuss connectedness and noetherianity of the Balmer spectrum of a right bounded derived category of finitely generated modules over a commutative ring.

- 29 中島 晴 久 (桜美林大自然)* Liftings of pseudo-reflection groups on invariant subrings of Krull domains of algebraic subtori under actions of reductive groups 10
- Haruhisa Nakajima Liftings of pseudo-reflection groups on invariant subrings of Krull domains of algebraic subtori under actions of reductive groups
(J. F. Oberlin Univ.)

概要 Pseudo-reflections of linear representations of groups can be extended to the affine group actions on Krull domains over an algebraically closed field K . Let G be an affine algebraic group over K with a reductive identity component G^0 acting regularly on a Krull K -domain R . Let T be an algebraic closed subtorus of G and suppose that $\mathcal{Q}(R)^T = \mathcal{Q}(R^T)$ of quotient fields. We will show: If G is the centralizer of T in G , then the pseudo-reflections of the action of G on R^T can be lifted to those on R . This seems to be the best possible result for the lifting of pseudo-reflections on the invariant ring of R of a normal connected subgroup to those on R .

13:00~14:00 特別講演

- 村井 聡 (阪大情報) 多様体の単体分割の持つ組合せ論的・代数的対称性
- Satoshi Murai (Osaka Univ.) Algebraic and combinatorial duality of triangulated manifolds

概要 The numbers of faces of a triangulated manifold satisfy a certain symmetry, which is known as Klee's Dehn–Sommerville equations. In this talk, I will show that Klee's Dehn–Sommerville equations can be algebraically explained as the Matlis duality of certain quotients of Stanley–Reisner rings using Goto's work on Buchsbaum rings.

9月13日(水) 第IX会場

9:15~11:15

- 30 板場 綾 子 (東京理大理) 二重矢を持つ巡回クイバーによる自己移入的 special biserial 多元環の有限条件 (Fg) 10
- Ayako Itaba (Tokyo Univ. of Sci.) Finiteness condition (Fg) for a self-injective special biserial algebra obtained by a circular quiver with double arrows

概要 Let K be an algebraically closed field. For a positive integer s , we consider a self-injective special biserial algebra Λ_s obtained by a circular quiver with s vertices and $2s$ arrows. This algebra Λ_s is a Koszul self-injective special biserial algebra for $s \geq 1$, but is not a weakly symmetric algebra for $s \geq 3$. Our purpose in this talk is to show that, for $s \geq 3$, Λ_s satisfies the finiteness condition (Fg) introduced by Erdmann–Holloway–Taillefer–Snashall–Solberg.

- 31 塚本真由(阪市大理) Strongly quasi-hereditary algebras and rejective subcategories 10
 Mayu Tsukamoto (Osaka City Univ.) Strongly quasi-hereditary algebras and rejective subcategories

概要 Ringel introduced a special class of quasi-hereditary algebras called right-strongly quasi-hereditary algebras, motivated by Iyama's finiteness theorem of representation dimensions of artin algebras. In this talk, we give characterizations of these algebras in terms of heredity chains and right rejective subcategories. As an application, we prove that any artin algebra of global dimension at most two is always right-strongly quasi-hereditary.

- 32 吉脇理雄 (静岡大理・阪市大数学研) On isomorphisms of generalized multifold extensions of algebras without
 浅芝秀人(静岡大理) nonzero oriented cycles 10
 木村真弓
 中島健(静岡大理)
 Michio Yoshiwaki (Shizuoka Univ./Osaka City Univ.) On isomorphisms of generalized multifold extensions of algebras without
 Hideto Asashiba (Shizuoka Univ.) nonzero oriented cycles
 Mayumi Kimura
 Ken Nakashima (Shizuoka Univ.)

概要 Let A be an algebra over an algebraically closed field \mathbb{k} with a basic set A_0 of primitive idempotents, which we regard as a \mathbb{k} -category with the object set A_0 . We denote by \hat{A} the repetitive category of A , whose object set is given by $\{x^{[i]} = (x, i) \mid x \in A_0, i \in \mathbb{Z}\}$ with the Nakayama automorphism ν of \hat{A} sending $x^{[i]}$ to $x^{[i+1]}$. We set $A^{[0]}$ to be the full subcategory of \hat{A} consisting of objects $x^{[0]}$ with $x \in A_0$, and $1^{[0]}: A \rightarrow \hat{A}$ the embedding sending x to $x^{[0]}$ for all $x \in A_0$.

Let $n \in \mathbb{Z}$. Then we show that an algebra of the form $\hat{A}/\langle\phi\rangle$, where ϕ is an automorphism of \hat{A} such that $\phi(A^{[0]}) = A^{[n]}$ is isomorphic to an algebra of the form $\hat{A}/\langle\widehat{\phi}_0\nu_A^n\rangle$, where $\widehat{\phi}_0$ is an automorphism of \hat{A} naturally induced from $\phi_0 := (1^{[0]})^{-1}\phi\nu_A^{-n}1^{[0]}$ if $eAe = \mathbb{k}$ for all $e \in A_0$.

- 33 小川泰朗(名大多元数理) Auslander–Reiten duality and recollements 10
 Yasuaki Ogawa (Nagoya Univ.) Auslander–Reiten duality and recollements

概要 Let k be a commutative field. For a given finite dimensional k -algebra, it is known that an idempotent induces a recollement of module categories. In this talk, we generalize this construction for a Krull–Schmidt k -linear category equipped with a nice duality. As an application, we provide another proof of Auslander–Reiten duality.

- 34 板垣智洋(東京理大理) Symmetric Hochschild extension algebras and normalized 2-cocycles .. 10
 Tomohiro Itagaki (Tokyo Univ. of Sci.) Symmetric Hochschild extension algebras and normalized 2-cocycles

概要 In this talk, we give a sufficient condition related to 2-cocycles for Hochschild extension algebras of bound quiver algebras by the standard duality module to be symmetric.

- 35 鯉江秀行 (東京理大理) 自己入射中山多元環に対する Hochschild extension algebra の quiver 表示
 板垣智洋 (東京理大理) 10
 眞田克典 (東京理大理)
 Hideyuki Koie (Tokyo Univ. of Sci.) The ordinary quivers of Hochschild extension algebras for self-injective
 Tomohiro Itagaki (Tokyo Univ. of Sci.) Nakayama algebras
 Katsunori Sanada (Tokyo Univ. of Sci.)

概要 Let T be a Hochschild extension algebra of a finite dimensional algebra A over a field K by the standard duality A -bimodule $\text{Hom}_K(A, K)$. We determine the ordinary quiver of T if A is a self-injective Nakayama algebra by means of the \mathbb{N} -graded 2nd Hochschild homology group $HH_2(A)$ in the sense of Sköldbberg.

- 36 西中恒和 (兵庫県立大経済) 群多元環の原始環性について 10
 Tsunekazu Nishinaka (Univ. of Hyogo) On primitivity of group algebras

概要 In my talk, we introduce a certain condition satisfied many infinite groups and show that the group algebra KG of a group G satisfying the condition is primitive for any field K .

- 37 河田成人 群環上の直既約加群のヴァーテックスについて 10
 (名古屋市大システム自然)
 Shigeto Kawata (Nagoya City Univ.) On vertices of indecomposable modules over group rings

概要 Let \mathcal{O} be a complete discrete valuation ring of characteristic zero with residue class field $k = \mathcal{O}/\pi\mathcal{O}$ of characteristic $p > 0$. Let $\mathcal{O}G$ be the group ring of a finite group G over \mathcal{O} . Suppose that P is a p -subgroup of G and Q ($\neq \{1_G\}$) is a proper normal subgroup of P . We show that there exists an indecomposable $\mathcal{O}G$ -lattice X with vertex P such that all vertices of the direct summands of a kG -module $X/\pi X$ are contained in Q .

- 38 速水孝夫 (北海学園大工)* Hochschild cohomology ring of the integral group ring of a split metacyclic group 10
 Takao Hayami (Hokkai-Gakuen Univ.) Hochschild cohomology ring of the integral group ring of a split metacyclic group

概要 We will determine the ring structure of the Hochschild cohomology $HH^*(\mathbb{Z}G)$ of the integral group ring of a split metacyclic group G of order 8ℓ for arbitrary integer $\ell \geq 2$ by giving the precise description of the integral cohomology ring $H^*(G, \mathbb{Z})$.

- 39 田阪文規 (鶴岡工高専)^b The number of simple modules in a block with Klein four hyperfocal subgroup 10
 Fuminori Tasaka The number of simple modules in a block with Klein four hyperfocal
 (Nat. Inst. of Tech., Tsuruoka Coll.) subgroup

概要 A 2-block of a finite group having a Klein four hyperfocal subgroup has the same number of irreducible Brauer characters as the corresponding 2-block of the normalizer of the hyperfocal subgroup.

11:30~12:00 代数学分科会総会

14:15~16:45

- 40 小松尚夫 (Wuhan Univ.) Incomplete Fubini numbers associated with determinants 10
Takao Komatsu (Wuhan Univ.) Incomplete Fubini numbers associated with determinants

概要 We study some properties of incomplete (restricted and associated) Fubini numbers. In particular, they have the natural extensions of the original Fubini numbers in the sense of determinants. We also introduce modified incomplete (restricted and associated) Bernoulli and Cauchy numbers and study characteristic properties.

- 41 藤本光史 (福岡教育大教育) Loop generators and factorization problem in $mn - 1$ puzzle groups .. 10
Mitsushi Fujimoto Loop generators and factorization problem in $mn - 1$ puzzle groups
(Fukuoka Univ. of Edu.)

概要 We introduce *the loop generators* corresponding to rotary operations in the $mn - 1$ puzzle. They are very useful to solve the $mn - 1$ puzzle and suitable to explain algorithms to solve it because the number of them is just $m - 1$ and the rotary operation is easy to manipulate. We show that God's number for the 8 puzzle in the loop generators is 16, and an experimental result using a factorization algorithm for 547 test instances of the 15 puzzle is reported. The result teaches us that the length of the solution by the loop generators is not long compared with optimal solutions using single-tile moves.

- 42 武田裕康 (北大理) Confluent hypergeometric systems associated with principal nilpotent
齋藤睦 (北大理) p -tuples 10
Hiroyasu Takeda (Hokkaido Univ.) Confluent hypergeometric systems associated with principal nilpotent
Mutsumi Saito (Hokkaido Univ.) p -tuples

概要 Kimura and Takano introduced confluent hypergeometric systems associated with centralizers of regular elements of $\mathfrak{gl}(n, \mathbb{C})$. We introduce hypergeometric systems associated with principal nilpotent p -tuples and show to deform integrands of solutions of this systems to that of Aomoto-Gel'fand systems.

- 43 北山秀隆 (和歌山大教育) Rationality problem for purely monomial group actions 10
Hidetaka Kitayama (Wakayama Univ.) Rationality problem for purely monomial group actions

概要 Let K be a field, n be a natural number and G be a finite subgroup of $GL(n; \mathbb{Z})$. In this talk, we will consider the rationality problem for purely monomial group actions, which asks whether the fixed field $K(x_1, \dots, x_n)^G$ under the purely monomial action of G is rational over K .

- 44 陶山大輔 (北大理) B 型 Weyl 部分配置の自由性と符号付きグラフ 10
M. Torielli (北大理)
辻栄周平 (北大理)
Daisuke Suyama (Hokkaido Univ.) The freeness of the Weyl subarrangements of type B and signed graphs
Michele Torielli (Hokkaido Univ.)
Shuhei Tsujie (Hokkaido Univ.)

概要 A Weyl arrangement is the hyperplane arrangement defined by a root system. R. P. Stanley gave a characterization of the freeness of the Weyl subarrangements of type A in terms of simple graphs. The Weyl subarrangements of type B can be represented by signed graphs. Any characterization of the freeness of them has not been known. However, characterizations of the freeness for a few restricted classes are known. In this talk, we give a characterization of the freeness and supersolvability of the Weyl subarrangements of type B under certain assumption.

- 45 竹ヶ原裕元 (室蘭工大工) Lefschetz invariants and Young characters for representations of the
 小田文仁 (近畿大理工) Coxeter groups of type B 10
 吉田知行 (北星学園大経済)
 Yugen Takegahara Lefschetz invariants and Young characters for representations of the
 (Muroran Inst. of Tech.) Coxeter groups of type B
 Fumihito Oda (Kinki Univ.)
 Tomoyuki Yoshida
 (Hokusei Gakuen Univ.)

概要 Let B_n be the Coxeter group of type B . In 1978, L. Geissinger and D. Kinch presented the concept of Young subgroups of B_n and showed that there exists a \mathbb{Z} -basis of the character ring $R(B_n)$ of B_n consisting of Young characters, which forces $R(B_n)$ to be isomorphic to the partial Burnside ring $\Omega(B_n, \widehat{\mathcal{U}}_n)$ relative to the set $\widehat{\mathcal{U}}_n$ of Young subgroups of B_n . The linear \mathbb{C} -characters of B_n are identified with reduced Lefschetz invariants which are units of $\Omega(B_n, \widehat{\mathcal{U}}_n)$.

- 46 齋藤 憲 (東北大情報) 単純グラフから構成される符号の分類 10
 Ken Saito (Tohoku Univ.) A classification of codes constructed from simple graphs

概要 Let us denote a finite field with 4 elements by \mathbb{F}_4 . We will introduce an additive code over \mathbb{F}_4 of length n defined by an additive subgroup of \mathbb{F}_4^n . It is known that every self-dual additive code can be represented by the adjacency matrix of a simple undirected graph. Danielsen and Parker (2006) classified all self-dual additive codes over \mathbb{F}_4 for lengths up to 12 by using graphs on up to 12 vertices. In this talk, we give a classification of codes having the largest minimum weight among the constructed additive codes from some graphs.

- 47 知念宏司 (近畿大理工) Divisible formal weight enumerator の構成 10
 Koji Chinen (Kinki Univ.) Construction of divisible formal weight enumerators

概要 Formal weight enumerators were introduced by M. Ozeki in 1997. In this talk, we propose an algorithm for the search of similar polynomials and show some examples.

- 48 知念宏司 (近畿大理工) Riemann 予想を満たさない extremal な多項式の構成 10
 Koji Chinen (Kinki Univ.) Construction of an extremal polynomial not satisfying the Riemann hypothesis

概要 Zeta functions for codes were introduced by I. Duursma in 1999 and were generalized to other invariant polynomials by the present author. One of the famous problems is whether extremal weight enumerators satisfy the Riemann hypothesis. In this talk, we give an example of an extremal invariant polynomial (not being related to a code) not satisfying the Riemann hypothesis.

- 49 安部利之 (愛媛大教育) ムーンシャイン頂点作用素代数の \mathbb{Z}_p -軌道体構成について 10
 Ching Hung Lam (Academia Sinica)
 山田裕理
 (一橋大経済・Academia Sinica)
 Toshiyuki Abe (Ehime Univ.) \mathbb{Z}_p -orbifold constructions of the Moonshine vertex operator algebra
 Ching Hung Lam (Academia Sinica)
 Hiromichi Yamada
 (Hitotsubashi Univ./Academia Sinica)

概要 Let V^\natural be the Moonshine vertex operator algebra which is a holomorphic vertex operator algebra of central charge 24 whose full automorphism group is the Monster simple group. We prove that for primes $p = 3, 5, 7, 13$, V^\natural is constructed from the Leech lattice vertex operator algebra by a \mathbb{Z}_p -orbifold construction.

- 50 柴田大樹(岡山理大理) On twisted algebraic loop groups and affine Kac–Moody groups ····· 10
 Taiki Shibata (Okayama Univ. of Sci.) On twisted algebraic loop groups and affine Kac–Moody groups

概要 Untwisted/twisted affine Lie algebras are well understood and have a lot of applications not only in mathematics but also in theoretical physics. On the other hand, infinite-dimensional “Lie groups” constructed from given affine Lie algebras (à la C. Chevalley), which we shall call affine Kac–Moody groups, seems to be less understood. D. Peterson and V. Kac (1983) mentioned and Y. Chen (1996) proved that an untwisted affine Kac–Moody group can be realized as a central extension of an algebraic loop group. In this talk, we generalize the result to all twisted cases. This is a joint work with J. Morita (University of Tsukuba) and A. Pianzola (University of Alberta).

17:00~18:00 特別講演

- 小寺諒介(京大理) Quantized Coulomb branches of Jordan quiver gauge theories and cyclotomic rational Cherednik algebras
 Ryosuke Kodera (Kyoto Univ.) Quantized Coulomb branches of Jordan quiver gauge theories and cyclotomic rational Cherednik algebras

概要 Braverman–Finkelberg–Nakajima gave a mathematically rigorous definition of the Coulomb branches of $3d \mathcal{N} = 4$ supersymmetric gauge theories. They are certain Poisson affine algebraic varieties and admit natural quantizations. In this talk we consider the quantized Coulomb branches associated with quiver gauge theories of Jordan type. We prove that they are isomorphic to the spherical parts of cyclotomic rational Cherednik algebras. This is a joint work with Hiraku Nakajima.

9月14日(木) 第IX会場

9:15~12:00

- 51 岩見智宏(九工大工)* Compactification of ceratin Picard-torsors as a variant of Boot-Strap type theorem ····· 15
 Tomohiro Iwami (Kyushu Inst. of Tech.) Compactification of ceratin Picard-torsors as a variant of Boot-Strap type theorem

概要 We study a compacification of quotient spaces by groupoid by the associated equivalent etale cohomology. As a corollary, we give certain type of a rigid Pic-torsor which appears in the studies of unirationality of supersingular K3 surfaces or the related moduli spaces by Ekedahl–Hyland–Shepherd–Barron, or Liedtke.

- 52 長峰孝典(新潟大自然) On automorphisms and coordinates in polynomial rings ····· 15
 Takanori Nagamine (Niigata Univ.) On automorphisms and coordinates in polynomial rings

概要 In this talk, we will explain some properties of coordinates in polynomial rings by introducing some concepts which are weaker than coordinates. In particular, in the polynomial ring in two variables over an algebraically closed field of characteristic zero, we show some relations between polynomials and their fibers on an affine plane.

- 53 黒田 茂 (首都大東京理工) Non-finitely generated polynomial subrings and birational modifications of \mathbf{G}_a -actions 15
 Shigeru Kuroda (Tokyo Metro. Univ.) Non-finitely generated polynomial subrings and birational modifications of \mathbf{G}_a -actions

概要 Let k be a field of characteristic zero, R a finitely generated k -domain with field of fractions $K \neq R$, and $K[\mathbf{x}] = K[x_1, \dots, x_n]$ the polynomial ring in n variables over K . In this talk, we discuss non-finite generation of the R -algebras $B \cap R[\mathbf{x}]$ for K -subalgebras B of $K[\mathbf{x}]$. This problem is closely related to Hilbert's fourteenth problem and its generalization by Zariski. One of our results implies that, if $n \geq 3$, then every non-trivial \mathbf{G}_a -action on the affine n -space over R can be 'converted' into a \mathbf{G}_a -action on the affine $(n+1)$ -space over R with non-finitely generated invariant ring.

- 54 小田部秀介 (東北大理) On a purely inseparable analogue of the Abhyankar Conjecture for affine curves 15
 Shusuke Otabe (Tohoku Univ.) On a purely inseparable analogue of the Abhyankar Conjecture for affine curves

概要 Let U be an affine smooth curve defined over an algebraically closed field of positive characteristic. The Abhyankar Conjecture (proved by Raynaud and Harbater in 1994) describes the set of finite quotients of Grothendieck's étale fundamental group of U . In this talk, I will consider a purely inseparable analogue of this problem, formulated in terms of Nori's profinite fundamental group scheme. I will give a partial answer to it.

- 55 古谷貴彦 (明海大歯) Maximal rigid objects in an orbit category arising from a tube 15
 山内雅司 (明海大歯)
 Takahiko Furuuya (Meikai Univ.) Maximal rigid objects in an orbit category arising from a tube
 Masashi Yamauchi (Meikai Univ.)

概要 In this talk, we introduce the notion of stable k -rigid objects in a triangulated category for $k \geq 1$. We then describe some properties of stable k -rigid objects in a higher cluster tube. As a main result, we completely determine the structures of stable 2-rigid objects in a higher cluster tube.

- 56 内場崇之 (早大理工) Geometric ruled surface の接続層の導来圏に定まる gluing stability condition について 15
 Takayuki Uchiba (Waseda Univ.) Gluing stability conditions on derived category of coherent sheaves on geometric ruled surfaces

概要 We give an explicit description of gluing stability conditions on geometric ruled surfaces by introducing gluing perversity. Moreover, we describe a destabilizing wall of skyscraper sheaves on ruled surfaces by deformation of stability conditions glued from $\widetilde{GL}^+(2, \mathbb{R})$ -translates of the standard stability condition on the base curve.

- 57 川谷康太郎 (阪大理) 純層とクライン特異点 15
 Kotaro Kawatani (Osaka Univ.) Pure sheaves and Kleinian singularities

概要 Ishii-Uehara classifies pure sheaves on the fundamental cycle of the Kleinian singularity A_n . The classification is an analogue of Grothendieck's classification of vector bundles on projective lines. We study the classification of pure sheaves on the other Kleinian singularities.

- 58 西村 陵 (埼玉大理工) 自己同型群の位数が d^2 となる非特異 d 次射影平面曲線 15
 Ryou Nishimura (Saitama Univ.) A smooth projective plane curve of degree d whose automorphism group has order d^2

概要 A smooth projective plane curve which satisfies the inequality $|\text{Aut}(C)| > d^2$ is well known. In this talk, we consider a smooth projective plane curve which satisfies the equality $|\text{Aut}(C)| = d^2$.

- 59 綾野孝則 (阪市大数学研) 種数 3 の超楕円曲線のシグマ因子上の有理型関数とその力学系への応用
 V. Matveevich Buchstaber 15
 (Steklov Inst. of Math.)
 Takanori Ayano (Osaka City Univ.) The field of meromorphic functions on sigma divisor of genus 3 hyper-
 Victor Matveevich Buchstaber elliptic curve and its application to dynamical system
 (Steklov Inst. of Math.)

概要 The field of meromorphic functions on the sigma divisor of a hyperelliptic genus 3 curve is described in terms of the gradient of its sigma function. Solutions of corresponding families of polynomial dynamical systems in \mathbb{C}^4 with two polynomial integrals are constructed as an application. These systems were introduced in the work of V. M. Buchstaber and A. V. Mikhailov on the base of commuting vector fields on the symmetric products of algebraic curves.

14:15~16:15

- 60 榎園 誠 (阪大 理) 平面曲線束のスロープ等式 15
 Makoto Enokizono (Osaka Univ.) Slope equality for plane curve fibrations

概要 A fibered surface whose general fiber is a smooth plane curve is called a plane curve fibration. In this talk, I will show that relative invariants for plane curve fibrations can be localized at a finite number of fiber germs and a certain equality between local invariants, which is called a slope equality, holds for these fibrations. As a corollary, we can define a local signature for plane curve fibrations.

- 61 榎園 誠 (阪大 理) 2次元超曲面特異点に対するダーフィー型の不等式 10
 Makoto Enokizono (Osaka Univ.) Durfee-type inequality for hypersurface surface singularities

概要 In 1978, A. H. Durfee conjectured that for a hypersurface surface singularity, six times its geometric genus does not exceed its Milnor number, which is nowadays called Durfee's strong conjecture. In this talk, I will show that an inequality among invariants of hypersurface surface singularities holds and then Durfee's strong conjecture is true for such a singularity with non-negative topological Euler number of the exceptional set of the minimal resolution. For the proof, we use the method of invariants of plane curve fibrations.

- 62 那須弘和 (東海大理) Obstructions to deforming curves on a prime Fano threefold 15
 Hirokazu Nasu (Tokai Univ.) Obstructions to deforming curves on a prime Fano threefold

概要 We prove that for every smooth prime Fano threefold V , the Hilbert scheme of smooth connected curves on V contains a generically non-reduced irreducible component of Mumford type.

- 63 鈴木 拓 (早大理工) ファノ多様体上の高階極小有理曲線族 15
 Taku Suzuki (Waseda Univ.) Higher order minimal families of rational curves on Fano manifolds

概要 In this talk, we introduce higher order minimal families H of rational curves associated to Fano manifolds X . We show that H is also a Fano manifold if the Chern characters of X satisfy some positivity conditions. We also provide a sufficient condition for Fano manifolds to be covered by higher rational manifolds.

- 64 奥村克彦 (早大理工) The Classification of SNC log symplectic structures on blow-up of projective spaces 15
 Katsuhiko Okumura (Waseda Univ.) The Classification of SNC log symplectic structures on blow-up of projective spaces

概要 A log symplectic structure is a generically symplectic Poisson structure with a reduced degeneracy divisor. The hypothesis that a Poisson structure is generically symplectic ensures that degeneracy locus become a divisor and then it is an anti-canonical divisor. Thus the diversity of generically symplectic Poisson structures measures positivity of anti-canonical class in some sense. The past few years, classification of log symplectic structures on higher-dimensional Fano varieties attracts considerable attention in this view point. In this talk, I will explain classifications of log symplectic structures with simple normal crossing degeneracy divisor on blowing-up of projective spaces along a linear subspace and give two new examples.

- 65 松澤陽介 (東大数理) On the dynamical and arithmetic degrees of self-maps of algebraic varieties 15
 Yohosuke Matsuzawa (Univ. of Tokyo) On the dynamical and arithmetic degrees of self-maps of algebraic varieties

概要 For a dominant rational self-map of an algebraic variety, one can define an invariant, called (first) dynamical degree, which measures the asymptotic behavior of the iterates of the map. On the other hand, when the variety is defined over a number field, one can associate to an orbit an invariant using Weil height function, called arithmetic degree, which measures the arithmetic complexity of the orbit. Kawaguchi and Silverman conjectured that the first dynamical degree and the arithmetic degree of a Zariski dense orbit coincide. We show that the arithmetic degree is less than or equals to the dynamical degree. We also prove that the conjecture is true for endomorphisms on surfaces. This is partially joint work with Kaoru Sano and Takahiro Shibata.

- 66 樋口伸宏 (横浜国大環境情報) 2線分の場合における central stream の境界成分の分類 15
 Nobuhiro Higuchi Classification of boundary components of central streams in the two slopes case
 (Yokohama Nat. Univ.)

概要 We classify the boundary components of the central stream for a Newton polygon consisting of two slopes, where one slope is less than $1/2$ and the other slope is greater than $1/2$. By our method, we can enumerate boundary components much faster than the enumeration by using the criterion obtained by Moonen–Wedhorn. Moreover, this method is expected to have many applications. Central streams and boundary components are described in terms of truncated Dieudonné modules of level one (abbreviated as DM_1 's). To study specializations of DM_1 's, we introduce a combinatorial tool which plays an important role in our proof.

16:30~17:30 特別講演

松谷 茂樹 (佐世保工高専) Euler–Bernoulli の弾性曲線 (elastica) とその一般化: 楕円関数の萌芽からアーベル関数論の再構築へ

Shigeki Matsutani
(Sasebo Nat. Coll. of Tech.) Elastica of Euler–Bernoulli and its generalization: from sprout of elliptic functions to reconstruction of Abelian function theory

概要 Elastica is an ideal thin elastic rod. Jacob Bernoulli proposed the elastica problem 1691: to determine every shape of elastica in a plane mathematically. To solve the problem, Jacob, Daniel Bernoulli and Euler discovered mathematical facts related to the lemniscate integral, the elastic energy (the oldest harmonic map), variational method, elliptic integrals, the moduli of elliptic curves. Finally Euler completely solved the problem 1744.

As its generalization, I have been studying the statistical mechanics of elastica to find the shape of DNA. Mathematically it means to investigate the geometrical structure of the loop space with the energy. It turned out that the structure is determined by the modified KdV flows and hyperelliptic functions including their moduli.

However, the hyperelliptic function theory is not sufficient to describe the generalized elastica problem whereas the shape of original elastica is completely described by elliptic function theory. Thus I have been also studying a reconstruction of the Abelian function theory for decades with coauthors.

In this talk, after I will show the history of elastica (including its relation to lemniscate) and its generalization, I mention the recent progress of the reconstruction of the Abelian function theory and its application to the generalization of elastica.

幾何学

9月11日(月) 第II会場

9:30~11:40

- 1 今野 北斗 (東大数理) Positive scalar curvature and Seiberg–Witten invariants of tuples of diffeomorphisms 15
 Hokuto Konno (Univ. of Tokyo) Positive scalar curvature and Seiberg–Witten invariants of tuples of diffeomorphisms

概要 We construct several invariants of tuples of commutative diffeomorphisms on a 4-manifold using the Seiberg–Witten theory for families. They are generalizations of Ruberman’s invariants of diffeomorphisms on a 4-manifold using 1-parameter gauge theory. We also give an application to the spaces of positive scalar curvature metrics for 4-manifolds which cannot be studied using Ruberman’s original invariant.

- 2 相野 眞行 (名大多元数理)^b Riemannian invariants that characterize rotational symmetries of the standard sphere 15
 Masayuki Aino (Nagoya Univ.) Riemannian invariants that characterize rotational symmetries of the standard sphere

概要 Inspired by the Lichnerowicz–Obata theorem for the first eigenvalue of the Laplacian, we define a new family of invariants $\{\Omega_k(g)\}$ for closed Riemannian manifolds. The value of $\Omega_k(g)$ sharply reflects the spherical part of the manifold. Indeed, $\Omega_1(g)$ and $\Omega_2(g)$ characterize the standard sphere.

- 3 森本 眞弘 (阪市大理) 横断的楕円型作用素と熱作用素 15
 Masahiro Morimoto (Osaka City Univ.) The heat operator of a transversally elliptic operator

概要 Let G be a connected compact Lie group. A G -invariant differential operator on a compact G -manifold is said to be transversally elliptic if it is elliptic in the directions transversal to the G -orbits. In this talk we study the heat operator of a transversally elliptic operator. After we review the spectral properties of a transversally elliptic operator, we define and investigate the character, that is a distribution on G generalizing the trace of the heat operator to the G -equivariant case.

- 4 瀬戸 樹 (名大多元数理) 分割された多様体上の巡回コサイクルと相対指数定理 15
 Tatsuki Seto (Nagoya Univ.) A cyclic cocycle and relative index theorems on partitioned manifolds

概要 In this talk, we extend Roe’s cyclic 1-cocycle to relative settings. We also state two relative index theorems for partitioned manifolds by using its cyclic cocycle, which are generalizations of index theorems on partitioned manifolds. One of these theorems is a variant of [arXiv:1411.6090, Theorem 3.3].

- 5 赤川 晋哉 (阪大理)^b Vanishing theorems of L^2 -cohomology groups on regular convex cones with the Cheng–Yau metrics 15
 Shinya Akagawa (Osaka Univ.) Vanishing theorems of L^2 -cohomology groups on regular convex cones with the Cheng–Yau metrics

概要 An open convex cone $\Omega \subset \mathbb{R}^n$ is said to be regular if Ω does not contain any full straight lines. We obtain vanishing theorems of L^2 -cohomology groups $L^2 H_{\partial}^{p,q}(\Omega, g)$ on a regular convex cone $\Omega \subset \mathbb{R}^n$ with the Cheng–Yau metric g for $p + q > n$ or $p > q$.

- 6 新井真人 (山形大理) 複素射影空間の余接束内の余等質性 1 特殊 Lagrange 部分多様体の構成
 馬場蔵人 (東京理大理工) 15
 Masato Arai (Yamagata Univ.) Special Lagrangian submanifolds and cohomogeneity one actions on the
 Kurando Baba (Tokyo Univ. of Sci.) complex projective space

概要 We give examples of cohomogeneity one special Lagrangian submanifolds in the cotangent bundle over the complex projective space, whose Calabi–Yau structure was given by Stenzel. For each example, we describe the condition of special Lagrangian as an ordinary differential equation. Our method is based on a moment map technique and the classification of cohomogeneity one actions on the complex projective space due to Takagi.

- 7 河井公大朗 (学習院大理) Frölicher–Nijenhuis cohomology on G_2 - and Spin(7)-manifolds 15
 Hông Vân Lê (CAS)
 L. Schwachhöfer (TU Dortmund)
 Kotaro Kawai (Gakushuin Univ.) Frölicher–Nijenhuis cohomology on G_2 - and Spin(7)-manifolds
 Hông Vân Lê (CAS)
 Lorenz Schwachhöfer (TU Dortmund)

概要 We show that a parallel differential form Ψ of even degree on a Riemannian manifold allows to define a natural differential both on $\Omega^*(M)$ and $\Omega^*(M, TM)$, defined via the Frölicher–Nijenhuis bracket. For instance, on a Kähler manifold, these operators are the complex differential and the Dolbeault differential, respectively. We investigate this construction when taking the differential w.r.t. the canonical parallel 4-form on a G_2 - and Spin(7)-manifold, respectively. We calculate the cohomology groups of $\Omega^*(M)$ and give a partial description of the cohomology of $\Omega^*(M, TM)$.

- 8 川村昌也 (高知工高専) Parabolic flows on almost complex manifolds 15
 Masaya Kawamura Parabolic flows on almost complex manifolds
 (Nat. Inst. of Tech., Kochi Coll.)

概要 We define two parabolic flows on almost complex manifolds, which coincide with the pluriclosed flow and the Hermitian curvature flow respectively on complex manifolds. We study the relationship between these parabolic evolution equations on a compact almost Hermitian manifold.

14:20~15:45

- 9 山本光 (東京理大) 特殊ラグランジュ部分多様体と変形エルミート・ヤン・ミルズ接続のミ
 ラー対応 15
 Hikaru Yamamoto (Tokyo Univ. of Sci.) Mirror correspondence between special Lagrangian submanifolds and
 deformed Hermitian Yang–Mills connections

概要 Leung, Yau and Zaslow defined deformed Hermitian Yang–Mills connections and gave a formal way to convert special Lagrangian submanifolds in X to deformed Hermitian Yang–Mills connections on W . In their paper, two conditions were assumed for simplicity. One is that X and W are actually lattice quotients of tangent and cotangent bundle of some common open subset B in \mathbb{R}^m . Another is that each Lagrangian submanifold can be written as a graph of a section of $X \rightarrow B$. In this talk, a way to glue their argument on a tropical manifold will be explained and a mild condition for Lagrangians will be mentioned.

- 10 中村 聡 (東北大理) トーリックファノ多様体における, 一般化されたケーラーアインシュタイン計量と一様安定性 15
Satoshi Nakamura (Tohoku Univ.) Generalized Kähler Einstein metrics and uniform stability for toric Fano manifolds

概要 We give a complete criterion for the existence of generalized Kähler Einstein metrics on toric Fano manifolds from view points of a uniform stability in a sense of GIT and the properness of a functional on the space of Kähler metrics.

- 11 野村 亮介 (東大数理) 標準束が semipositive なコンパクト Kähler 多様体に対する宮岡–Yau の不等式 15
Ryosuke Nomura (Univ. of Tokyo) Miyaoka–Yau inequality for compact Kähler manifolds with semipositive canonical bundles

概要 The Miyaoka–Yau inequality is an inequality for Chern classes. In this talk, we prove it for compact Kähler manifold with semipositive canonical bundle.

- 12 高橋 良輔 (東北大理) Anti-canonically balanced 計量の安定性 15
斎藤 俊輔 (東北大 AIMR)
Ryosuke Takahashi (Tohoku Univ.) Stability of anti-canonically balanced metrics
Shunsuke Saito (Tohoku Univ.)

概要 Anti-canonically balanced metrics are quantization of Kähler–Einstein metrics on Fano manifolds. We introduce a new algebro-geometric stability on Fano manifolds and show that the existence of anti-canonically balanced metrics implies our stability. The relation between our stability and others is also discussed.

- 13 高橋 良輔 (東北大理) Smooth approximation of the modified conical Kähler–Ricci flow 15
Ryosuke Takahashi (Tohoku Univ.) Smooth approximation of the modified conical Kähler–Ricci flow

概要 On a Fano manifold M , the conical Kähler–Ricci flow (CKRF) evolves Kähler metrics while preserving cone singularities along an effective divisor D . When D is smooth and M admits a conical Kähler–Einstein metric, the limiting behavior of CKRF is studied by Liu–Zhang. However, allowing D to have simple normal crossing support changes the whole situation. In this talk, we consider the case when D is simple normal crossing and M admits a conical Kähler–Ricci soliton. In order to study the limiting behavior of CKRF, we construct the regularized flow of CKRF modified by a holomorphic vector field from the view point of the gradient flow interpretation with respect to the modified log/twisted Mabuchi K -energy.

16:00~17:00 特別講演

- 新田 泰文 (東工大理工) 標準ケーラー計量の存在問題と多様体の安定性について
Yasufumi Nitta (Tokyo Tech) On the existence of canonical Kähler metrics and stability

概要 We will talk about the relationship between the existence of “canonical” Kähler metrics and algebraic geometric stabilities of polarized manifolds. It is conjectured that the existence of constant scalar curvature Kähler metrics will be equivalent to the notion of K-polystability, which is known as the Yau–Tian–Donaldson conjecture. For the case of Kähler–Einstein metrics on Fano manifolds, it was solved affirmatively by Chen–Donaldson–Sun. However, for general polarizations, the above conjecture is still open. In this talk, we shall discuss on the recent development for this problem and some versions. (for extremal Kähler metrics, generalized Kähler–Einstein metrics, etc.)

9月12日(火) 第Ⅲ会場

10:10~10:25 2017年度日本数学会幾何学賞授賞式

10:30~11:30 2017年度日本数学会幾何学賞受賞特別講演 (トポロジー分科会と合同)

小林 治

Weyl のゲージ理論, Schwarz 微分, そしてある球面定理

Kobayashi Osamu

Weyl's gauge theory, the Schwarzian derivative and a sphere theorem

概要 For many years I have been interested in conformal differential geometry and projective differential geometry. In this talk I would like to explain what these geometries mean at the present day and what can be expected in the future. I am now, as of June 2017, planning the talk and it will be concerned chiefly with a new conformal invariant which is similar to Yamabe's conformal invariant in many respects. In addition a conjecture, a sphere theorem, will be presented. If time permits I touch upon projective differential geometry and discuss some complements to Weyl's setting. This talk as a whole is a derivation from H. Weyl's "Reine Infinitesimalgeometrie" and subsequent developments by K. Yano, H. Yamabe and M. Obata.

13:00~14:00 2017年度日本数学会幾何学賞受賞特別講演 (トポロジー分科会と合同)

作間 誠 (広島大理) 3次元多様体のファイバー曲面とヘガード曲面

Makoto Sakuma (Hiroshima Univ.) Fiber surfaces vs Heegaard surfaces of 3-manifolds

概要 Though fiber surfaces and Heegaard surfaces have completely different natures, we can find various analogies between them. We describe the analogies from the view points of (1) the branched fibration theorem, (2) monodromy groups, (3) McShane's identity and (4) geometric structures.

9月13日(水) 第Ⅱ会場

10:00~11:40

14 山本航平 (東北大理) Cheeger 定数とパーコレーション 15

Kohei Yamamoto (Tohoku Univ.) Cheeger constant and percolation process

概要 Percolation process is a kind of probability theory. Let $G = (V, E)$ be a connected graph, and fix a parameter $p \in [0, 1]$, then we consider each edge $e \in E$ to be open with probability p independently. We have a subgraph of G which consisted only of open edges, it has some connected components. In this talk, we discuss the relationship between the number of connected components and Cheeger constant.

15 佐藤健治 (玉川大工) 高次元球面の単体の垂心の存在と一意性 15

Kenzi Satô (Tamagawa Univ.) The existence and uniqueness of orthocenters of simplices of high-dimensional spheres

概要 We consider orthocenters of simplices of the unit sphere of the n -dimensional Euclidean space. For $n = 3$, orthocenters always exist for all simplices, but for $n \geq 4$, they do not necessarily exist. Moreover, unlike the case of the Euclidean space, it is possible that there exist infinite numbers of orthocenters. In this talk, we give characterizations of the existence and the uniqueness of orthocenters.

- 16 只野 誉 (東京理大理) Some compactness theorems via m -Bakry-Émery and m -Modified Ricci curvatures with negative m 15
 Homare Tadano (Tokyo Univ. of Sci.) Some compactness theorems via m -Bakry-Émery and m -Modified Ricci curvatures with negative m

概要 In this talk, stimulated by Ohta and Takatsu, and Wylie, we shall establish some new compactness theorems for complete Riemannian manifolds via m -Bakry-Émery and m -modified Ricci curvatures with negative m . Our compactness theorems may be considered as natural generalizations of the classical compact theorems due to Ambrose, Galloway, and Cheeger-Gromov-Taylor.

- 17 椋野 純一 (名大多元数理) 正の曲率作用素を持つ擬 Riemann 多様体の基本群について 15
 Junichi Mukuno (Nagoya Univ.) On the fundamental group of semi-Riemannian manifolds with positive curvature operator

概要 We consider semi-Riemannian submersions $\pi : (E, g) \rightarrow (B, -g_B)$ under the condition with (B, g_B) Riemannian, the fiber closed Riemannian, and the horizontal distribution integrable. Then we prove that, if the non-spacelike geodesically complete semi-Riemannian manifold E has some positivity of curvature, then the fundamental group of the fiber is finite. Moreover we construct an example of semi-Riemannian submersions with some positivity of curvature, non-integrable horizontal distribution, and the finiteness of the fundamental group of the fiber.

- 18 印南 信宏 (新潟大理) 射影的ランダース変換とカットローカス 15
 Nobuhiro Innami (Niigata Univ.) Projective Randers change and cut locus

概要 Let g be a Riemannian metric on a manifold M . We call a Finsler metric $F(x, y) = \sqrt{g(x)(y, y)} + \omega(y)$ a Randers metric where ω is a 1-form on M . Two Finsler metrics are said to be pointwise projectively related if they have the same geodesics as point sets. If the 1-form ω is closed, then F is pointwise projectively related to g . We see that the invariance of the cut loci w.r.t g and F implies the exactness of a closed 1-form ω .

- 19 陶 辰 (東北大理) グラフに関する離散曲面の収束理論 15
 Chen Tao (Tohoku Univ.) A convergence theory of discrete surface on graphs

概要 In present presentation, we attempt to identify a hidden smooth surface for a given discrete surface by providing a convergence theorem of the sequence of subdivisions of a network. We prove the sequence of the Goldberg-Coxeter subdivisions of a trivalent network realized in 3-dim Euclidean space by harmonic maps are consisting of a Cauchy sequence in the Hausdorff topology. As an application, we study the Mackay Crystal and estimate the convergence of subdivisions and their normal vectors.

14:20~15:45

- 20 三浦 幸平 (宮城大基盤教育) On the moduli of isotropic and helical minimal immersions between spheres 10
 G. Toth (Rutgers Univ., Camden)
 Kouhei Miura (Miyagi Univ.) On the moduli of isotropic and helical minimal immersions between spheres
 Gabor Toth (Rutgers Univ., Camden)

概要 We first give a simple criterion for (the lowest order) isotropy of a spherical minimal immersion in terms of orthogonality relations in the third (ordinary) derivative of the image curves. This is then applied in the main result of this talk which gives a full characterization of isotropic $SU(2)$ -equivariant spherical minimal immersions of S^3 into the unit sphere of real and complex $SU(2)$ -modules. Specific examples include the polyhedral minimal immersions of which the icosahedral minimal immersion is isotropic whereas its tetrahedral and octahedral cousins are not.

- 21 佐藤 雄一郎 (首都大東京理工) 擬ユークリッド空間内の極小線織面の分類について 10
 Yuichiro Sato (Tokyo Metro. Univ.) On the classification of ruled minimal surfaces in pseudo-Euclidean space

概要 This talk gives, in generic situations, a complete classification of ruled minimal surfaces in pseudo-Euclidean space with arbitrary index. In addition, we discuss the condition for ruled minimal surfaces to exist, and give a counter-example on the problem of Bernstein type. We see that there are very fruitful ruled minimal surfaces in four dimensional Minkowski space or four dimensional pseudo-Euclidean space with neutral metric, i.e. having index 2. In particular, it should be remarkable that some of those ruled minimal surfaces are embedded in three dimensional subspace with degenerate metric of pseudo-Euclidean space.

- 22 大栗 正弘 (中大理工)* チェビシェフ作用素が対角化可能でない3次元局所等質非退化中心アファイン超曲面 10
 Masahiro Ooguri (Chuo Univ.) Three-dimensional locally homogeneous nondegenerate centroaffine hypersurfaces with nondiagonalizable Tchebychev operator

概要 We determine three-dimensional locally homogeneous nondegenerate centroaffine hypersurfaces with nondiagonalizable Tchebychev operator.

- 23 安本 真士 (阪市大数学研) 離散時間的極小曲面と離散波動方程式 15
 Masashi Yasumoto (Osaka City Univ.) Discrete timelike minimal surfaces and discrete wave equations

概要 In the continuous case, a timelike immersion with vanishing mean curvature in 3-dimensional Minkowski space is called a timelike minimal surface. Timelike minimal surfaces are highly related to linear and nonlinear wave equations. In this talk we briefly introduce a theory of discrete timelike surfaces. In particular, by a reparametrization of discrete surfaces, we show that each coordinate function of a discrete timelike minimal surface satisfies a discrete wave equation. This result provides not only the geometric meaning of special solutions for a discrete wave equation but also a new representation formula for discrete timelike minimal surfaces.

- 24 榎本 一之 (東京理大基礎工) 曲線の混合全曲率 15
 伊藤 仁一 (熊本大教育)
 Kazuyuki Enomoto Total mixed curvature of curves in Euclidean space
 (Tokyo Univ. of Sci.)
 Jin-ichi Itoh (Kumamoto Univ.)

概要 The total mixed curvature of a curve in E^3 is defined as the integral of $\sqrt{\kappa^2 + \tau^2}$, where κ is the curvature and τ is the torsion. The total mixed curvature is the length of the spherical curve defined by the principal normal vector field. We study the infimum of the total mixed curvature in a set of curves, where the endpoints and the principal normal vectors at the endpoints are prescribed. In our previous works, similar problems have been studied for the unit tangent vector and for the binormal vector.

- 25 黒瀬 俊 (関西学院大理工) ミンコフスキ平面上および2次元ド・ジッター空間上の曲線から等積中
 松浦 望 (福岡大理) 心アフィン平面曲線へのある変換 15
 Takashi Kurose (Kwansei Gakuin Univ.) Certain transformations from curves on a Minkowski plane and on a
 Nozomu Matsuura (Fukuoka Univ.) two-dimensional de Sitter space to equicentroaffine plane curves

概要 Geometric counterparts of the Miura transformation between the Korteweg–de Vries (KdV) equation and the defocusing modified KdV equation are given, by using the fact that the KdV equation arises from certain time-evolutions of equicentroaffine curves and the defocusing modified KdV equation from those of curves on a Minkowski plane and on a two-dimensional de Sitter space.

16:00~17:00 特別講演

服部 広 大 (慶 大 理 工)^b リッチ平坦多様体の無限遠点における接錐の非一意性について

Kota Hattori (Keio Univ.) The nonuniqueness of the tangent cones at infinity of Ricci-flat manifolds

概要 For a metric space (X, d) , the Gromov–Hausdorff limit of $(X, a_n d)$ as $a_n \rightarrow 0$ is called the tangent cone at infinity of (X, d) . Although the tangent cone at infinity always exists if (X, d) is a complete Riemannian manifold with nonnegative Ricci curvature, the uniqueness does not hold in general. Colding and Minicozzi showed the uniqueness under the assumption that (X, d) is a Ricci-flat manifold satisfying some additional conditions. In this talk, I will explain some examples of noncompact complete hyper-Kähler manifolds who have several tangent cones at infinity, and determine the moduli spaces of them.

函数論

9月11日(月) 第VII会場

9:00~12:00

- 1 須川 敏 幸 (東北大情報) 超幾何函数の幾何的性質について 10
 Li-Mei Wang
 (Univ. of Int. Business and Econ.)
 Toshiyuki Sugawa (Tohoku Univ.) On geometric properties of hypergeometric functions
 Li-Mei Wang
 (Univ. of Int. Business and Econ.)

概要 In this talk, we will see spirallikeness (including starlikeness) of the shifted hypergeometric function $F(z) = {}_2F_1(a, b; c; z)$ with complex parameters a, b, c . First, we observe the asymptotic behaviour of the hypergeometric function around the point $z = 1$ to obtain necessary conditions for F to be λ -spirallike for a given λ with $-\pi/2 < \lambda < \pi/2$. We next give sufficient conditions for F to be λ -spirallike. More general results will also be given in the talk if time permits.

- 2 須川 敏 幸 (東北大情報) 双曲計量による領域の凸性の特徴づけ 10
 Toshiyuki Sugawa (Tohoku Univ.) Characterizations of convexity of a domain in terms of the hyperbolic metric

概要 Let Ω be a domain in the complex plane with hyperbolic metric $\lambda_\Omega(z)|dz|$ of Gaussian curvature -4 . Mejia and Minda proved that Ω is (Euclidean) convex if and only if $d(z, \partial\Omega)\lambda_\Omega(z) \geq 1/2$ for $z \in \Omega$, where $d(z, \partial\Omega)$ denotes the Euclidean distance from z to the boundary $\partial\Omega$. In the present talk, we give spherical and hyperbolic counterparts of this result in terms of the spherical/hyperbolic density of the hyperbolic metric $\lambda_\Omega(z)|dz|$. A key idea is to obtain a geometric characterization of such convex domains relative to the spherical/hyperbolic metric.

- 3 柴 雅 和 (広島大*) 開リーマン面の閉リーマン面への等角的埋め込み — Closings と流体力学
 山口 博 史 (滋賀大*) 的周期行列 15
 Masakazu Shiba (Hiroshima Univ.*) Conformal embeddings of an open Riemann surface into closed ones of
 Hiroshi Yamaguchi (Shiga Univ.*) the same genus — closings and hydrodynamic period matrices

概要 Let R be an open Riemann surface of finite genus $g (\geq 1)$ and χ_R be a canonical homology basis of R modulo dividing cycles. A closing of (R, χ_R) is, roughly speaking, a triplet $[S, \chi_S, \iota]$ consisting of a closed Riemann surface S of genus g , a canonical homology basis χ_S , and a conformal mapping $\iota: R \rightarrow S$ which induces the prescribed correspondence between χ_R and χ_S . Denote by C the set of closings of (R, χ_R) , and let \mathfrak{M} be the set of the period matrices (τ_{jk}) of (S, χ_S) , $[S, \chi_S, \iota] \in C$. For any $(a_1, a_2, \dots, a_g) \in \mathbb{R}^g$ with $\sqrt{a_1^2 + a_2^2 + \dots + a_g^2} \neq 0$ the set $\{\sum a_j a_k \tau_{jk} \mid (\tau_{jk}) \in \mathfrak{M}\}$ is a closed disk in \mathbb{H} . We show among other things that $\partial\mathfrak{M}$ is described by the generalized period matrices derived from holomorphic differentials with hydrodynamically specific boundary behavior.

- 4 宮地 秀 樹 (阪大理) Harmonic differentials for infinitesimal deformations of singular Euclidean structures 15
 Hideki Miyachi (Osaka Univ.) Harmonic differentials for infinitesimal deformations of singular Euclidean structures

概要 In this talk, I will describe the infinitesimal deformations of singular flat structures defined from generic holomorphic quadratic differentials under the de Rham theoretic framework.

- 5 宮地 秀樹 (阪大 理) A formula of the Levi form of Teichmüller distance 15
 Hideki Miyachi (Osaka Univ.) A formula of the Levi form of Teichmüller distance

概要 I will give a formula of the Levi form of the Teichmüller distance on the Teichmüller space.

- 6 小森 洋平 (早大 教育) On Schwarz automorphic functions 15
 久保 翔太 (早大 高等学院)
 Yohei Komori (Waseda Univ.) On Schwarz automorphic functions
 Shota Kubo
 (Waseda Univ. Senior High School)

概要 Let Γ be the (a, b, c) -hyperbolic triangle group acting on $\hat{\mathbf{D}}$, the unit disk \mathbf{D} with the set of cusps of Γ . Then the quotient space $\hat{\mathbf{D}}/\Gamma$ is isomorphic to the Riemann sphere $P^1(\mathbf{D})$ which induces a meromorphic function on \mathbf{D} . It is called the Schwarz automorphic function, and we write it down explicitly in terms of a, b and c .

- 7 松崎 克彦 (早大 教育) タイヒミュラー空間の商ベアス埋め込みの単射性について 15
 Katsuhiko Matsuzaki (Waseda Univ.) Injectivity of the quotient Bers embedding of Teichmüller space

概要 The Bers embedding of the Teichmüller space is a map into the Banach space of corresponding holomorphic quadratic differentials. This induces a complex Banach manifold structure to the Teichmüller space. If we take a subspace of the universal Teichmüller space, we can usually project down the Bers embedding to a well-define map from the quotient Teichmüller space to the quotient Banach space. We call this the quotient Bers embedding but its injectivity is not a trivial matter. In this talk, we consider several cases where the injectivity holds true.

- 8 中村 豪 (愛知工大 工) 熱方程式に関する Dirichlet 問題の多項式解 15
 鈴木 紀明 (名城大 理工)
 Gou Nakamura (Aichi Inst. of Tech.) Polynomial solution to Dirichlet problems for the heat equation
 Noriaki Suzuki (Meijo Univ.)

概要 We consider polynomial solution to Dirichlet problems for the heat equation, where polynomials are in two variables x and t with real coefficients. Our interest is to determine a polynomial $\psi(x, t)$ such that for any polynomial $f(x, t)$ there exists a heat polynomial $u(x, t)$ which is equal to $f(x, t)$ on the curve $\psi(x, t) = 0$ in the xt -plane. In our previous work we determined ψ of degree at most two and showed that there exist no such ψ of degree 3. In this talk we show that there exist no such ψ of degree greater than 3.

- 9 相川 弘明 (北大 理) Dichotomy of global capacity density 15
 Hiroaki Aikawa (Hokkaido Univ.) Dichotomy of global capacity density

概要 We show that the limit of the lower capacity density is equal either to 0 or to 1.

- 10 西尾 昌治 (阪市大 理) Weighted polyharmonic and polyparabolic Bergman spaces on the upper
 下村 勝孝 (茨城大 理) half space 15
 Masaharu Nishio (Osaka City Univ.) Weighted polyharmonic and polyparabolic Bergman spaces on the upper
 Katsunori Shimomura (Ibaraki Univ.) half space

概要 Polyharmonic functions are solutions of the iterated Laplace equation. In this talk, we discuss spaces of polyharmonic functions together with iterated parabolic operators on the upper half space of the Euclidean space. After explaining some basic properties of polyharmonic functions and parabolic operators of fractional order, we introduce weighted polyharmonic and polyparabolic Bergman spaces, and shall discuss their relations and reproducing properties.

- 11 下村勝孝(茨城大理) Caloric morphism with Bateman space mapping for radial metrics 15
Katsunori Shimomura (Ibaraki Univ.) Caloric morphism with Bateman space mapping for radial metrics

概要 Caloric morphisms are transformations preserving solutions of heat equation. Bateman mappings are conformal in semi-euclidean spaces. In this talk, we shall discuss problems whether there exist caloric morphisms with Bateman space mapping for radial semi-riemannian metrics.

14:15~15:00

- 12 片方江(一関工高専) Entire functions whose Julia sets include any finitely many copies of quadratic Julia sets 15
Koh Katagata Entire functions whose Julia sets include any finitely many copies of quadratic Julia sets
(Ichinoseki Nat. Coll. of Tech.)

概要 We show that for any finite collection of quadratic Julia sets, there exist a polynomial and a transcendental entire function whose Julia sets include copies of the given quadratic Julia sets. In order to prove the result, we construct quasiregular maps with required dynamics and employ the quasiconformal surgery to obtain the desired functions.

- 13 川平友規(東工大理工) Almost conformal copies of the Julia sets in the Mandelbrot set 15
木坂正史(京大人間環境)
Tomoki Kawahira (Tokyo Tech) Almost conformal copies of the Julia sets in the Mandelbrot set
Masashi Kisaka (Kyoto Univ.)

概要 We show that there are quasiconformal copies of the Cantor Julia sets embedded in the boundary of the Mandelbrot set, whose dilatations are arbitrarily close to one. Indeed, these embeddings are also close to complex affine maps. It implies that these copies are “superfine”.

- 14 上野康平(大同大)* Böttcher coordinates for holomorphic skew products 15
Kohei Ueno (Daido Univ.) Böttcher coordinates for holomorphic skew products

概要 For a holomorphic skew product with a superattracting fixed point, we construct a Böttcher coordinate on an invariant open set whose closure contains the fixed point.

15:20~16:20 特別講演

- R. Korhonen Delay differential Painlevé equations and difference Nevanlinna theory
(Univ. of Eastern Finland)
Risto Korhonen Delay differential Painlevé equations and difference Nevanlinna theory
(Univ. of Eastern Finland)

概要 Necessary conditions are obtained for certain types of rational delay differential equations to admit a transcendental meromorphic solution of hyper-order less than one. The equations obtained include delay Painlevé equations and equations solved by elliptic functions. Difference analogue of Nevanlinna theory is a central tool in the proofs of the main results. An overview of this theory, as well as some of its applications to difference Painlevé equations, are also presented.

16:30~17:30 特別講演

- 西尾昌治(阪市大理) 放物型方程式に対するポテンシャル論と関数空間
Masaharu Nishio (Osaka City Univ.) Potential theory and function spaces for parabolic equations

概要 We consider a parabolic operator $L^{(\alpha)} = \partial_t + (-\Delta)^\alpha$ on \mathbf{R}^{n+1} for $0 < \alpha \leq 1$ and $n \in \mathbf{N}$. When $\alpha = 1$, $L^{(1)}$ is the heat operator, and otherwise, $L^{(\alpha)}$ is a non-local operator. When $\alpha = 1/2$, the operator $L^{(\alpha)}$ is called the Poisson operator and closely related with harmonic functions on \mathbf{R}^{n+1} . In this talk, after recalling the potential theory for $L^{(\alpha)}$ to define $L^{(\alpha)}$ -harmonic functions, we discuss function spaces of $L^{(\alpha)}$ -harmonic functions, called the parabolic Bergman space and the parabolic Bloch space.

9月12日(火) 第VII会場

9:00~12:00

- 15 中根 静男 (東京工芸大) On formal normal forms of holomorphic germs at super-saddle fixed points 15
 Shizuo Nakane On formal normal forms of holomorphic germs at super-saddle fixed points
 (Tokyo Polytechnic Univ.)

概要 We consider a class of rigid holomorphic germs on \mathbf{C}^2 at super saddle fixed points. Their formal normal forms are given by Ruggiero. We investigate the convergence/divergence of their formal conjugacies to the normal forms. It turns out that, in most cases, the formal conjugacies diverge. We also show convergence result under some assumptions.

- 16 田島 慎一 (筑波大数理物質) Integral dependence relation と半擬斉次孤立特異点の b -関数 15
 加藤 満生 (琉球大教育)
 鍋島 克輔 (徳島大理工)
 Shinichi Tajima (Univ. of Tsukuba) B -functions of semi-quasihomogeneous hypersurface singularities and
 Mitsuo Kato (Univ. of Ryukyus) integral dependence relations
 Katsusuke Nabeshima
 (Tokushima Univ.)

概要 Annihilators in the ring of analytic linear partial differential operators associated with a μ -constant deformation of isolated hypersurface singularities are considered. Algorithmic methods of computing annihilators and b -functions are described for semi-quasihomogeneous singularities. Key ingredients of the proposed methods are local cohomology classes and integral dependence relations.

- 17 鍋島 克輔 (徳島大理工) 収束冪級数環での integral number の計算 —グレブナー基底 vs 代数的局
 田島 慎一 (筑波大数理物質) 所コホモロジー類— 10
 Katsusuke Nabeshima Algorithms for computing integral numbers in a ring of convergent
 (Tokushima Univ.) power series —Gröbner bases vs Local cohomology—
 Shinichi Tajima (Univ. of Tsukuba)

概要 We present new algorithms for computing integral numbers w.r.t. an ideal in a ring of convergent power series. The problems of solving the integral numbers can be regarded as the ideal membership problems in the ring of convergent power series. In this talk, we give two methods for solving the membership problems. One is utilizing Gröbner bases and the another is utilizing local cohomology classes. We also address the question of how to generalize the methods to parametric cases.

- 18 田島 慎一 (筑波大数理物質) Transformation law による Grothendieck local residue の計算 15
 鍋島 克輔 (徳島大理工)
 Shinichi Tajima (Univ. of Tsukuba) Computing Grothendieck local residues via transformation law
 Katsusuke Nabeshima
 (Tokushima Univ.)

概要 We give an algorithm for computing Grothendieck local residues via transformation law. Actually, we need syzygy, standard bases, ideal quotient, local cohomology etc, to get Grothendieck local residues. Thus, we give the relations in the talk. Furthermore, we give some computation examples.

- 19 梅野 高司 (九州産大工) 代数体から得られるトロイダル群 15
 Takashi Umeno (Kyushu Sangyo Univ.) Toroidal groups defined by algebraic number fields

概要 Let p be any prime number. The toroidal group defined by $Q(\sqrt[p]{p})$ has no non-constant meromorphic functions on it. On the other hand, the toroidal group defined by $Q(\sqrt[p]{p})$ is a quasi-Abelian variety.

- 20 小池 貴之 (京大 理)* Complex $K3$ surfaces containing Levi-flat hypersurfaces 10
Takayuki Koike (Kyoto Univ.) Complex $K3$ surfaces containing Levi-flat hypersurfaces

概要 We show the existence of a complex $K3$ surface X which is not a Kummer surface and has a one-parameter family of Levi-flat hypersurfaces in which all the leaves are dense. We construct such X by patching two open complex surfaces obtained as the complements of tubular neighborhoods of elliptic curves embedded in blow-ups of the projective planes at general nine points.

- 21 千葉 優作 (お茶の水女大基幹) The extension of holomorphic functions on a non-pluriharmonic locus 15
Yusaku Tiba (Ochanomizu Univ.) The extension of holomorphic functions on a non-pluriharmonic locus

概要 Let $n \geq 4$ and let Ω be a bounded hyperconvex domain in \mathbb{C}^n . Let φ be a negative exhaustive smooth plurisubharmonic function on Ω . We show that any holomorphic function defined on a connected open neighborhood of the support of $(i\partial\bar{\partial}\varphi)^{n-3}$ can be extended to the holomorphic function on Ω .

- 22 濱田 英隆 (九州産大理工) Bounded support points for the Carathéodory families in several complex variables 10
Hidetaka Hamada Bounded support points for the Carathéodory families in several complex variables
(Kyushu Sangyo Univ.)

概要 In this talk, we will show that the shearing process recently introduced by Bracci can be generalized to $\mathcal{N}_A(\mathbb{B}^2)$ and $\mathcal{M}_g(\mathbb{B}^2)$, where A is a diagonal matrix whose diagonal elements are λ and 1 with $\lambda \in [1, 2)$ and $g \in H(\mathbb{U})$ is a convex (univalent) function with real coefficients such that $g(0) = 1$, $\Re g(\zeta) > 0$ for all $\zeta \in \mathbb{U}$ and $\mathbb{U}(1, a_0) \subseteq g(\mathbb{U})$, where a_0 is a constant defined by g . We also give the results for $\mathcal{M}_g(\mathbb{U}^2)$.

- 23 濱田 英隆 (九州産大理工) Bounded support points for the families with parametric representation in several complex variables 10
Hidetaka Hamada Bounded support points for the families with parametric representation in several complex variables
(Kyushu Sangyo Univ.)

概要 In this talk, we will show that the shearing process recently introduced by Bracci can be generalized to $S_A^0(\mathbb{B}^2)$, $S_g^0(\mathbb{B}^2)$, where A is a diagonal matrix whose diagonal elements are λ and 1 with $\lambda \in [1, 2)$ and $g \in H(\mathbb{U})$ is a convex (univalent) function with real coefficients such that $g(0) = 1$, $\Re g(\zeta) > 0$ for $\zeta \in \mathbb{U}$ and $\mathbb{U}(1, a_0) \subseteq g(\mathbb{U})$. As a corollary, we obtain bounded support points for these families. This result is in contrast to the one dimensional case, where all support points of S are unbounded. Also, our result shows the existence of bounded support points for various subclasses of $S^*(\mathbb{B}^2)$ and that $S_A^0(\mathbb{B}^2) \neq S^0(\mathbb{B}^2)$. We also give a result for $S_g^0(\mathbb{U}^2)$ and $S_g^*(\mathbb{U}^2)$.

- 24 濱田 英隆 (九州産大理工) The reachable families and the support points in several complex variables 10
Hidetaka Hamada The reachable families and the support points in several complex variables
(Kyushu Sangyo Univ.)

概要 In this talk, we show the relation of the reachable families and the support points of $S_A^0(\mathbb{B}^2)$ (or, $S_g^0(\mathbb{B}^2)$) and apply it to show that $\tilde{\mathcal{R}}_{\log M}(\text{id}_{\mathbb{B}^2}, \mathcal{N}_A(\mathbb{B}^2)) \neq S_A^0(\mathbb{B}^2, M)$ and $\tilde{\mathcal{R}}_{\log M}(\text{id}_{\mathbb{B}^2}, \mathcal{M}_g(\mathbb{B}^2)) \neq S_g^0(\mathbb{B}^2, M)$, where A is a diagonal matrix whose diagonal elements are λ and 1 with $\lambda \in [1, 2)$ and $g \in H(\mathbb{U})$ is a convex (univalent) function with real coefficients such that $g(0) = 1$, $\Re g(\zeta) > 0$ for all $\zeta \in \mathbb{U}$ and $\mathbb{U}(1, a_0) \subseteq g(\mathbb{U})$. This result provides a basic difference between the theory of bounded univalent mappings on the unit disc \mathbb{U} and that on the unit ball \mathbb{B}^n , $n \geq 2$.

- 25 清水 悟 (東北大理)* 可解な自己同型群をもつチューブ領域 15
Satoru Shimizu (Tohoku Univ.) Tube domains with solvable groups of automorphisms

概要 Let T_Ω be a tube domain in \mathbf{C}^n with polynomial infinitesimal automorphisms and $\mathfrak{g}(T_\Omega)$ the Lie algebra of all complete holomorphic vector fields on T_Ω . By definition, every element of $\mathfrak{g}(T_\Omega)$ has the form of a polynomial vector field. The investigation into the tube domain T_Ω such that $\mathfrak{g}(T_\Omega)$ is solvable has significance to the general study of tube domains with polynomial infinitesimal automorphisms. We have made an experimental investigation into such a case previously. In this talk, we discuss the general structure of solvable $\mathfrak{g}(T_\Omega)$, which gives a development to the previous investigation.

- 26 野口潤次郎 (東大*)^b A remark on Oka's coherence without Weierstrass' preparation theorem and the Oka theory 15
Junjiro Noguchi (Univ. of Tokyo*) A remark on Oka's coherence without Weierstrass' preparation theorem and the Oka theory

概要 The proofs of Oka's Coherence Theorems are based on Weierstrass' Preparation (division) Theorem. Here we observe that a Weak Coherence of Oka proved without Weierstrass' Preparation (division) Theorem, but only with *power series expansions* is sufficient to prove Oka's Jôku-Ikô and hence Cousin I, II, holomorphic extensions, and Levi's Problem, as far as the domain spaces are non-singular. The proof of the Weak Coherence of Oka is almost of linear algebra. We will present some new or simplified arguments in the proofs.

13:00~14:00 特別講演

- 本田 竜 広 (広島工大工) Distortion theorems for holomorphic mappings on bounded symmetric domains
Tatsuhiro Honda Distortion theorems for holomorphic mappings on bounded symmetric domains
(Hiroshima Inst. of Tech.) domains

概要 In this talk, we will generalize distortion theorems for normalized holomorphic functions on the unit disc in \mathbf{C} to normalized holomorphic mappings on bounded symmetric domains in a higher dimensional complex Banach space.

函数方程式論

9月11日(月) 第VIII会場

9:00~12:00

- 1 松本圭司(北大理) Relative twisted (co)homology groups associated with Lauricella's F_D 12
 Keiji Matsumoto (Hokkaido Univ.) Relative twisted (co)homology groups associated with Lauricella's F_D

概要 We define a relative twisted homology group $H_1(T, D; \mathcal{L})$ isomorphic to the space of local solutions to Lauricella's hypergeometric system F_D for any parameters. We define a relative twisted cohomology group $H^1(T, D; \mathcal{L})$ as its dual space. We show that $H^1(T, D; \mathcal{L})$ is isomorphic to three kinds of twisted de Rham cohomology groups. We define an intersection form between relative twisted homology groups and that between relative twisted cohomology groups, and show their compatibility.

- 2 泉英明(千葉工大情報) Abel 関数方程式の漸化式を用いた解法 12
 Hideaki Izumi (Chiba Inst. of Tech.) Solving Abel equation by using recurrence relations

概要 In this talk, we deal with recurrence relations of the form

$$a_{n+1} - a_n = P(a_n),$$

where P is a polynomial consisting of $\deg \geq 2$ terms or more generally,

$$a_{n+k} - a_n = P(a_n, a_{n+1}, a_{n+2}, \dots, a_{n+k-1})$$

and find general terms which converge to 0.

- 3 筒井亨(千葉大理) 確定特異点型偏微分方程式の特性特異初期値問題の解の特異性について 12
 Toru Tsutsui (Chiba Univ.) Propagation of regular singularities in a complex analytic characteristic initial value problem

概要 We consider a characteristic initial value problem of a class of second order linear partial differential equation with regular singular initial data in the complex domain. We express the solution by means of series of hypergeometric functions, and show that the solution has regular singularities on three intersecting hypersurfaces. We also clarify the structure of analytic continuation of the solution.

- 4 小原功任(金沢大理工) 非心複素 Wishart 行列の最大固有値の分布関数の超幾何微分方程式による計算 9
 高山信毅(神戸大理) F. Danufane (金沢大自然)
 Katsuyoshi Ohara (Kanazawa Univ.) Numerical evaluation of the distribution function of the largest root of
 Takayama Nobuki (Kobe Univ.) complex non-central Wishart matrices
 Fadil Danufane (Kanazawa Univ.)

概要 Kang and Alouini gave a determinant formula for the cumulative distribution function of the largest root of complex non-central Wishart matrices in their study of a wireless communication system with multiple antennas. The entries of the determinant formula are expressed in terms of a hypergeometric function in 2 variables. We give an asymptotic formula of the hypergeometric function and give a stable numerical analysis scheme to evaluate the hypergeometric function.

- 5 岩木 耕平 (名大多元数理) (合流型) 超幾何微分方程式の Voros 係数の位相的漸化式による表示とそ
 小池 達也 (神戸大理) の応用 12
 竹井優美子 (神戸大理)
 Kohei Iwaki (Nagoya Univ.) On the expression of Voros coefficients for (confluent) hypergeometric
 Tatsuya Koike (Kobe Univ.) differential equations in terms of the topological recursion and its ap-
 Yumiko Takei (Kobe Univ.) plications

概要 Voros coefficients are important objects in exact WKB analysis to study global behaviors of solutions of differential equations. In this talk we will report that the Voros coefficients for (confluent) hypergeometric differential equations are given by the generating functions of free energies defined in terms of the Eynard–Orantin topological recursion. From these results, we can give concrete forms of the free energies for algebraic equations related to these equations.

- 6 五十嵐 光 On solutions of ultradiscrete Painlevé II equation with parity variables
 竹村剛一 (中大理工) 12
 Hikaru Igarashi On solutions of ultradiscrete Painlevé II equation with parity variables
 Kouichi Takemura (Chuo Univ.)

概要 We introduce a simultaneous ultradiscrete Painlevé II equation with parity variables, which is shown to be more suitable for studying two-parameter solutions than the single second-order ultradiscrete Painlevé II equation with parity variables. We investigate several types of two-parameter solutions and the solutions which are related with the ultradiscrete limit of determinant type solutions of q -Painlevé II.

- 7 川越大輔 (京大情報) Propagation of boundary-induced discontinuity in stationary radiative
 陳逸昆 (Taiwan Univ.) transfer 9
 Daisuke Kawagoe (Kyoto Univ.) Propagation of boundary-induced discontinuity in stationary radiative
 I-Kun Chen (Taiwan Univ.) transfer

概要 We consider the boundary value problem of the stationary transport equation, an integro-differential equation, in the slab domain of general dimensions. In this talk, we discuss the relation between discontinuity of the incoming boundary data and that of the solution to the boundary value problem. We introduce two conditions posed on the boundary data so that discontinuity of the boundary data propagates along the positive characteristic lines. We also introduce an example in two dimensional case which shows that piecewise continuity of the boundary data is not a sufficient condition for the main result.

- 8 一ノ瀬 弥 (信州大理) Dirac 方程式に対する経路積分の相対論的共変性について 12
 Wataru Ichinose (Shinshu Univ.) On the relativistic covariance of the Feynman path integral for the Dirac
 equation

概要 In recent days the Feynman path integral has been constructed mathematically for the Dirac equation. In this talk we will show that this Feynman path integral is relativistically covariant, i.e. has the property of spinor under the Lorentz transformations. First we give the representation of the fundamental solution by Fourier transformation with respect to momentum variables $p \in \mathbb{R}^4$ to the free Dirac equation, as of the Feynman propagator. Then, our proof can be completed by means of the theories of both the Dirac matrices and the Lie group.

- 9 柴山允瑠 (京大情報) ケプラー型ポテンシャル系のエネルギー固定問題における周期解の存在
 12

Mitsuru Shibayama (Kyoto Univ.) Periodic solutions for a prescribed-energy problem of Keplerian-type potential systems

概要 We study the existence of periodic solutions for a prescribed-energy problem of Hamiltonian systems whose potential function has a singularity at the origin like $-1/|q|^\alpha (q \in \mathbb{R}^N)$. It is known that there exist generalized periodic solutions which may have collisions, and the number of possible collisions has been estimated. In this talk we provide a new estimation of the number of collisions. Especially we show that the obtained solutions have no collision if $N \geq 2$ and $\alpha > 1$.

- 10 宇佐美広介 (岐阜大工)* 一般化された単振り子の方程式の解の減衰性 12
 Hiroyuki Usami (Gifu Univ.) Decay of solutions to generalized pendulum equations

概要 Decay of solutions to generalized pendulum equations are considered. Explicit sufficient conditions are given for solutions of such equations to decay at the infinity.

- 11 矢ヶ崎一幸 (京大情報) Fold-Hopf 分岐に対する標準形の非可積分性 12
 Kazuyuki Yagasaki (Kyoto Univ.) Nonintegrability of the normal form of the fold-Hopf bifurcation

概要 We consider the unfolding of the codimension-two fold-Hopf bifurcation and prove its meromorphic nonintegrability in the meaning of Bogoyavlenskij for almost all parameter values. Our proof is based on a generalized version of the Morales–Ramis–Simó theory for non-Hamiltonian systems and related variational equations up to second order are used.

- 12 上村豊 (東京海洋大海洋) 破裂 2-ソリトン 12
 Yutaka Kamimura A bursting 2-soliton
 (Tokyo Univ. of Marine Sci. and Tech.)

概要 It is shown that each 2-soliton obtained by an inverse scattering method with respect to an energy dependent Schrödinger equation has a finite life span, in the reflectionless case where the transmission coefficient has two poles on the imaginary axis.

- 13 大塚厚二 ^b 境界値問題の特異点集合形状最適化とソボレフ空間トレース定理との関係
 (広島国際学院大総合教育センター) 12
 Kohji Ohtsuka Relation of shape optimization of singular points in boundary value
 (Hiroshima Kokusai Gakuin Univ.) problems and trace theorem in Sobolev space

概要 The shape of the object is determined by materials and environment. The materials is described by the boundary value problem for partial differential equation with given functions (the environment) and the shape given by the boundary. We already constructed the theory of shape optimization of sets of singular points which determines the shape of the object and the method of numerical calculations exists. I will talk about that the strength of the singular point give the influence to the shape optimization process that the trace theorem of Sobolev spaces. On the contrary, we can observe the strength of a singular point with a shape optimization process.

- 14 橋詰雅斗 (阪市大理) Embedding on the Strauss's radial compactness lemma 9
 佐野めぐみ (阪市大理)
 Masato Hashizume (Osaka City Univ.) Embedding on the Strauss's radial compactness lemma
 Megumi Sano (Osaka City Univ.)

概要 We consider the embedding related to Strauss's compactness lemma. We study the sufficiently condition of compactness and non-compactness for the embedding from the radial Sobolev space to the Lebesgue space with variable exponent.

14:15~16:15

- 15 L. Cavallina (東北大情報) 二相ねじり剛性最適化問題の球対称解について 12
 Lorenzo Cavallina (Tohoku Univ.) On the spherically symmetric solutions to the two-phase torsional rigidity optimization problem

概要 We consider the unit ball $\Omega \subset \mathbb{R}^N$ ($N \geq 2$) filled with two materials with different conductivities. We perform shape derivatives up to the second order to find out precise information about locally optimal configurations with respect to the torsional rigidity functional. In particular we analyse the role played by the configuration obtained by putting a smaller concentric ball inside Ω . In this case the stress function admits an explicit form which is radially symmetric: this allows us to compute the sign of the second order shape derivative of the torsional rigidity functional with the aid of spherical harmonics. Depending on the ratio of the conductivities a symmetry breaking phenomenon occurs.

- 16 大枝和浩(早大GEC) Coexistence steady states of a prey-predator model with population flux
 久藤衡介(電通大情報理工) by attractive transition 12
 Kazuhiro Oeda (Waseda Univ.) Coexistence steady states of a prey-predator model with population flux
 Kousuke Kuto by attractive transition
 (Univ. of Electro-Comm.)

概要 This talk is concerned with the Dirichlet problem of a diffusive Lotka–Volterra prey-predator system with population flux by attractive transition. We study the global bifurcation structure of positive stationary solutions. Moreover, we discuss the asymptotic behavior of positive stationary solutions as the nonlinear diffusion coefficient approaches infinity.

- 17 柴田徹太郎(広島大工) Oscillatory structures of bifurcation curves for semilinear eigenvalue problems 12
 Tetsutaro Shibata (Hiroshima Univ.) Oscillatory structures of bifurcation curves for semilinear eigenvalue problems

概要 We study the global and local behavior of bifurcation curves for nonlinear eigenvalue problems which include some oscillatory nonlinear term g . We consider the case where λ is parameterized by the maximum norm $\alpha = \|u_\lambda\|_\infty$ of the solution u_λ corresponding to λ and is represented as $\lambda = \lambda(g, \alpha)$. Especially, we restrict our attention to the case where $\lambda(g, \alpha) \rightarrow \pi^2/4$ as $\alpha \rightarrow \infty$. We establish several precise asymptotic formulas for $\lambda(g, \alpha)$ as $\alpha \rightarrow \infty$ and $\alpha \rightarrow 0$ with the exact second terms to understand well the total structures of the bifurcation curves.

- 18 宮本安人(東大数理) ある半線形楕円型方程式の非球対称特異解の存在について 9
 Yasuhito Miyamoto (Univ. of Tokyo) On the existence of infinitely many nonradial solutions of a semilinear elliptic equation

概要 We construct countably infinitely many nonradial singular solutions of the problem

$$\Delta u + e^u = 0 \text{ in } \mathbb{R}^N \setminus \{0\}, \quad 4 \leq N \leq 10$$

of the form $u(r, \sigma) = -2 \log r + \log 2(N-2) + v(\sigma)$, where $v(\sigma)$ depends only on $\sigma \in \mathbb{S}^{N-1}$. To this end we construct countably infinitely many solutions of

$$\Delta_{\mathbb{S}^{N-1}} v + 2(N-2)(e^v - 1) = 0, \quad 4 \leq N \leq 10,$$

using ODE techniques.

- 19 A. Pomponio (Politecnico di Bari) Quasilinear elliptic equations of Born–Infeld type 12
 渡辺達也 (京都産大理)
 Alessio Pomponio (Politecnico di Bari) Quasilinear elliptic equations of Born–Infeld type
 Tatsuya Watanabe
 (Kyoto Sangyo Univ.)

概要 In this talk, we study, with variational technique, the existence of positive solutions for quasilinear elliptic equations of Born–Infeld type. We obtain the existence result for a large class of nonlinearities.

- 20 渡辺宏太郎 (防衛大) \mathbb{S}^n 上の薄い円環領域における Brezis–Nirenberg 問題の非球対称な分岐解
 塩路直樹 (横浜国大工) の構成について 9
 Kotaro Watanabe Bifurcation and symmetry breaking for Brezis–Nirenberg problem on
 (Nat. Defense Acad. of Japan) the thin annulus of the n -dimensional sphere
 Naoki Sioji (Yokohama Nat. Univ.)

概要 In this talk, we consider the Brezis–Nirenberg problem on the thin annulus of the standard sphere \mathbb{S}^n . By the consideration of Morse index of linearized equation, we construct bifurcation solution from radially symmetric solution. To obtain the results, uniqueness of the positive radial solution plays an important role.

- 21 高橋太 (阪市大理) Critical and subcritical fractional Trudinger–Moser type inequalities on
 \mathbb{R} 9
 Futoshi Takahashi (Osaka City Univ.) Critical and subcritical fractional Trudinger–Moser type inequalities on
 \mathbb{R}

概要 In this talk, we are concerned with the critical and subcritical Trudinger–Moser type inequalities for functions in a fractional Sobolev space $H^{1/2,2}$ on the whole real line. We prove the relation between two inequalities and discuss the attainability of the suprema.

- 22 高橋太 (阪市大理) On a weighted Trudinger–Moser type inequality on the whole space and
 Van Hoang Nguyen related maximizing problem 9
 (Inst. de Math. de Toulouse)
 Futoshi Takahashi (Osaka City Univ.) On a weighted Trudinger–Moser type inequality on the whole space and
 Van Hoang Nguyen related maximizing problem
 (Inst. de Math. de Toulouse)

概要 We establish a weighted Trudinger–Moser type inequality with the full Sobolev norm constraint on the whole Euclidean space. Main tool is the singular Trudinger–Moser inequality on the whole space recently established by Adimurthi and Yang, and a transformation of functions. We also discuss the existence and non-existence of maximizers for the associated variational problem.

- 23 梶木屋龍治 (佐賀大理工) 穴の空いた領域における p ラプラス方程式の対称解 12
 Ryuji Kajikiya (Saga Univ.) Symmetric solutions of p -Laplace equations in hollow domains

概要 In this lecture, we study the p -Laplace equation in a hollow symmetric domain. Let H and G be closed subgroups of the orthogonal group such that $H \subset G$ and $H \neq G$. Then we prove the existence of a positive solution which is H invariant and G non-invariant.

- 24 坂口 茂 (東北大情報)* 不変等温面を伴う二相導体と関連する楕円型優決定問題 12
 Shigeru Sakaguchi (Tohoku Univ.) Two-phase heat conductors with a stationary isothermic surface and their related elliptic overdetermined problems

概要 In the previous paper in 2016, we considered a two-phase heat conductor in \mathbb{R}^N with $N \geq 2$ consisting of a core and a shell with different constant conductivities. Among other things, when the medium outside the two-phase conductor has a possibly different conductivity, we treated the Cauchy problem for $N \geq 3$ with the initial condition where the conductor has temperature 0 and the outside medium has temperature 1. It was shown that if there is a stationary isothermic surface in the shell near the boundary, then the structure of the conductor must be spherical. Here we report that the same proposition holds true even when $N = 2$, and as by-products, we can give other proofs of all the previous results of that paper in $N(\geq 2)$ dimensions and prove a symmetry theorem on their related two-phase elliptic overdetermined problems.

16:30~17:30 特別講演

- O. Lisovyi (Univ. de Tours) Painlevé functions, Fredholm determinants and combinatorics
 Oleg Lisovyi (Univ. de Tours) Painlevé functions, Fredholm determinants and combinatorics

9月12日(火) 第VIII会場

9:15~12:00

- 25 川上翔矢 (埼玉大理工) 結び目エネルギーの離散化 12
 Shoya Kawakami (Saitama Univ.) A discretization of the knot energies

概要 “Knot energy” was proposed to answer the question of what is a “good” figure in a given knot type. A requirement of the knot energy is that the better the figure of a knot is, the lesser the value of its energy is. Jun O’Hara gave a definition of a family of knot energies satisfying such a property in 1991. One of his energies is invariant under Möbius transformations and is called the Möbius energy. To study the Möbius energy, several discrete versions have been proposed, for example, by Kim–Kusner in 1993 and by Simon in 1994. Rawdon–Simon in 2006, Rawdon–Worthington in 2010, and Scholtes in 2014 showed their convergence. In this talk, discrete energies not only of the Möbius energy but of the general energies are given, and their convergence is discussed.

- 26 郡司克徳 (埼玉大理工) 分解された Möbius エネルギーに対するLojasiewicz 不等式について ... 12
 Katsunori Gunji (Saitama Univ.) The Lojasiewicz inequalities for decomposed Möbius energies

概要 The Möbius energy is one of the knot energies defined by O’Hara. Blatt showed the global existence and convergence of the gradient flow of the Möbius energy near stationary points. The Lojasiewicz inequality played an important role for proving such results. On the other hand, by work of Ishizeki–Nagasawa, it is known that the Möbius energy can be decomposed into parts which keep the Möbius invariance. In this study, the Lojasiewicz inequality is proved for each decomposed part of the Möbius energy. In an appropriate function space setting, we can show that the 2nd variations of the decomposed Möbius energies have L^2 -representations and, using a result of Chill, the energies satisfy the Lojasiewicz inequality.

- 27 S. Blatt (Paris Lodron Univ. Salzburg) 分解された Möbius エネルギーの Möbius 不変なエネルギー密度による別表現 9
 長澤壯之 (埼玉大理工)
 Simon Blatt (Paris Lodron Univ. Salzburg) Alternative representation of the decomposed Möbius energies with the Möbius invariant densities
 Takeyuki Nagasawa (Saitama Univ.)

概要 The Möbius energy, defined by O'Hara, is one of the knot energies, and named after the Möbius invariant property which was shown by Freedman–He–Wang. It is also known that the energy can be decomposed into three parts keeping the Möbius invariance, proved by Ishizeki–Nagasawa. Though the decomposed energies are Möbius invariant, their densities are not. In this talk, the authors announce that the decomposed energies have alternative representation with the Möbius invariant densities. Using the fact that the cross ratio is invariant under the Möbius transformation, we define new Möbius invariant energies whose densities can be written by the cross ratio. Furthermore we show that these coincide with the decomposed energies.

- 28 S. Blatt (Paris Lodron Univ. Salzburg) Möbius エネルギーの Möbius 不変な離散化と分解 9
 長澤壯之 (埼玉大理工)
 Simon Blatt (Paris Lodron Univ. Salzburg) A Möbius invariant discretization and decomposition of the Möbius energy
 Takeyuki Nagasawa (Saitama Univ.)

概要 The Möbius energy, defined by O'Hara, is named after the Möbius invariant property which was shown by Freedman–He–Wang. The energy can be decomposed into three parts, each of which is Möbius invariant, proved by Ishizeki–Nagasawa. Several discrete versions of Möbius energy, that is, corresponding energies for polygons, are known, and it showed that they converge to the continuum version as the number of vertices to infinity. However already-known discrete energies lost the property of Möbius invariance, nor the Möbius invariant decomposition. Here a new discretization of the Möbius energy is proposed. It has the Möbius invariant property, and can be decomposed into the Möbius invariant components which converge to the original components of decomposition in the continuum limit.

- 29 高野耕太 (埼玉大理工) 一般化された平均曲率ベクトルの積分平均極限を用いた幾何学的表示 . . . 9
 Kouta Takano (Saitama Univ.) A geometric representation of the generalized mean curvature vector using a limit of integral means

概要 The generalized mean curvature is usually represented by the first variation of the varifold. A limit of integral means of the discretization of the classical mean curvature vector suggests a new representation of the generalized mean curvature without using the variation. A similar idea applicable to varifolds satisfying some regularity conditions. As a result, a new geometric representation of the generalized mean curvature is obtained, which has geometric meaning. The approximate tangent space of the measure on Euclidean space is used, however the tangential element of the varifold is not needed for our representation. Consequently, it has the advantage of giving rise to a definition of the generalized mean curvature vector for general measure on Euclidean space.

- 30 佐藤 龍一 (東北大理) The Cauchy problem for the Finsler heat equation 12
 赤木 剛朗 (東北大理)
 石毛 和弘 (東北大理)
 Ryuichi Sato (Tohoku Univ.) The Cauchy problem for the Finsler heat equation
 Goro Akagi (Tohoku Univ.)
 Kazuhiro Ishige (Tohoku Univ.)

概要 Let H be a norm of \mathbf{R}^N and H_0 the dual norm of H . Denote by Δ_H the Finsler–Laplace operator defined by $\Delta_H u := \operatorname{div}(H(\nabla u)\nabla_\xi H(\nabla u))$. In this paper we prove that the Finsler–Laplace operator Δ_H acts as a linear operator to H_0 -radially symmetric smooth functions. Furthermore, we obtain an optimal sufficient condition for the existence of the solution of the Cauchy problem to the Finsler heat equation

$$\partial_t u = \Delta_H u, \quad x \in \mathbf{R}^N, \quad t > 0,$$

where $N \geq 1$ and $\partial_t := \partial/\partial t$.

- 31 関 行宏 (九大数理) Type II blow-up mechanisms in a semilinear heat equation with critical Joseph–Lundgren exponent, Part II: neutral case 12
 Yukihiro Seki (Kyushu Univ.) Type II blow-up mechanisms in a semilinear heat equation with critical Joseph–Lundgren exponent, Part II: neutral case

概要 We are concerned with blow-up mechanisms in a semilinear heat equation

$$u_t = \Delta u + |u|^{p-1}u, \quad x \in \mathbf{R}^N, \quad t > 0,$$

where $p > 1$ is a constant. It is well known that type II blow-up does occur if $N \geq 11$ and $p > p_{JL}$, where p_{JL} stands for the Joseph–Lundgren exponent. I will report a recent result on type II blow-up for $p = p_{JL}$.

- 32 山本 宏子 (明大MIMS) 半線形波動方程式の反応拡散近似 12
 二宮 広和 (明大総合数理)
 Hiroko Yamamoto (Meiji Univ.) Reaction-diffusion approximation of a semilinear wave equation
 Hirokazu Ninomiya (Meiji Univ.)

概要 Reaction-diffusion systems are one of nonlinear parabolic systems and are often used as models described chemical reaction system, combustion system, and so on. If the number of unknown variables of a reaction-diffusion system increases, the dynamics of a reaction-diffusion system may become complicated. In order to study the complexity of the dynamics, we consider the following question: What kinds of systems can we approximate by reaction-diffusion systems? In this talk, we introduce a reaction-diffusion system which approximates a semilinear wave equation. The proof is based on the energy estimates.

- 33 谷口 雅治 (岡山大自然) Allen–Cahn 方程式における角錐形進行波の一意性と安定性 12
 Masaharu Taniguchi (Okayama Univ.) The uniqueness and stability of pyramidal traveling fronts in the Allen–Cahn equations

概要 Traveling front solutions with pyramidal shapes are studied in the Allen–Cahn equation (Nagumo Equation) in the n -dimensional Euclidean space. Here n is any integer that is greater or equal to 3. The existence of pyramidal traveling fronts was shown by [1] and [3]. The uniqueness and stability was shown by [2] for $n = 3$. In this work, I report the uniqueness and stability of pyramidal traveling fronts for n that are greater or equal to 3.

- 34 藤江健太郎 (東京理大理)^b No critical nonlinear diffusion in 1D quasilinear Keller–Segel system ··· 12
T. Cieřlak (IMPAN)
Kentarou Fujie (Tokyo Univ. of Sci.) No critical nonlinear diffusion in 1D quasilinear Keller–Segel system
Tomasz Cieřlak (IMPAN)

概要 This talk deals with the fully parabolic 1D chemotaxis system with diffusion $1/(1+u)$. We prove that the above mentioned nonlinearity, despite being a natural candidate, is not critical. It means that for such a diffusion any initial condition, independently on the magnitude of mass, generates global-in-time solution. In view of our theorem one sees that one-dimensional Keller–Segel system is essentially different than its higher-dimensional versions. In order to prove our theorem we establish a new Lyapunov-like functional associated to the system. The information we gain from our new functional (together with some estimates based on the well-known old Lyapunov functional) turns out to be rich enough to establish global existence for the initial-boundary value problem.

- 35 永井敏隆 (広島大*) Boundedness of solutions to a parabolic-elliptic chemotaxis model in \mathbb{R}^2
山田哲也 (福井工高専) with critical mass ··········· 12
Toshitaka Nagai (Hiroshima Univ.*) Boundedness of solutions to a parabolic-elliptic chemotaxis model in \mathbb{R}^2
Tetsuya Yamada
(Fukui Nat. Coll. of Tech.) with critical mass

概要 We consider the Cauchy problem of a parabolic-elliptic chemotaxis model in \mathbb{R}^2 . Our purpose is to discuss the boundedness of nonnegative solutions to the Cauchy problem in the critical case where the total mass of the initial data is 8π .

- 36 Xinru Cao (Paderborn Univ.) Global existence and stabilization in a 3D two-species chemotaxis-Stokes
来間俊介 (東京理大理) system with competitive kinetics ··········· 12
水上雅昭 (東京理大理)
Xinru Cao (Paderborn Univ.) Global existence and stabilization in a 3D two-species chemotaxis-Stokes
Shunsuke Kurima (Tokyo Univ. of Sci.) system with competitive kinetics
Masaaki Mizukami
(Tokyo Univ. of Sci.)

概要 This talk deals with a 3D two-species chemotaxis-Stokes system with competitive kinetics. A single-species case was studied by e.g., Winkler (2012), Tao–Winkler (2015, 2016) and Lankeit (2016). However, there has not been rich results on coupled two-species-fluid systems. Recently, Hirata–K.–Mizukami–Yokota (2017) proved global existence of classical solutions for a 2D two-species chemotaxis-Navier–Stokes system. The present work asserts global existence and behaviour of classical solutions for the case of two species in 3D.

- 37 水上雅昭 (東京理大理) Improvement of conditions for asymptotic stability in a fully parabolic
two-species chemotaxis-competition model ··········· 12
Masaaki Mizukami
(Tokyo Univ. of Sci.) Improvement of conditions for asymptotic stability in a fully parabolic
two-species chemotaxis-competition model

概要 This talk is concerned with asymptotic behavior of solutions to a fully parabolic two-species chemotaxis-competition model. Bai and Winkler proved asymptotic behavior in the system under some conditions and special setting in 2016. Recently, the conditions assumed in the previous work were improved (M., 2017); however, this result did not give a complete improvement. The main result of this talk asserts complete improvement of the conditions assumed in Bai–Winkler (2016) and M. (2017) for asymptotic stability.

13:00~14:00 特別講演

- 足達 慎二 (静岡大工) 半線形楕円型方程式の正值解の一意性・非退化性とその応用
 Shinji Adachi (Shizuoka Univ.) On the uniqueness and the non-degeneracy of positive solutions for a class of semilinear elliptic equations in \mathbb{R}^N and its applications

概要 We survey some recent results for the uniqueness and the non-degeneracy of positive solutions for a class of semilinear elliptic equations in \mathbb{R}^N . Especially, we show the uniqueness and the non-degeneracy of positive solutions in the cases where nonlinear terms may have sublinear growth at infinity. As applications, we obtain the uniqueness and the non-degeneracy of positive solutions for some quasilinear elliptic equations.

9月13日(水) 第VIII会場

9:15~12:00

- 38 熊谷 大雅 (早大教育) Asymptotics of the solutions of Hamilton–Jacobi equations with large drift term 9
 Taiga Kumagai (Waseda Univ.) Asymptotics of the solutions of Hamilton–Jacobi equations with large drift term

概要 In this talk, we consider the asymptotic behavior of solutions to Hamilton–Jacobi equations with large drift term in an open subset of two dimensional Euclidean space, where the set is determined through a Hamiltonian and the drift is given by the Hamiltonian vector field. Under some growth assumptions on the Hamiltonian, in the case where the Hamiltonian has degenerate critical points, we establish the convergence of solutions of the Hamilton–Jacobi equations and identify the limit of the solutions as the solution of systems of ordinary differential equations on a graph. The graph has many line segments more than four at a node.

- 39 小杉 卓裕 (東北大理)^b Equivalence of viscosity solutions between obstacle problems and gradient constraint problems 12
 Takahiro Kosugi (Tohoku Univ.) Equivalence of viscosity solutions between obstacle problems and gradient constraint problems

概要 It is showed that the unique viscosity solutions of fully nonlinear elliptic partial differential equations under gradient constraint coincides with that of the equation with suitably selected bilateral obstacles. To this end, it is necessary to obtain the Lipschitz estimates on viscosity solutions of bilateral obstacle problems.

- 40 水谷 治哉 (阪大理) 尺度臨界な Schrödinger 作用素に対するレゾルベント評価とその応用 II 12
 Haruya Mizutani (Osaka Univ.) Resolvent estimates for scaling-critical Schrödinger operators and applications II

概要 We discuss a recent progress on uniform resolvent estimates for Schrödinger operators with scaling-critical potentials and their applications to global-in-time Strichartz estimates for the Cauchy problem of the Schrödinger equation and Keller type eigenvalue bounds for non-self-adjoint Schrödinger operators with complex-valued potentials.

- 41 太田 雅人 (東京理大理) Strong instability of standing waves for nonlinear Schrödinger equations with a partial confinement 12
 Masahito Ohta (Tokyo Univ. of Sci.) Strong instability of standing waves for nonlinear Schrödinger equations with a partial confinement

概要 We study the instability of standing wave solutions for nonlinear Schrödinger equations with a one-dimensional harmonic potential in dimension $N \geq 2$. We prove that if the nonlinearity is L^2 -critical or supercritical in dimension $N - 1$, then any ground states are strongly unstable by blowup.

- 42 星 埜 岳 (阪 大 理) Analyticity of solutions to the nonlinear Schrödinger equations without gauge invariance 9

Gaku Hoshino (Osaka Univ.) Analyticity of solutions to the nonlinear Schrödinger equations without gauge invariance

概要 We study analyticity of global solutions to NLS without gauge invariance.

- 43 星 埜 岳 (阪 大 理) Analytic smoothing effect for global solutions to a system of Schrödinger equations with large data 9

Gaku Hoshino (Osaka Univ.) Analytic smoothing effect for global solutions to a system of Schrödinger equations with large data

概要 We study analytic smoothing effect for a system of Schrödinger equations.

- 44 百 名 亮 介 (早 大 理 工)* Hartree 方程式のある種の適切性について 9
星 埜 岳 (阪 大 理)

Ryosuke Hyakuna (Waseda Univ.) A well-posedness result for the Hartree equation
Gaku Hoshino (Osaka Univ.)

概要 In this talk, we consider the Cauchy problem of the Hartree equation and we discuss the local existence when data are not characterized by any kind of square integrability.

- 45 木 下 真 也 (名 大 多 元 数 理) Local well-posedness for the Cauchy problem of the Klein–Gordon–Zakharov system in two and three dimensions 12

Shinya Kinoshita (Nagoya Univ.) Local well-posedness for the Cauchy problem of the Klein–Gordon–Zakharov system in two and three dimensions

概要 We consider the Cauchy problem of 2D and 3D Klein–Gordon–Zakharov system with very low regularity initial data. We prove the bilinear estimates which are crucial to get the local in time well-posedness. The estimates are established by the Fourier restriction norm method. We utilize the nonlinear version of the classical Loomis–Whitney inequality.

- 46 佐 川 侑 司 (阪 大 理) A sharp lower bound for the lifespan of small solutions to the Schrödinger
砂 川 秀 明 (阪 大 理) equation with a subcritical power nonlinearity 12
保 田 舜 介

Yuji Sagawa (Osaka Univ.) A sharp lower bound for the lifespan of small solutions to the Schrödinger
Hideaki Sunagawa (Osaka Univ.) equation with a subcritical power nonlinearity
Shunsuke Yasuda

概要 Let T_ε be the lifespan for the solution to the Schrödinger equation on \mathbb{R}^d with a subcritical power nonlinearity and the initial data in the form $\varepsilon\varphi(x)$. We provide a sharp lower bound estimate for T_ε as $\varepsilon \rightarrow +0$ which can be written explicitly by the initial data and the nonlinearity. This is an improvement of the previous result by H. Sasaki [Adv. Diff. Eq. 14 (2009), 1021–1039].

- 47 岸 本 展 (京 大 数 理 研) Ill-posedness of the third order NLS equation with Raman scattering
堤 誉 志 雄 (京 大 理) term 12

Nobu Kishimoto (Kyoto Univ.) Ill-posedness of the third order NLS equation with Raman scattering
Yoshio Tsutsumi (Kyoto Univ.) term

概要 We consider the nonlinear Schrödinger equation with third order dispersion and intrapulse Raman scattering term. Without the Raman scattering term, the associated Cauchy problem is known to be locally well-posed in Sobolev spaces. We show that the Raman scattering term causes the ill-posedness of the Cauchy problem (nonexistence of local-in-time solutions) in Sobolev spaces. We also mention the unique solvability of the Cauchy problem in the analytic function space.

- 48 岡本 葵 (信州大工) 確率化された初期値をもつエネルギー臨界非線形 Schrödinger 方程式の
Tadahiro Oh (Univ. of Edinburgh) 初期値問題の可解性 12
O. Pocovnicu (Heriot-Watt Univ.)
Mamoru Okamoto (Shinshu Univ.) Probabilistic Cauchy problem for the energy critical nonlinear Schrödinger
Tadahiro Oh (Univ. of Edinburgh) equations
Oana Pocovnicu (Heriot-Watt Univ.)

概要 We consider the Cauchy problem for the nonlinear Schrödinger equations (NLS) with non-algebraic nonlinearities on the Euclidean space. In particular, we study the energy-critical NLS in \mathbb{R}^d ($d = 5, 6$) and energy-critical NLS without gauge invariance and prove that they are almost surely locally well-posed with respect to randomized initial data below the energy space.

- 49 青木 和貴 (阪大理) 4階非線形 Schrödinger 方程式の時間大域解の存在について 12
Kazuki Aoki (Osaka Univ.) Global existence of solutions for fourth-order nonlinear Schrödinger equations

概要 We consider the Cauchy problem for the fourth-order nonlinear Schrödinger equations with supercritical power nonlinearities. The class of the fourth-order nonlinear Schrödinger equations describe deep water wave dynamics. We prove global existence of small solutions in one or two space dimensions. This is an improvement of the result by N. Hayashi, J. A. Mendez-Navarro, and P. I. Naumkin. [Commun. Contemp. Math. 18(3), 1550035, 24 pp (2016)]

- 50 瓜屋 航太 (岡山理大理)* 3次元における斉次型臨界非線形項をもつ非線形 Schrödinger 方程式の解
眞崎 聡 (阪大基礎工) の長距離散乱について 12
宮崎 隼人 (津山工高専)
Kota Uriya (Okayama Univ. of Sci.) Long range scattering for nonlinear Schrödinger equations with critical
Satoshi Masaki (Osaka Univ.) homogeneous nonlinearity in three dimensions
Hayato Miyazaki
(Tsuyama Nat. Coll. of Tech.)

概要 We consider the final state problem for the nonlinear Schrödinger equation with a homogeneous nonlinearity of the critical order which is not necessarily a polynomial in three dimensions. We give a sufficient condition on the nonlinearity for that the corresponding equation admits a solution that behaves like a free solution with or without a logarithmic phase correction. This is the extension of the previous result in one and two dimensions. Moreover, we present a candidate of the second asymptotic profile to the solution.

- 51 橋本伊都子 多次元空間上におけるバーガーズ方程式の球対称問題の漸近挙動について 9
(関西大システム理工・阪市大数学研)
松村 昭孝 (阪大情報)
Itsuko Hashimoto Asymptotic behavior toward nonlinear waves for radially symmetric solutions of multi-dimensional Burgers equation
(Kansai Univ./Osaka City Univ.)
Akitaka Matsumura (Osaka Univ.)

概要 This talk is concerned with the asymptotic behaviors of radially symmetric solutions for multi-dimensional Burgers equation on the exterior domain in n -dimensional space, where the boundary and far field conditions are prescribed. In a case where the corresponding 1-D Riemann problem for the non-viscous part admits a shock wave, we show the solution tends toward a superposition of stationary wave and rarefaction wave as time goes to infinity, and also show the decay rate estimate. Furthermore, for $n = 3$, we give the complete classification of the asymptotic states, which includes even a superposition of stationary wave and viscous shock wave.

12:15~12:30 2017年度日本数学会解析学賞授賞式

14:15~16:15

- 52 側島基宏 (東京理大理工)* Diffusion phenomena for the wave equation with space-dependent damping term growing at infinity 9
 若杉勇太 (愛媛大理工) Motohiro Sobajima Diffusion phenomena for the wave equation with space-dependent damping term growing at infinity
 (Tokyo Univ. of Sci.)
 Yuta Wakasugi (Ehime Univ.)

概要 We study the asymptotic behavior of solutions to the wave equation with damping depending on the space variable and growing at the spatial infinity. We prove that the solution is approximated by that of the corresponding heat equation as time tends to infinity.

- 53 側島基宏 (東京理大理工) Weighted energy estimates for wave equation with space-dependent damping term for slowly decaying initial data 9
 若杉勇太 (愛媛大理工) Motohiro Sobajima Weighted energy estimates for wave equation with space-dependent damping term for slowly decaying initial data
 (Tokyo Univ. of Sci.)
 Yuta Wakasugi (Ehime Univ.)

概要 In this talk we consider the wave equation with space-dependent damping coefficient $a(x) = |x|^{-\alpha}$ ($\alpha \in [0, 1)$) in an exterior domain Ω having a smooth boundary. Weighted energy estimates with weight functions like polynomials are given and these decay rate are almost sharp, even when the initial data do not have compact support in Ω . The crucial idea is to use special solution of $\partial_t u = |x|^\alpha \Delta u$ with a polynomial decay.

- 54 片山聡一郎 (阪大理) A note on the decay property for dissipative nonlinear wave equations in one space dimension 9
 若狭恭平 (室蘭工大工) B. Yordanov (北大国際連携機構)
 Soichiro Katayama (Osaka Univ.) A note on the decay property for dissipative nonlinear wave equations in one space dimension
 Kyouhei Wakasa (Muroran Inst. of Tech.)
 Borislav Yordanov (Hokkaido Univ.)

概要 In this talk, we consider the Cauchy problem for the dissipative nonlinear wave equation in one space dimension. In the work of the dissipative nonlinear wave equation, it is well-studied by using the energy estimates. The purpose of this talk is to show the point-wise estimates of solutions. Such kind of estimates describe the characteristics of the wave equation and precious information. Making use of these estimates, we can also get the result of the energy decay.

- 55 吉田夏海 Decay properties of solutions toward a multiwave pattern for the scalar conservation law with the Ostwald–de Waele-type viscosity 12
 (立命館大OIC総合研究機構) Natsumi Yoshida (Ritsumeikan Univ.) Decay properties of solutions toward a multiwave pattern for the scalar conservation law with the Ostwald–de Waele-type viscosity

概要 We study the asymptotic decay of solutions toward a multiwave pattern (rarefaction wave and viscous contact wave) of the Cauchy problem for the one-dimensional viscous conservation law where the far field states are prescribed. Especially, we deal with the case when the flux function is convex or concave but linearly degenerate on some interval, and also the viscosity is a nonlinearly degenerate one. The proof is given by a technical time-weighted energy method and the careful estimates for the interactions between the nonlinear waves.

- 56 吉田夏海 (立命館大 OIC 総合研究機構) Large time behavior of solutions toward viscous shock waves to the Cauchy problem for the scalar conservation law with nonlinear flux and viscosity 12

Natsumi Yoshida (Ritsumeikan Univ.) Large time behavior of solutions toward viscous shock waves to the Cauchy problem for the scalar conservation law with nonlinear flux and viscosity

概要 We study the large time asymptotics of solutions to the Cauchy problem for the one-dimensional scalar viscous conservation law where the far field states are prescribed. Especially, we deal with the case when the flux function is a non-convex nonlinear function, and also the viscosity is a nonlinear function. The proof is given using a technical weighted energy method associated with the nonlinearity of the flux and the viscosity.

- 57 杉山裕介 (東京理大理) On the global existence of weak solutions to 1D degenerate quasilinear wave equations 12
Yunguang Lu (Hangzhou Normal Univ.)

Yusuke Sugiyama (Tokyo Univ. of Sci.) On the global existence of weak solutions to 1D degenerate quasilinear wave equations
Yunguang Lu (Hangzhou Normal Univ.)

概要 We consider the existence and the nonexistence of global generalized (nonnegative) solutions of the nonlinearly degenerate wave equations $\partial_t^2 u = \partial_x(u^{2a}\partial_x u)$ with the nonnegative initial data $u_0(x) \geq 0$ and $a > 0$. This result is an extension of the results in the second author's paper, where the existence and the nonexistence of the unique global classical solution were studied with a threshold on $\int_{-\infty}^{\infty} u_1(x)dx$ and the non-degeneracy condition $u_0(x) \geq \delta_0 > 0$ on the initial data.

- 58 肥田野久二男 (三重大教育)* Global existence for a system of quasi-linear wave equations in 3D satisfying the weak null condition 12
横山和義 (北海道科学大工)
Kunio Hidano (Mie Univ.) Global existence for a system of quasi-linear wave equations in 3D satisfying the weak null condition
Kazuyoshi Yokoyama (Hokkaido Univ. of Sci.)

概要 We show global existence of small solutions to the Cauchy problem for a system of quasi-linear wave equations in three space dimensions. The feature of the system lies in that it satisfies the weak null condition, though we permit the presence of some quadratic nonlinear terms which do not satisfy the null condition. Due to the presence of such quadratic terms, the standard argument no longer works for the proof of global existence. To get over this difficulty, we extend the ghost weight method of Alinhac so that it works for the system under consideration.

- 59 星賀 彰 (静岡大工)* 伝播速度が異なる 2 次元準線形波動方程式系の滑らかな解のライフスパンについて 9
Akira Hoshiga (Shizuoka Univ.) Lower bound of the lifespan of solutions to systems of quasi-linear wave equations with multiple propagation speeds

概要 We consider the Cauchy problem of systems of quasi-linear wave equations in 2-dimensional space. We assume that the propagation speeds are distinct and that the nonlinearities contain quadratic and cubic terms of the first and second order derivatives of the solution. We know that if the all quadratic and cubic terms of nonlinearities satisfy strong null-condition, then there exists a global solution for sufficiently small initial data. In this paper, we study about the lifespan of the smooth solution, when the cubic terms in the quasi-linear nonlinearities do not satisfy the strong null-condition.

- 60 佐々木多希子 (明大理工) 非線形波動方程式系の爆発曲線について 9
Takiko Sasaki (Meiji Univ.) The blow-up curve for a system of nonlinear wave equations

概要 We study a blow-up curve for a system of nonlinear wave equations. The purpose of this talk is to show that the blow-up curve is a C^1 curve if the initial values are large and smooth enough. To prove the result, we convert the system into a first order system, and then apply a modification of the method of Caffarelli and Friedman (1986).

- 61 小池 開 自由分子流中の物体運動に平面壁が与える影響 12
(慶大理工・理化学研 AIP)

Kai Koike (Keio Univ./RIKEN) Wall effect on the motion of a body immersed in a free molecular flow

概要 Satellites, International Space Station, orbital debris all move in highly rarefied atmosphere. This motivates the study of fluid-structure interaction in highly rarefied gas. Caprino et al. [Comm. Math. Phys., **264** (2006), 167–189] analyzed 1-D motion of a rigid body in a free molecular flow. They proved algebraic convergence of the body's velocity $V(t)$ to the terminal velocity V_∞ . My question is whether the asymptotic behavior changes if there is a fixed wall behind the moving body. Caprino et al. considered the motion in the whole space \mathbb{R}^d ; I considered the motion in the half space \mathbb{R}_+^d . I proved that the approach to the terminal velocity changes despite the fact that the body and the fixed wall go away infinitely.

16:30~17:30 特別講演

伊藤 弘道 (東京理大理) き裂を含む領域における偏微分方程式の解析

Hiromichi Itou (Tokyo Univ. of Sci.) On analysis for partial differential equations in a cracked domain

概要 Theory of partial differential equations has been developed mainly in smooth domains. In non-smooth domains such as polyhedral or cracked domains, mathematical difficulties appear because domains have singular points. Then, it is important to analyze the precise behavior of the solution of the partial differential equations with boundary conditions near the singular points. This kind of analysis has possibility of application in various fields of science and engineering such as fracture problems, inverse problems (nondestructive evaluation) and so on.

In this talk, first we have an overview of crack problems from both sides of mathematics and mechanics. Second we introduce some convergent series expansions of solutions of a boundary value problem at a crack tip in a linearized elasticity model. Next, we note that the study of cracks within the context of the linearized theory of elasticity has a drawback due to an inconsistency with regard to the strain, namely the strain becomes infinite at the crack tip. Then we consider a boundary value problem in a nonlinear elastic body that exhibits limiting small strain, which does not suffer from the inconsistency. For this problem we introduce the concept of a non-smooth viscosity solution, called generalized solution, which is described as generalized variational inequalities and coincides with the weak solution in the smooth case. Lastly, we mention future and ongoing researches in crack problems and their applications.

9月14日(木) 第VIII会場

9:00~12:00

- 62 高村博之 (公立はこだて未来大) 劣 Strauss 指数をもつ半線形消散波動方程式の解の爆発と lifespan 評価
 頼宇安 (Lishui Univ.) 12
Hiroyuki Takamura Blow-up and lifespan estimate of solutions of semilinear damped wave
 (Future Univ. Hakodate) equations with sub-Strauss exponent
 Lai Ning-An (Lishui Univ.)

概要 It is well-known that the critical exponent for semilinear damped wave equations is Fujita exponent when the damping is effective. Introducing a multiplier for the time-derivative of the spatial integral of unknown functions, we succeed to employ the analysis on semilinear wave equations and to prove a blow-up result for semilinear damped wave equations with sub-Strauss exponent when the damping is in the scattering range.

- 63 加藤正和 (室蘭工大工) 空間3次元上の時間減衰する重み付き非線形波動方程式の解の時間大域
 櫻庭みく 存在と爆発 12
Masakazu Kato Global existence and blow-up for wave equations with weighted nonlin-
 (Muroran Inst. of Tech.) ear term in 3D
 Miku Sakuraba

概要 We consider the global existence and blow-up of small data solutions for the wave equation with weighted nonlinear term in three space dimensions. We obtain upper and lower bound of the lifespan of solutions to the problem.

- 64 今井啄人 (公立はこだて未来大) 空間2次元におけるスケール不変な半線形消散波動方程式の解のライフ
 高村博之 (公立はこだて未来大) スパン評価 12
 加藤正和 (室蘭工大工)
 若狭恭平 (室蘭工大工)
Takuto Imai (Future Univ. Hakodate) The lifespan of solutions to semilinear damped wave equations with
 Hiroyuki Takamura scale invariant in two space dimension
 (Future Univ. Hakodate)
 Masakazu Kato
 (Muroran Inst. of Tech.)
 Kyouhei Wakasa
 (Muroran Inst. of Tech.)

概要 In this talk, we report the sharp upper and lower bound of the lifespan of solutions to semilinear damped wave equations in the scale invariant case with a special constant in two space dimensions. The result is divided into two cases up to the total integral of the sum of the initial data.

- 65 寺本有花 (九大数理) Bifurcation of the compressible Taylor vortex 12
 隠居良行 (九大数理)
 西田孝明 (京大情報)
 Yuka Teramoto (Kyushu Univ.) Bifurcation of the compressible Taylor vortex
 Yoshiyuki Kagei (Kyushu Univ.)
 Takaaki Nishida (Kyoto Univ.)

概要 We consider the Couette–Taylor problem, a flow between two concentric cylinders, whose inner cylinder is rotating with uniform speed and the outer one is at rest. If the rotating speed is sufficiently small, the Couette flow (laminar flow) is stable. When the rotating speed increases, beyond a certain value of the rotating speed, a vortex flow pattern (Taylor vortex) appears. Mathematically, this phenomena can be formulated as a bifurcation problem. In this talk the Couette–Taylor problem is considered for the compressible Navier–Stokes equation and the bifurcation of the Taylor vortex is proved when the Mach number is sufficiently small.

- 66 相木雅次 (東京理大理工) Motion of a vortex filament on a slanted plane 12
 Masashi Aiki (Tokyo Univ. of Sci.) Motion of a vortex filament on a slanted plane

概要 We consider a nonlinear model equation, known as the localized induction equation, describing the motion of a vortex filament immersed in an incompressible and inviscid fluid. The talk will report on the global-in-time unique solvability of an initial-boundary value problem describing the motion of a vortex filament on a slanted plane. The proof relies on the careful analysis of the shape of the filament near the boundary, and this will be the main focus of this talk.

- 67 李煥元 (東大数理) 3次元非斉次磁気流体方程式の大域的強解 12
 Huanyuan Li (Univ. of Tokyo) Global strong solutions to the three dimensional density-dependent MHD equations

概要 In this talk, we consider an initial boundary value problem for the three dimensional nonhomogeneous incompressible magnetohydrodynamic equations with density-dependent viscosity and resistivity coefficients over a bounded smooth domain. Global in time unique strong solution is proved to exist when the initial vorticity and current density are both suitably small in some Sobolev space with arbitrary large initial mass density, and the vacuum of initial density is also allowed.

- 68 杉山裕介 (東京理大理) 変数係数摩擦項を持つ1次元 Euler 方程式の解の爆発 12
 Yusuke Sugiyama (Tokyo Univ. of Sci.) Blow-up of solutions to the 1D Euler equation with valuable damping coefficient

概要 We consider a blow-up problem for the 1D Euler equation with time and space dependent damping. We investigate sufficient conditions on initial data and the rate of spatial or time-like decay of the coefficient of damping for the occurrence of the finite time blow-up. In particular, our sufficient conditions ensure that the derivative blow-up occurs in finite time with the solution itself and the pressure bounded. Our method is based on simple estimates with Riemann invariant.

- 69 ブレジナヤン (東工大理工) On measure-valued solutions to the complete Euler system 12
 E. Feireisl (Czech Acad. of Sci.)
Jan Březina (Tokyo Tech) On measure-valued solutions to the complete Euler system
 Eduard Feireisl (Czech Acad. of Sci.)

概要 We introduce the concept of dissipative measure-valued solution to the complete Euler system describing the motion of an inviscid compressible fluid. These solutions are characterized by a parameterized (Young) measure and a dissipation defect in the total energy balance. A dissipative measure-valued solution can be seen as the most general concept of solution to the Euler system retaining its structural stability. In particular, we show that a dissipative measure-valued solution necessarily coincides with a classical one on its life span provided they share the same initial data.

- 70 渡邊圭市 (早大理工) Compressible-incompressible two phase flow of Korteweg type with phase transition: model problem 9
 Keiichi Watanabe (Waseda Univ.) Compressible-incompressible two phase flow of Korteweg type with phase transition: model problem

概要 In this talk, we show the existence of \mathcal{R} -bounded solution operator families for a resolvent problem arising from the motion, where one fluid is a capillary compressible viscous flow and the other is an incompressible viscous flow. Moreover, we show that the regularity of density is W_q^3 with respect to the space variable, although it is W_q^1 in the usual case.

- 71 梶原直人 (東大数理) 最大正則性理論を用いた phase-field Navier–Stokes 方程式の可解性及び定常解の安定性について 12
 Naoto Kajiwara (Univ. of Tokyo) Solvability and stability for the phase-field Navier–Stokes equations via maximal regularity

概要 We study the dynamics of vesicle membranes in incompressible viscous fluids. We prove existence and uniqueness of the local strong solution for this model coupling of the Navier–Stokes equations with a phase field equation in an L_p - L_q setting via maximal regularity. Moreover we have that the solution is real analytic in time and space. It is also shown that the variational strict stable solution is exponentially stable, provided the product of the viscosity coefficient and the mobility constant is large.

- 72 中村憲史 (筑波大数理物質) 双曲型 Stokes 方程式の解に対する局所エネルギー減衰定理 9
小林孝行 (阪大基礎工)
久保隆徹 (筑波大数理物質)
Kenji Nakamura (Univ. of Tsukuba) Local energy decay estimate of solutions to the hyperbolic type Stokes equations
Takayuki Kobayashi (Osaka Univ.)
Takayuki Kubo (Univ. of Tsukuba)

概要 In this talk, we discuss a local energy decay estimate of solutions to the initial-boundary value problem for the hyperbolic type Stokes equations of incompressible fluid flow.

- 73 齋藤平和 (早大理工) Maximal L_p - L_q regularity for a compressible fluid model of Korteweg type on general domains 9
 Hirokazu Saito (Waseda Univ.) Maximal L_p - L_q regularity for a compressible fluid model of Korteweg type on general domains

概要 In this talk, we would like to consider a linearized system on general domains Ω arising from a compressible fluid model of Korteweg type. The boundary of Ω consists of two parts S, Γ with $\text{dist}(S, \Gamma) > 0$. One imposes the free boundary condition on Γ , while the non-slip condition on S . It is admissible that $S = \emptyset$ or $\Gamma = \emptyset$ in this talk. We show the maximal L_p - L_q regularity for the linearized system.

- 74 古川 賢 (東大数理)^b 3次元 Oseen 渦型 Navier–Stokes 流の漸近安定性 …………… 9
 Ken Furukawa (Univ. of Tokyo) On asymptotic stability of the three dimensional Oseen type Navier–Stokes flow

概要 In this talk, we construct three dimensional Oseen type Navier–Stokes flows in Euclidean space and vertically periodic space and show their asymptotic stability in vertically periodic space under large initial perturbation.

- 75 鶴見 裕之 (早大理工) Extension criterion via partial components of vorticity on strong solutions to the Navier–Stokes equations in higher dimensions …………… 12
 Hiroyuki Tsurumi (Waseda Univ.) Extension criterion via partial components of vorticity on strong solutions to the Navier–Stokes equations in higher dimensions

概要 We consider the extension criterion of strong solutions to the Navier–Stokes equations in \mathbb{R}^N . It is proved that among $\frac{N(N-1)}{2}$ components of the vorticity, $[\frac{N}{2}]$ components are negligible for the criterion whether the time local solutions can be extended beyond the critical time. Our result may be regarded as generalization to the higher dimensional case of Chae–Choe and Kozono–Yatsu in the 3D case which showed that only two components in L^q , $\frac{3}{2} < q \leq \infty$, of the vorticity contribute to such an extension criterion. Furthermore, the critical case $q = \infty$ originally treated by Kato–Ponce in \mathbb{R}^N is also generalized in such a way that $[\frac{N}{2}]$ components of vorticity are redundant for the extension criterion.

- 76 澤田 宙広 (岐阜大工) 空間無限遠方で1次増大する初期値に対するプリミティブ方程式の局所可解性 …………… 12
 Okihiro Sawada (Gifu Univ.) Locally-in-time well-posedness of the primitive equations with linearly growing initial data

概要 The primitive equations with linearly growing initial data in the horizontal direction are concerned. The existence theory of mild solutions is established in certain interpolation spaces. A semi-group of Ornstein–Uhlenbeck type is investigated in Lebesgue spaces, including its smoothing property. Constructing mild solutions, the fixed point arguments of Fujita–Kato type are used.

14:15~16:15

- 77 筒井 容平 (信州大理) Asymptotic stability of stationary solutions to the Navier–Stokes equations in Besov spaces …………… 9
 岡部 孝宏 (弘前大教育) J. M. Cunanan (埼玉大理工)
 Youhei Tsutsui (Shinshu Univ.) Asymptotic stability of stationary solutions to the Navier–Stokes equations in Besov spaces
 Takahiro Okabe (Hirosaki Univ.) Jayson Mesitas Cunanan (Saitama Univ.)

概要 We give an asymptotic stability result for the incompressible Navier–Stokes equations in Besov spaces with sub/super critical smoothness. Following the argument by Kozono–Yamazaki (1995), we establish smoothing estimates for the semigroup generated by the Laplacian with a perturbation. Applying that, a critical estimate is proved, which is the main ingredient for the proof of the stability result.

- 78 小 菌 英 雄 (早大基幹理工)* Finite energy of generalized suitable weak solutions to the Navier–Stokes equations and the Liouville type theorem 9
 寺 澤 祐 高 (名大多元数理)
 若 杉 勇 太 (愛媛大理工)
 Hideo Kozono (Waseda Univ.) Finite energy of generalized suitable weak solutions to the Navier–Stokes equations and the Liouville type theorem
 Yutaka Terasawa (Nagoya Univ.)
 Yuta Wakasugi (Ehime Univ.)

概要 Introducing a new notion of generalized suitable weak solutions, we first prove validity of the energy inequality for such a class of weak solutions to the Navier–Stokes equations in the whole space \mathbb{R}^n . Although we need certain growth condition on the pressure, we may treat the class even with infinite energy quantity except for the initial velocity.

- 79 菱 田 俊 明 (名大多元数理)^b Large time behavior of a generalized Oseen evolution operator, with applications to the Navier–Stokes flow past a rotating obstacle 12
 Toshiaki Hishida (Nagoya Univ.) Large time behavior of a generalized Oseen evolution operator, with applications to the Navier–Stokes flow past a rotating obstacle

概要 Consider the motion of a viscous incompressible fluid in a 3D exterior domain when a rigid body moves with a prescribed time-dependent translational and angular velocities. We develop decay estimates of the evolution operator, which provides a solution of the linearized non-autonomous system, and then apply them to the Navier–Stokes initial value problem.

- 80 菱 田 俊 明 (名大多元数理)^b Navier–Stokes flow past a rigid body: attainability of steady solutions as limits of unsteady weak solutions 12
 P. Maremonti (Campania 大)
 Toshiaki Hishida (Nagoya Univ.) Navier–Stokes flow past a rigid body: attainability of steady solutions as limits of unsteady weak solutions
 Paolo Maremonti
 (Univ. degli Studi della Campania)

概要 Consider the Navier–Stokes flow in 3D exterior domains, where the translational velocity of the body becomes $-u_\infty$ after some finite time. For the starting problem raised by Finn, we study some generalized situation in which unsteady solutions start from large motions being in L^3 . We then conclude that the steady solutions are still attainable as limits of evolution of those fluid motions provided u_∞ is small enough.

- 81 柴 田 良 弘 (早大理工) Local well-posedness for the Magnetohydrodynamics in the different two liquids case 9
 Yoshihiro Shibata (Waseda Univ.) Local well-posedness for the Magnetohydrodynamics in the different two liquids case

概要 In this talk, I would like to talk about local well-posedness for the magnetohydrodynamic equations in the two different liquids case. Since two divergence free conditions are over determined, I consider the jump condition for the divergence of magnetic field on the interface. This is a new aspect in treating the MHD. After transforming the time dependent unknown domain to the reference domain by Lagrange transformation, using the maximal L_p - L_q regularity for the linearized equations we prove a local in time unique existence theorem.

- 82 柴田良弘 (早大理工) Maximal L_p - L_q regularity theorem for the linearized electro-magnetic field equations 9

Yoshihiro Shibata (Waseda Univ.) Maximal L_p - L_q regularity theorem for the linearized electro-magnetic field equations

概要 In this talk, I will present the maximal L_p - L_q regularity theorem for the linearized electro-magnetic field equations. In solving MHD equations, linearized equations are decoupled, because the coupling terms are semi-linear. Thus, as linearized equations, we treat Stokes equations with interface conditions and linear electro-magnetic equations with interface conditions. Stokes equations with interface conditions have been studied by Pruess and Shimonett, and Maryani and Saito, so that I treat only the linear electro-magnetic equations with interface conditions.

- 83 柴田良弘 (早大理工) Local well-posedness for the two component flow 9
Yoshihiro Shibata (Waseda Univ.) Local well-posedness for the two component flow

概要 In this talk, I would like to talk about local in time unique existence theorem for the two components flow. I used the modelling due to Vicent Giovangigli: Multicomponent flow modeling, Birkhäuser. After transforming the equations to one component case by using the Giovangigli transformation, I proved the local well-posedness for the resultant system of equations by using Lagrange transformation and the maximal L_p - L_q regularity theorem for the linearized equations. This is a joint work with Ewelina Zatorska (Imperial College of London).

- 84 柴田良弘 (早大理工) Global well-posedness for the two component flow 9
Yoshihiro Shibata (Waseda Univ.) Global well-posedness for the two component flow

概要 In this talk, I would like to talk about a global in time unique existence theorem for the reduced system of equations from the two component flow. A key is to prove the exponential decay estimate in some quotient space of the linearized equations. Some special structure of equations guarantees this exponential decay property.

- 85 岡田 晃 (京大人間環境) Navier–Stokes 方程式の強解の正則性と漸近挙動 12
小 藺 英雄 (早大理工)
清 水 扇 丈 (京大人間環境)
Akira Okada (Kyoto Univ.) C^∞ regularity of strong solutions to the Navier–Stokes equations and
Hideo Kozono (Waseda Univ.) its decay property
Senjo Shimizu (Kyoto Univ.)

概要 Global mild solution u to the Navier–Stokes equation with the small initial data $u(0) \in L_n(\mathbb{R}^n)$ is constructed by Kato. We show u becomes infinitely differentiable with respect to space having the decay property $\|A^m u(t)\|_{L_p} = O(t^{-\frac{n}{2}(\frac{1}{n}-\frac{1}{p})-m})$ as $t \rightarrow \infty$ for all $n \leq p < \infty$.

- 86 金子健太 (早大理工) スケール不変 Besov 空間における Navier–Stokes 方程式の定常解について 12
小 藺 英雄 (早大理工)
清 水 扇 丈 (京大人間環境)
Kenta Kaneko (Waseda Univ.) Stationary solution to the Navier–Stokes equations in the scaling invariant Besov space
Hideo Kozono (Waseda Univ.)
Senjo Shimizu (Kyoto Univ.)

概要 We consider the stationary problem of the Navier–Stokes equations in \mathbb{R}^n for $n \geq 3$. We show existence, uniqueness and regularity of solutions in the homogeneous Besov space $\dot{B}_{p,q}^{-1+\frac{n}{p}}$ which is the scaling invariant one. As a corollary of our results, a self-similar solution is obtained. For the proof, several bilinear estimates are established. The essential tool is based on the paraproduct formula and the imbedding theorem in homogeneous Besov spaces.

16:30~17:30 特別講演

高田 了 (九大数理) Dispersive estimates for rotating fluids and stably stratified fluids
Ryo Takada (Kyushu Univ.) Dispersive estimates for rotating fluids and stably stratified fluids

概要 In this talk, we consider the initial value problems for the rotating Navier–Stokes equations and the stably stratified Boussinesq equations. We establish the sharp dispersive estimates for the linear propagators related to the rotation and the stable stratification. As applications, we give explicit relations between the size of initial data and the angular/buoyancy frequency which ensure the unique existence of global solutions to the above systems. Consequently, it is shown that the size of initial data can be taken large in proportion to the speed of rotation and the strength of stable stratification.

実函数論

9月13日(水) 第VII会場

9:30~12:00

- 1 原 宇信 (北大理・首都大東京理工) 逆ヘルダー型不等式の補間の性質とその Harnack の不等式への応用について 15
 Takano Hara (Hokkaido Univ./Tokyo Metro. Univ.) Interpolation properties of reverse Hölder inequalities and their applications to Harnack inequalities

概要 In this talk, we consider a backward self-improvement property of the reverse Hölder inequality with increasing support

$$\left(\frac{1}{\mu(B)} \int_B u^s d\mu \right)^{1/s} \leq C \left(\frac{1}{\mu(2B)} \int_{2B} u^p d\mu \right)^{1/p},$$

where $B = B(x_0, R)$ and $2B = B(x_0, 2R)$ are balls in a metric measure space (X, d, μ) , u is a nonnegative function on $2B$ and $0 < p < s \leq \infty$ and C are positive constants.

- 2 本田 あおい (九工大情報工) 可積分単関数の生成する L_0 の線形部分空間 15
 岡崎 悦明 (フuzzyシステム研)
 Aoi Honda (Kyushu Inst. of Tech.) Linear subspace of L_0 generated by integrable step functions
 Yoshiaki Okazaki (Fuzzy Logic Systems Inst.)

概要 Let $(\Omega, \mathcal{A}, \mu), \mu(\Omega) = +\infty$, be an infinite measure space and $L_0 = L_0(\Omega, \mathcal{A}, \mu)$ be the space of all real valued measurable functions on $(\Omega, \mathcal{A}, \mu)$. We introduce the closed linear subspace M_0 of L_0 generated by the integrable step functions. We give a characterization of M_0 .

- 3 河邊 淳 (信州大工) Choquet 積分に対する Vitali の収束定理 15
 Jun Kawabe (Shinshu Univ.) The Vitali convergence theorem for Choquet integrals

概要 The Vitali theorem for uniformly integrable functions is fundamental in Lebesgue integration theory and contains other important convergence theorems for abstract Lebesgue integral. The purpose of this talk is to formulate Vitali type theorems for the Choquet integral and its symmetric and asymmetric extensions with respect to a nonadditive measure. The bounded convergence theorem and the dominated convergence theorem for Choquet integrals are obtained as corollaries to our Vitali type theorems.

- 4 高阪 史明 (東海大理) 上に有界な曲率を持つ完備測地的距離空間における spherically nonspreading 写像の不動点 15
 Fumiaki Kohsaka (Tokai Univ.) Fixed points of spherically nonspreading mappings in complete geodesic metric spaces with curvature bounded above

概要 Fixed points of spherically nonspreading mappings in complete geodesic metric spaces with curvature bounded above.

In this talk, we propose the concepts of spherically nonspreading mappings and firmly spherically nonspreading mappings in complete CAT(1) spaces and obtain fixed point and convergence theorems for them. The resolvent of a proper lower semicontinuous convex function is a typical example of these mappings. As applications, we study the problem of minimizing convex functions in such spaces.

- 5 松下 慎也 (秋田県立大) 近接点法の収束について 15
 Shin-ya Matsushita (Akita Pref. Univ.) On the convergence of the proximal point algorithm

概要 Throughout this talk, let H be a Hilbert space and let $f : H \rightarrow (-\infty, \infty]$ be a proper lower semicontinuous convex function. We assume that f is bounded below. The proximal point algorithm is an approximation method for finding a minimizer of f . In this talk, we consider the convergence rate of the proximal point algorithm.

- 6 田村 高幸 (千葉大社会科学) On uniform non-squareness of direct sums of Banach spaces 15
 加藤 幹雄 (九工大*)
 Takayuki Tamura (Chiba Univ.) On uniform non-squareness of direct sums of Banach spaces
 Mikio Kato (Kyushu Inst. of Tech.*)

概要 We shall characterize the uniform non-squareness of the ψ -direct sum $(X_1 \oplus \cdots \oplus X_N)_\psi$ of Banach spaces X_1, \dots, X_N , where ψ is a convex function on the N -simplex Δ_N satisfying certain conditions. To do this we shall introduce a new class of convex functions.

- 7 鈴木 智成 (九工大工) τ -distance の再定義 15
 Tomonari Suzuki Redefinition of τ -distance
 (Kyushu Inst. of Tech.)

概要 We will talk about the redefinition of τ -distance.

- 8 厚 芝 幸子 (山梨大教育人間) Attractive point theorems and convergence theorems for some classes of
 nonlinear mappings in Hilbert spaces 15
 Sachiko Atsushiba Attractive point theorems and convergence theorems for some classes of
 (Univ. of Yamanashi) nonlinear mappings in Hilbert spaces

概要 In this talk, we prove attractive point theorems for nonlinear mappings. Using the ideas of attractive points and acute points, we also prove weak and strong convergence theorems for nonlinear mappings by some iterative methods.

- 9 渡辺 俊一 (東京情報大総合情報) Fixed point theorems in ordered metric spaces and applications to non-
 linear boundary value problems 15
 Toshikazu Watanabe Fixed point theorems in ordered metric spaces and applications to non-
 (Tokyo Univ. of Information Sci.) linear boundary value problems

概要 In this talk, we extend the concept of mixed monotone mappings and then we consider certain fixed point theorems for a pair of mappings in metric spaces with a partial ordering. As an application, we study existence of solutions for the following fourth-order two-point boundary value problems for elastic beam equations.

14:15~15:45

- 10 新井 龍太郎 (茨城大理) Commutators of Calderón–Zygmund and generalized fractional integral
 中井 英一 (茨城大理) operators on generalized Morrey spaces with variable growth condition
 15
 Ryutaro Arai (Ibaraki Univ.) Commutators of Calderón–Zygmund and generalized fractional integral
 Eiichi Nakai (Ibaraki Univ.) operators on generalized Morrey spaces with variable growth condition

概要 We discuss the boundedness of the commutators $[b, T]$ and $[b, I_\rho]$ on generalized Morrey spaces with variable growth condition, where T is a Calderón–Zygmund operator, I_ρ is a generalized fractional integral operator and b is a function in generalized Campanato spaces with variable growth condition.

- 11 竹本 奈央 (奈良女大人間文化) ウェーブレットの逆変換公式について 15
森 藤 紳 哉 (奈良女大理)
 Nao Takemoto (Nara Women's Univ.) Some variations on wavelet reconstruction formulae
 Shinya Moritoh (Nara Women's Univ.)

概要 We consider some variations on wavelet reconstruction formulae. An alternative formula was considered by Lebedeva and Postnikov in 2014. One of the aims of the talk is to give a multidimensional version of their formula.

- 12 飯田 毅 士 (福島工高専) On sufficient conditions for the boundedness of the fractional maximal operator between weighted L^p -spaces with different weights 15
 Takeshi Iida
 (Fukushima Nat. Coll. of Tech.) On sufficient conditions for the boundedness of the fractional maximal operator between weighted L^p -spaces with different weights

概要 In this talk, we discuss the boundedness of the fractional maximal operator between weighted L^p -spaces with different weights. The B_p -condition which is introduced by Pérez is also necessary and sufficient condition for the boundedness of the fractional maximal operator. As application of this theorem is related to the classical weighted inequality of Fefferman–Stein.

- 13 齋藤 洋樹 (日大理工) Maximal operators with the weighted Hausdorff content 15
田中 仁 (筑波技術大)
渡辺 俊一 (東京情報大総合情報)
 Hiroki Saito (Nihon Univ.) Maximal operators with the weighted Hausdorff content
 Hitoshi Tanaka
 (Tsukuba Univ. of Tech.)
 Toshikazu Watanabe
 (Tokyo Univ. of Information Sci.)

概要 In this talk, we first introduce the d -dimensional weighted Hausdorff content with arbitrary weight on \mathbb{R}^n . Then we establish the Fefferman–Stein type inequalities for the fractional maximal operator with the weighted Hausdorff content. Further, we discuss the boundedness of the fractional maximal operator on Choquet–Lorentz spaces.

- 14 宮地 晶彦 (東京女大現代教養)^b Exotic class の双線形擬微分作用素の有界性について 15
富田 直人 (阪大理)
 Akihiko Miyachi
 (Tokyo Woman's Christian Univ.) Boundedness of bilinear pseudo-differential operators with exotic symbols
 Naohito Tomita (Osaka Univ.)

概要 We give the sharp boundedness result for bilinear pseudo-differential operators in $L^p \times L^q$ to L^r , $1/p + 1/q = 1/r \leq 1$, in the case that the symbols satisfy the Hörmander condition with $0 \leq \rho < 1$.

16:00~17:00 特別講演

- 中村 昭宏 (東海大海洋) Nonharmonic Fourier 級数と Riesz 基底
 Akihiro Nakamura (Tokai Univ.) Nonharmonic Fourier series and Riesz bases

概要 The study of nonharmonic Fourier series was initiated by Paley and Wiener (1934). We take up the problems of the stability of Riesz basis properties of complex exponential systems in $L^2[-\pi, \pi]$. We consider the sequence $\{\lambda_n\}$ with perturbations of some subsequence of integers and investigate whether the system $\{e^{i\lambda_n t}\}$ becomes a Riesz basis in $L^2[-\pi, \pi]$. We use some stability theorems and the criterion obtained by B. S. Pavlov for $\{e^{i\lambda_n t}\}$ to become a Riesz basis. Finally, we refer to the existences of the bases which are not Riesz bases.

9月14日(木) 第VII会場

9:30~12:00

- 15 J. Lankeit (Paderborn Univ.) How far does small chemotactic interaction perturb the Fisher–KPP
 水上雅昭 (東京理大理) dynamics on bounded convex domains? 15
 Johannes Lankeit (Paderborn Univ.) How far does small chemotactic interaction perturb the Fisher–KPP
 Masaaki Mizukami dynamics on bounded convex domains?
 (Tokyo Univ. of Sci.)

概要 This work is concerned with the question that “how far does small chemotactic interaction perturb the Fisher–KPP dynamics?”. A chemotaxis system with logistic source was studied by e.g., Winkler (2010, 2014) and Zheng (2016). However, there are still many open problems about the chemotaxis system. On the other hand, the Fisher–KPP system has been studied extensively. Thus the development of this work will enable us to see new properties of solutions for the chemotaxis system. The main result of this talk gives convergence of solutions for the chemotaxis system to solutions for the Fisher–KPP system on bounded convex domains.

- 16 T. Black (Paderborn Univ.) Global existence in a Keller–Segel–(Navier–)Stokes system with singular
 J. Lankeit (Paderborn Univ.) sensitivity 15
 水上雅昭 (東京理大理)
 Tobias Black (Paderborn Univ.) Global existence in a Keller–Segel–(Navier–)Stokes system with singular
 Johannes Lankeit (Paderborn Univ.) sensitivity
 Masaaki Mizukami
 (Tokyo Univ. of Sci.)

概要 This talk is concerned with global existence of solutions to a Keller–Segel–(Navier–)Stokes system with singular sensitivity. In the fluid-free case, Winkler established global existence of classical solutions under some condition in 2011, and Fujie showed that the global solutions are bounded in 2015. However, a Keller–Segel system with singular sensitivity coupled with a Navier–Stokes equation has not been studied. The main result of this talk gives that the same condition assumed in Winkler’s result (2011) leads to global existence in the system.

- 17 元田大志 (京都教育大) 力学的境界条件下での退化放物型方程式への抽象発展方程式論からの接近
 深尾武史 (京都教育大) 15
 Taishi Motoda (Kyoto Univ. of Edu.) Abstract approach to degenerate parabolic equations with dynamic bound-
 Takeshi Fukao (Kyoto Univ. of Edu.) ary conditions

概要 The existence and uniqueness problem for a degenerate parabolic equation with dynamic boundary condition is discussed. Follows from the previous works by A. Damlamian (1977), the abstract theory of evolution equation, governed by the subdifferential of proper, lower-semicontinuous and convex functional, can be applied. Then, the suitable setting of function spaces and duality mapping is needed. One of the key point is the assumption of growth condition for the maximal monotone graph which characterizes the original degenerate diffusion.

- 18 加納理成 (高知大教育) 硬化現象を記述する1次元モデルの可解性について 15
深尾武史 (京都教育大)
Risei Kano (Kochi Univ.) The existence of solutions for the one-dimensional hardening model
Takeshi Fukao (Kyoto Univ. of Edu.)

概要 In this talk, we treat some parabolic problem with related to the quasi-variational inequality. The unknown functions $u = u(t, x)$ and $\sigma = \sigma(t, x)$ describe the displacement and stress, respectively in the one-dimensional interval $(0, L)$. The system stands for the hardening problem that the materials are hardened by plasticity. That is derived from the perfect plasticity model introduced by Duvaut–Lions. In the perfect plasticity model, the function which stands for threshold value in the plastic deformation is a constant. In this talk, we discuss the solvability for the above model under the situation that threshold function depending upon time or unknown function.

- 19 深尾武史 (京都教育大) Cahn–Hilliard approach to nonlinear diffusion equations on unbounded
来間俊介 (東京理大理) domains 15
横田智巳 (東京理大理)
Takeshi Fukao (Kyoto Univ. of Edu.) Cahn–Hilliard approach to nonlinear diffusion equations on unbounded
Shunsuke Kurima (Tokyo Univ. of Sci.) domains
Tomomi Yokota (Tokyo Univ. of Sci.)

概要 This talk deals with nonlinear diffusion equations under Neumann boundary conditions in a unbounded domain with smooth bounded boundary. Recently, Kurima–Yokota (2017) proved existence of solutions to these equations with growth conditions for diffusion terms. The present work asserts that we can solve the original problem by passing to the limit in the approximate problem without growth conditions in a unbounded domain.

- 20 深尾武史 (京都教育大) 領域内部の方程式を補助条件とする境界上での Cahn–Hilliard 方程式の
P. Colli (Pavia Univ.) 可解性について 15
Takeshi Fukao (Kyoto Univ. of Edu.) Cahn–Hilliard equation on the boundary with bulk condition
Pierluigi Colli (Pavia Univ.)

概要 The well-posedness for a system of partial differential equations and dynamic boundary conditions is discussed. This system is a sort of transmission problem between the dynamics in the bulk Ω and on the boundary Γ . The Poisson equation for the chemical potential, the Allen–Cahn equation for the order parameter in the bulk Ω are considered as auxiliary conditions for solving the Cahn–Hilliard equation on the boundary Γ .

- 21 山崎教昭 (神奈川大工) New class of doubly nonlinear evolution equations governed by double
剣持信幸 (ワルシャワ大ICM) time-dependent subdifferentials 15
白川健 (千葉大教育)
Noriaki Yamazaki (Kanagawa Univ.) New class of doubly nonlinear evolution equations governed by double
Nobuyuki Kenmochi (Univ. of Warsaw) time-dependent subdifferentials
Ken Shirakawa (Chiba Univ.)

概要 We discuss a new class of doubly nonlinear evolution equations governed by time-dependent subdifferentials in uniformly convex Banach spaces, and establish an abstract existence result of solutions. Also, we give some applications to nonlinear PDEs with gradient constraint for time-derivatives.

- 22 中屋敷亮太 (千葉大理) 力学的境界条件を課す場合の Allen–Cahn 型特異拡散方程式に対する数
白川 健 (千葉大教育) 学解析 15
Ryota Nakayashiki (Chiba Univ.) Qualitative properties of the solution to Allen–Cahn type equations with
Ken Shirakawa (Chiba Univ.) singularities subject to dynamic boundary condition

概要 In this talk, we consider coupled system of nonlinear PDEs. The system consists of an Allen–Cahn type equation with singular diffusion in a bounded spatial domain Ω , and another Allen–Cahn type equation on the smooth boundary $\partial\Omega$. The coupled PDEs are transmitted via the dynamic boundary condition. The objective of this study is to achieve a mathematical treatment to analyze the systems for singular diffusion equations and the dynamic boundary conditions. Now, the results concerned with the well-posedness of the system, involved in the representation of solution and comparison principal and the continuous association between solutions to our system and those in regular systems, are reported in forms of some Main Theorems.

- 23 白川 健 (千葉大教育) 結晶粒界運動の数学モデルの 1 次元定常解の解構造 15
渡邊 紘 (大分大理工)
中屋敷亮太 (千葉大理)
S. Moll (Univ. Valencia)
Ken Shirakawa (Chiba Univ.) Structures of steady-state solutions to a one-dimensional mathematical
Hiroshi Watanabe (Oita Univ.) model of grain boundary motion
Ryota Nakayashiki (Chiba Univ.)
Salvador Moll (Univ. Valencia)

概要 In this talk, we consider a system of one-dimensional elliptic type boundary value problems, denoted by (S_∞) . The system corresponds to a one-dimensional steady-state problem for the mathematical model of grain boundary motion, proposed by [Kobayashi et al, Phys. D, 140 (2000), 141–150], and one of characteristics is in the point that the inhomogeneous Dirichlet type boundary condition is imposed for the crystalline orientation. On this basis, we set the objectives of the talk as follows: (A) to show the structures of all steady-state solutions, including physically-important ones; (B) to clarify the base-structure of steady-states with the physical importance; (C) the verification of the SBV-regularity for the steady-state solutions.

14:15~16:00

- 24 愛木 豊彦 (日本女大理) コンクリート中性化過程に現れる 1 次元水分輸送方程式に対する制御問題
S. A. Timoshin 15
(Siberian Branch Russian Acad. Sci.)
Toyohiko Aiki (Japan Women's Univ.) Control problem for the one-dimensional moisture transport equation
Sergey A. Timoshin appearing in concrete carbonation process
(Siberian Branch Russian Acad. Sci.)

概要 We consider a system of partial differential equations describing a mass conservation law for moisture in a porous medium. This type of systems can be found in concrete carbonation process and already proposed and studied by Kumazaki–Aiki. In the system the relationship between the relative humidity and the degree of saturation is described by a play operator. In this talk we consider a real time control problem for the above system. The aim of the problem is to control a solution of the system by putting a multi-valued operator into the differential equation. Here, I will discuss about a physical background for the control and establish the existence of a solution to the control problem.

- 25 喜多航佑 (早大理工) On some elliptic systems arising from a nuclear reactor model 15
大谷光春 (早大理工)
坂本浩紀
 (日立GEニュークリア・エナジー(株))
Kosuke Kita (Waseda Univ.) On some elliptic systems arising from a nuclear reactor model
Mitsuharu Ôtani (Waseda Univ.)
Hiroki Sakamoto
 (Hitachi-GE Nuclear Energy, Ltd.)

概要 We consider a stationary problem of a certain reaction diffusion system arising from a nuclear reactor model, which consists of two unknown functions representing the neutron density and the temperature in nuclear reactors. In Gu–Wang (1994, 1996), they studied this problem with some boundary conditions (homogeneous Dirichlet–Dirichlet conditions and homogeneous Neumann–Robin conditions) and prove the solvability and the uniqueness of ordered positive solution. In this talk, we impose Robin and power type nonlinear boundary conditions on the problem and show the existence and the uniqueness results similar to the previous results. We rely on Krasnoselskii’s type fixed point theorem due to lack of the variational structure.

- 26 黒田隆徳 (早大理工) Solvability of complex Ginzburg–Landau equations with non-dissipative
大谷光春 (早大理工) terms in general domains 15
Takanori Kuroda (Waseda Univ.) Solvability of complex Ginzburg–Landau equations with non-dissipative
Mitsuharu Ôtani (Waseda Univ.) terms in general domains

概要 We consider the following complex Ginzburg–Landau equation, (CGL)₋:

$$u_t(t, x) - (\lambda + i\alpha)\Delta u - (\kappa + i\beta)|u|^{q-2}u - \gamma u = f(t, x) \quad \text{on } [0, T) \times \Omega,$$

where $\lambda, \kappa > 0$; $\alpha, \beta, \gamma \in \mathbb{R}$; i denotes the imaginary unit; $T > 0$; $f : [0, T) \times \Omega \rightarrow \mathbb{C}$ is a given external force and Ω is general, possibly unbounded domain. Our approach to (CGL)₋ is to regard our equation as a parabolic equation in a product Hilbert space $(L^2(\Omega))^2$ over \mathbb{R} with $-\lambda\Delta u$ being a principal term, and when $\kappa > 0$, our nonlinear term should be treated as a non-monotone perturbation. For a general domain Ω , it is difficult to handle such kind of perturbations, because of the lack of compactness.

- 27 渡邊 紘 (大分大理工) ある退化放物型-楕円型方程式系の可解性 15
Hiroshi Watanabe (Oita Univ.) Solvability of degenerate parabolic-elliptic systems

概要 We consider the initial value problem (CP) for degenerate parabolic-elliptic systems with variable coefficients. The systems are coupled with strongly degenerate parabolic equations and elliptic equations. Strongly degenerate parabolic equations are regarded as a linear combination of the time-dependent conservation laws (quasilinear hyperbolic equations) and the porous medium type equations (nonlinear degenerate parabolic equations). Thus, this equation has both properties of hyperbolic equations and those of parabolic equations. In this talk, we formulate entropy solutions to (CP) and show the existence and uniqueness of the solutions.

- 28 熊崎 耕太 (苫小牧工高専) 多孔質媒体内での水分膨張過程を記述するある自由境界問題について .. 15
Kota Kumazaki On a free boundary problem for moisture swelling process in porous
 (Tomakomai Nat. Coll. of Tech.) materials

概要 In this talk, we propose a mathematical model for moisture swelling process in concrete materials. Moisture swelling process appear in, for instance, frost damage in concrete materials which is a nonlinear phenomenon to give rise to crack inside of concrete. Our model consists of a diffusion equation for moisture in a one microscopic hole of concrete and a free boundary problem for the front of moisture region. In this talk, we discuss the existence and uniqueness of a solution of this problem.

- 29 都 築 寛 (広島修道大経済) Initial-boundary value problems for Vlasov–Poisson systems with angle error in magnetic field 15
- Yutaka Tsuzuki Initial-boundary value problems for Vlasov–Poisson systems with angle error in magnetic field
(Hiroshima Shudo Univ.)

概要 In this talk, we deal with existence of solutions to Vlasov–Poisson systems in a half-space with external magnetic force horizontal to a wall. In 2013, Skubachevskii gives local-in-time solvability to the system. Moreover, in 2017, global-in-time solutions were obtained by effectively using the magnetic force whose direction is horizontal to the wall. This talk provides an existence result for the system where the magnetic force has angle error in the vertical direction.

16:15～17:15 特別講演

- 都 築 寛 (広島修道大経済) Global existence of solutions to Vlasov–Poisson equations with external magnetic field in a half space
- Yutaka Tsuzuki Global existence of solutions to Vlasov–Poisson equations with external magnetic field in a half space
(Hiroshima Shudo Univ.)

概要 This talk is concerned with solvability of Vlasov–Poisson equations in a half-space with external magnetic field. In 2013, local-in-time existence of solutions to the equation was proved by Skubachevskii. However, the result shows the fact that the existence time T is exponentially small, which means the plasma can reach a wall and melt it in the extremely short time. The purpose of this talk is to obtain existence result for the equation with a very large time T . Moreover this talk provides global-in-time solvability for the equation with a more strict condition for the magnetic force whose direction is horizontal to the wall.

函数解析学

9月11日(月) 第VI会場

14:15~16:45

- 1 岩田 順 敬 (東工大科学技術創成研究院) 無限小生成作用素の対数表現に関する非線形性 15

Yoritaka Iwata (Tokyo Tech) Nonlinearity as for the logarithmic representation of infinitesimal generators

概要 The logarithmic representation of invertible evolution families is introduced in Ref.[1]. The nonlinearity related to the logarithmic representation is discussed in terms of its similarity with the Coles–Hopf transform.

Ref: [1] Yoritaka Iwata, “Infinitesimal generators of invertible evolution families”, *Methods Funct. Anal. Topology* **23** 1 (2017) 26–36.

- 2 和田 和 幸 (八戸工高専)^b N -体シュタルクハミルトニアンのスペクトル 15

Kazuyuki Wada (Nat. Inst. of Tech., Hachinohe Coll.) Spectrum of N -body Stark Hamiltonians

概要 We consider N -body Stark Hamiltonians. By Herbst et al, it is shown that both pure point spectrum and singular continuous spectrum of N -body Stark Hamiltonians are empty. Their method is based on Mourre’s positive commutator method. We prove this result by applying the theory of generalized canonical commutation relations.

- 3 蘆田 聡 平 (京 大 理) Absence of singular continuous spectrum and propagation estimates of multistate Schrödinger operators 15

Sohei Ashida (Kyoto Univ.) Absence of singular continuous spectrum and propagation estimates of multistate Schrödinger operators

概要 We study multistate Schrödinger operators related to molecular dynamics. We consider potentials which do not necessarily decay including those homogeneous of degree zero. We prove absence of the singular continuous spectrum and propagation estimates which mean the scattering at speed larger than a positive constant and decay of the states with potentials higher than considered energy at infinity. We also consider the multistate Schrödinger operators with many-body structures. We obtain the Mourre estimate and the minimal velocity estimate for the many-body operators. The lower bound of the velocity is determined by the distance between the energy and thresholds below the energy.

- 4 森 岡 悠 (同志社大理工) Interior transmission eigenvalue problems on compact manifolds with smooth boundary 15

Hisashi Morioka (Doshisha Univ.) Interior transmission eigenvalue problems on compact manifolds with smooth boundary
Naotaka Shoji (Univ. of Tsukuba)

概要 We consider an interior transmission eigenvalue (ITE) problem on some compact C^∞ -Riemannian manifolds with smooth boundary. In particular, we do not assume that two domains are diffeomorphic, but we impose some conditions of Riemannian metrics and indices of refraction on the boundary. Then we prove the discreteness of the set of ITEs, the existence of infinitely many ITEs, and its Weyl type lower bound.

- 5 川本昌紀 (東京理大理) 時間減衰する調和ポテンシャルをもつ Schrödinger 方程式の解に対する
 米山泰祐 (東京理大理) Strichartz 型評価 15
 Masaki Kawamoto (Tokyo Univ. of Sci.) Strichartz estimates for harmonic potential with time-decaying coefficient
 Taisuke Yoneyama (Tokyo Univ. of Sci.)

概要 In this talk, we prove the Strichartz estimates for the Schrödinger equations with a harmonic potential with a time-decaying coefficient by introducing the time weighted Lebesgue space.

- 6 新國裕昭 (前橋工科大) 多重結合からなるジグザグナノチューブ上の周期的シュレディンガー作用素のスペクトルについて 15
 Hiroaki Niikuni Spectra of periodic Schrödinger operators on zigzag nanotubes with multiple chemical bonds
 (Maebashi Inst. of Tech.)

概要 In this talk, we study the spectral structure of periodic Schrödinger operators on zigzag nanotubes with multiple chemical bonds. Utilizing the Floquet–Bloch theory for the corresponding quantum graph, we see that the spectrum has the band-gap structure. Namely, the spectrum consists of flat band and the absolutely continuous spectrum. In the talk, we see the difference between the case of the single bond and the multiple bonds.

- 7 山岸弘幸 (産業技術高専) 弾性基盤上の張力をかけた棒のたわみの 2 点境界値問題と対応するソボレフ不等式の最良定数 3 15
 Hiroyuki Yamagishi The best constant of Sobolev inequality corresponding to a bending problem of a beam under tension on an elastic foundation 3
 (Tokyo Metropolitan Coll. of Indus. Tech.)

概要 We consider two-point boundary value problems for bending of a beam supported by uniformly distributed springs with spring constant $q > 0$ on a fixed floor under tension $p > 0$. The tension is relatively strong, that is $(p/2)^2 > q$. We have treated periodic, Dirichlet, Dirichlet–Neumann and Neumann boundary conditions and found their Green functions. As an application, we have found the best constants of the corresponding Sobolev inequality, which are equal to the maximum of diagonal values of Green functions.

- 8 伊藤宏 (愛媛大理工)* 発散するポテンシャルをもつパウリ作用素の平方根について 15
 Hiroshi Ito (Ehime Univ.) On the square root of the Pauli operator with a diverging potential

概要 We first study the spectral properties of two relativistic Hamiltonians; one is the square root of a Pauli operator with an electric potential growing polynomially at infinity, and the other differs from it only in the sign of the potential. We next give a resonance free region for the latter. Moreover, we show that resonances (eigenvalues) of each of them converge to resonances (eigenvalues) of the corresponding Pauli operators with the same potential in the nonrelativistic limit.

- 9 伊藤宏 (愛媛大理工)* 発散するポテンシャルをもつ Dirac 作用素のレゾナンス 10
 Hiroshi Ito (Ehime Univ.) Resonances of Dirac operators with a diverging potential

概要 We investigate resonance free regions of Dirac operators with a bounded magnetic potential and an electric potential diverging at infinity with the help of the dilation analytic method and the FW transform. In this work two square roots of c -dependent Pauli operators play an important role.

17:00~18:00 特別講演

川本昌紀 (東京理大理) 時間周期パルス磁場に対する散乱理論

Masaki Kawamoto (Tokyo Univ. of Sci.) Scattering theory for periodically pulsed magnetic field

概要 We study the quantum dynamics of a charged particle in the plane in the presence of a periodically pulsed magnetic field perpendicular to the plane. We show that by controlling the cycle when the magnetic field is switched on and off appropriately, the result of the asymptotic completeness of wave operators can be obtained under the assumption that the potential V satisfies the decaying condition $|V(x)| \leq C(1 + |x|)^{-\rho}$, for some $\rho > 0$.

9月12日(火) 第VI会場

9:45~12:00

10 富樫瑠美 (長岡工高専) 末梢スペクトルと積を用いた写像の特徴づけ について 15

Rumi Togashi Characterization of maps by peripheral spectra and multiplication

(Nagaoka Nat. Coll. of Tech.)

概要 We introduce the condition related to the peripheral spectrum and the multiplication for maps between uniform algebras to be linear and isometric. We can describe maps related to algebra isomorphisms by some property of peripheral spectra and multiplication. We also generalize weakly peripherally-multiplicative maps and peripherally monomial-preserving maps and give some examples.

11 三浦 毅 (新潟大理) $C^1([0, 1])$ 上の等距離写像 15

高木 啓行 (信州大理)

Takeshi Miura (Niigata Univ.) Isometries on $C^1([0, 1])$

Hiroyuki Takagi (Shinshu Univ.)

概要 Let $C^1([0, 1])$ be a linear space of all continuously differentiable complex valued functions on the closed unit interval $[0, 1]$. $C^1([0, 1])$ is a Banach space with respect to the following norms: $\|f\|_C = \sup_{t \in [0, 1]} (|f(t)| + |f'(t)|)$, $\|f\|_\Sigma = \|f\|_\infty + \|f'\|_\infty$ and $\|f\|_\sigma = |f(0)| + \|f'\|_\infty$, where $\|\cdot\|_\infty$ denotes the supremum norm on $[0, 1]$. We give the characterization of surjective isometries on $C^1([0, 1])$ with respect to the above norms.

12 渡辺 恵一 (新潟大理) A. A. Ungar のメビウスジャイロベクトル空間について 15

Keiichi Watanabe (Niigata Univ.) On the Möbius gyrovector spaces by A. A. Ungar

概要 For the Möbius gyrovector spaces introduced by A. A. Ungar, we reveal the structure of finitely generated gyrovector subspaces, present a notion of orthogonal gyrodecomposition with respect to any gyrovector subspace which is closed under the Poincare metric. Moreover, we show a concrete procedure to obtain orthogonal gyroexpansion in a Möbius gyrovector space, like as the classical orthogonal expansion in a Hilbert space.

13 羽鳥 理 (新潟大自然) Peculiar homomorphisms on admissible quadruples 15

Osamu Hatori (Niigata Univ.) Peculiar homomorphisms on admissible quadruples

概要 We study unital homomorphisms on admissible quadruples. In particular, we exhibit results that every unital homomorphism between admissible quadruples with certain conditions on maximal ideal spaces is of type BJ.

- 14 田端 亮 (有明工高専) Schur の immanant 不等式と Littlewood–Richardson 対応 …………… 15
 Ryo Tabata (Ariake Nat. Coll. of Tech.) Schur’s immanant inequalities and Littlewood–Richardson’s correspondence

概要 Immanants are generalizations of the determinant and the permanent, and are labeled by Young diagrams. The limit of immanants of a correlation matrix is an interesting problem in terms of inequality problems, whose origins are Schur’s inequality and Lieb’s permanental dominance conjecture. In this talk, we give some results of the limit of immanants depending on the arms and legs of the Young diagrams, applying the Littlewood–Richardson rule, which is one of the most important property to describe the representations of the symmetric group. Also, we observe the behavior of the Littlewood–Richardson rule that becomes simple under some conditions.

- 15 C. Reyes-Bustos (九大数理) Spectral degeneracies in the asymmetric quantum Rabi model …………… 15
 木本一史 (琉球大理)
 若山正人 (九大数理)
 Cid Reyes-Bustos (Kyushu Univ.) Spectral degeneracies in the asymmetric quantum Rabi model
 Kazufumi Kimoto (Univ. of Ryukyus)
 Masato Wakayama (Kyushu Univ.)

概要 In this talk, the authors prove the existence of spectral degeneracies in the asymmetric quantum Rabi model (AQRM) when the symmetry-breaking parameter ϵ is a half-integer. The degeneracy had been previously established for the case $\epsilon = \frac{1}{2}$ by Wakayama (2017) and verified experimentally the general case by Li and Batchelor in 2015. The main result is established by the study of certain (so-called constraint) polynomials appearing from the finite-dimensional irreducible representations of $\mathfrak{sl}_2(\mathbb{R})$ in the representation theoretical picture of the AQRM. Two independent proofs are given of the main result, each one giving a better understanding of the structure of the spectrum of the AQRM.

- 16 示野信一 (関西学院大理工) Small K -type に対する Harish-Chandra の c -関数 …………… 15
 織田 寛 (拓殖大工)
 Nobukazu Shimeno Harish-Chandra’s c -functions for small K -types
 (Kwansei Gakuin Univ.)
 Hiroshi Oda (Takushoku Univ.)

概要 We study elementary spherical functions on a non-compact real simple Lie group of finite center associated with a small K -type in the sense of Wallach. We prove that in most cases, the radial parts of elementary spherical functions for small K -types are written by hypergeometric functions of Heckman and Opdam. As an application, we give explicit formulae for Harish-Chandra’s c -functions for small K -types and obtain the inversion formulae for the spherical transforms for small K -types.

- 17 中濱良祐 (東大数理) Intertwining operators between holomorphic discrete series representations …………… 15
 Ryosuke Nakahama (Univ. of Tokyo) Intertwining operators between holomorphic discrete series representations

概要 In this talk the speaker presents the result on the explicit construction of embedding maps between two holomorphic discrete series representations. Today we mainly deal with the embedding of the holomorphic discrete series representation of $U(s', s'')$ into that of $Sp(s, \mathbb{R})$, where $s = s' + s''$.

13:00~14:00 特別講演

久保利久 (龍谷大経済) Differential symmetry breaking operators of $O(n, 1)$ for differential forms
 Toshihisa Kubo (Ryukoku Univ.) Differential symmetry breaking operators of $O(n, 1)$ for differential forms

概要 Let X be a smooth manifold and Y a smooth submanifold of X . Take $G' \subset G$ to be a pair of Lie groups that act transitively on $Y \subset X$, respectively. Suppose that $\mathcal{V} \rightarrow X$ and $\mathcal{W} \rightarrow Y$ are G - and G' -equivariant vector bundles over X and Y with fibers V and W , respectively. Then we call a differential operator $\mathcal{D}: C^\infty(X, \mathcal{V}) \rightarrow C^\infty(Y, \mathcal{W})$ between the spaces of smooth sections a *differential symmetry breaking operator* (differential SBO) if \mathcal{D} is G' -intertwining.

In the last year, for the setting $(G, G', V, W) = (O(n+1, 1), O(n, 1), \wedge^i(\mathbb{C}^n), \wedge^j(\mathbb{C}^{n-1}))$ with $n \geq 3$, we completely classified the differential SBOs with their explicit formulas. In other words, for any $0 \leq i \leq n$ and $0 \leq j \leq n-1$, we classified all the differential SBOs $\mathcal{D}^{i \rightarrow j}: \mathcal{E}^i(S^n) \rightarrow \mathcal{E}^j(S^{n-1})$ from the space of differential i -forms $\mathcal{E}^i(S^n)$ over the standard Riemann sphere S^n to that of differential j -forms $\mathcal{E}^j(S^{n-1})$ over the totally geodesic hypersphere S^{n-1} . In this talk we would like to discuss how we classify such operators. This is a joint work with T. Kobayashi and M. Pevzner.

9月13日(水) 第VI会場

9:00~12:00

- 18 藤本将行 (大阪教育大) Matrix Wielandt inequality via the matrix geometric mean 15
 瀬尾祐貴 (大阪教育大教育)

Masayuki Fujimoto Matrix Wielandt inequality via the matrix geometric mean
 (Osaka Kyoiku Univ.)
 Yuki Seo (Osaka Kyoiku Univ.)

概要 In this talk, by virtue of the matrix geomtric mean and the polar decomposition, we present new Wielandt type inequalities for matrices of any size. To this end, based on results due to J. I. Fujii, we reform a matrix Cauchy–Schwarz inequality, which differs from ones due to Marshall and Olkin.

- 19 藤井淳一 (大阪教育大教育) 負パラメータのツァリス相対作用素エントロピー 15
 瀬尾祐貴 (大阪教育大教育)

Junichi Fujii (Osaka Kyoiku Univ.) Tsallis relative operator entropy with negative parameters
 Yuki Seo (Osaka Kyoiku Univ.)

概要 Tsallis relative operator entropy was firstly formulated by Fujii and Kamei as an operator version of Uhlmann's relative entropy. Afterwards, Yanagi, Kuriyama and Furuichi reformulated Tsallis relative operator entropy as an operator version of Tsallis relative entropy. In this talk, we define Tsallis relative operator entropy with negative parameters of (non-invertible) positive operators on a Hilbert space and show some properties.

20 遠山宏明 (前橋工科大) The n -th divergence on the path $A \natural_t B$ 15
 伊佐浩史 (前橋工科大)
 伊藤公智 (前橋工科大)
 亀井栄三郎
 渡邊雅之 (前橋工科大)
 Hiroaki Tohyama The n -th divergence on the path $A \natural_t B$
 (Maebashi Inst. of Tech.)
 Hiroshi Isa (Maebashi Inst. of Tech.)
 Masatoshi Ito (Maebashi Inst. of Tech.)
 Eizaburou Kamei
 Masayuki Watanabe
 (Maebashi Inst. of Tech.)

概要 Let A and B be bounded positive invertible operators on a Hilbert space and let $\Psi_{A,B}(t) \equiv A \natural_t B$ be an operator valued smooth function, where $A \natural_t B \equiv A^{\frac{1}{2}}(A^{-\frac{1}{2}}BA^{-\frac{1}{2}})^t A^{\frac{1}{2}}$ ($t \in \mathbb{R}$) is a path passing through A and B . We consider the following functions $\Psi_{A,B}^{[n]} : \mathbb{R}^2 \rightarrow B(\mathcal{H})$: $\Psi_{A,B}^{[1]}(x, y) \equiv \frac{\Psi_{A,B}(x) - \Psi_{A,B}(y)}{x - y}$ and $\Psi_{A,B}^{[n]}(x, y) \equiv \frac{\Psi_{A,B}^{[n-1]}(x, y) - \Psi_{A,B}^{[n-1]}(y, y)}{x - y}$ ($n \geq 2$). Since Petz–Bregman divergence $D_{FK}(A|B)$ can be represented by $\Psi_{A,B}^{[1]}(1, 0) - \Psi_{A,B}^{[1]}(0, 0)$, we can give the n -th Petz–Bregman divergence $D_{FK}^{[n]}(A|B) \equiv \Psi_{A,B}^{[n]}(1, 0) - \Psi_{A,B}^{[n]}(0, 0)$. Moreover, we treat the n -th divergences related with some operator valued divergences defined by the difference between the relative operator entropies.

21 伊藤公智 (前橋工科大) Estimations of the Lehmer mean by the Heron mean 15
 Masatoshi Ito (Maebashi Inst. of Tech.) Estimations of the Lehmer mean by the Heron mean

概要 As generalizations of the arithmetic and geometric means for positive real numbers a and b , the power difference mean $J_q(a, b) = \frac{q}{q+1} \frac{a^{q+1} - b^{q+1}}{a^q - b^q}$, the Lehmer mean $L_q(a, b) = \frac{a^{q+1} + b^{q+1}}{a^q + b^q}$, and the Heron mean $K_q(a, b) = (1 - q)\sqrt{ab} + q \frac{a+b}{2}$ are well known. Recently, we have shown estimations of the power difference mean by the Heron mean.

In this talk, similarly to these results, we get estimations of the Lehmer mean by the Heron mean. In other words, we obtain the greatest value $\alpha = \alpha(q)$ and the least value $\beta = \beta(q)$ such that the double inequality $K_\alpha(a, b) < L_q(a, b) < K_\beta(a, b)$ holds for any $q \in \mathbb{R}$. We can also obtain operator inequalities for bounded linear operators on a Hilbert space.

22 山崎文明 (東洋大理工) Upper and lower bounds, and operator monotonicity of an extension of
 伊藤公智 (前橋工科大) the Petz–Hasegawa function 10
 古田孝之 (弘前大*)
 柳田昌宏 (東京理大理)
 Takeaki Yamazaki (Toyo Univ.) Upper and lower bounds, and operator monotonicity of an extension of
 Masatoshi Ito (Maebashi Inst. of Tech.) the Petz–Hasegawa function
 Takayuki Furuta (Hirosaki Univ.*)
 Masahiro Yanagida
 (Tokyo Univ. of Sci.)

概要 In this talk, we shall consider an extension of the Petz–Hasegawa function. In fact, we shall give upper and lower bounds, and operator monotonicity of this function with elementary proofs.

- 23 渚 勝 (千葉大理) 正定値関数と作用素ノルム不等式 15
 イمامヌグラハアルバニア
 (インドネシア教育大)

Masaru Nagisa (Chiba Univ.) Positive definite functions and operator norm inequalities
 Imam Nugraha Albania
 (Univ. Pendidikan Indonesia)

概要 Let $a_1 \geq a_2 \geq \dots \geq a_K > 0$ and $b_1 \geq b_2 \geq \dots \geq b_K > 0$. We consider the function

$$g(x) = \prod_{i=1}^K \frac{\sinh a_i x}{\sinh b_i x}.$$

It is known that the positive definiteness of such functions are related to some operator norm inequalities. In the case of $K = 2$, g is positive definite if and only if $a_1 \leq b_1$ and $a_1 + a_2 \leq b_1 + b_2$. Unfortunately we do not know such an equivalent condition when $K \geq 3$. We consider the condition which related to the positive definiteness of g .

- 24 大野修一 (日本工大)* The Toeplitzness of weighted composition operators 10
 Shūichi Ohno (Nippon Inst. of Tech.) The Toeplitzness of weighted composition operators

概要 We will consider the asymptotic toeplitzness associated with weighted composition operators on the Hardy–Hilbert space H^2 .

- 25 榎本雅俊 * 2つの部分空間の同配置問題 15
 綿谷安男 (九大数理)

Masatoshi Enomoto Configuration problem of two subspaces
 Yasuo Watatani (Kyushu Univ.)

概要 P. R. Halmos has presented his famous results of a classification of 2 subspaces under unitary equivalence. In this talk we discuss two subspaces under more weak equivalence. We point out that operator ranges are crucially important to study a classification of 2 subspaces under more weak equivalence. Under this equivalence, we give continuously many non-isomorphic examples of systems of 2 subspaces. We also give a relation between operator ranges, Hilbert representations of A_2 Dynkin quiver and particular systems of 3 subspaces.

- 26 梶原毅 (岡山大環境) シェルピンスキカーペットに付随する C^* -環の解析 15
 綿谷安男 (九大数理)

Tsuyoshi Kajiwara (Okayama Univ.) C^* -algebra associated with Sierpinski carpet
 Yasuo Watatani (Kyushu Univ.)

概要 In this talk, we present analysis of the core of the C^* -algebras associated with self-similar maps associated with Sierpinski carpet. Although, the branch set is a infinite set and the structure of branching is complex for Sierpinski carpet, we can classify finite traces and ideals of the core, and can describe the matrix representation of the core.

- 27 須藤隆洋 (琉球大理) The K-theory for the group and subgroup C^* -algebras of the special or general linear groups over integers 15

Takahiro Sudo (Univ. of Ryukyus) The K-theory for the group and subgroup C^* -algebras of the special or general linear groups over integers

概要 We consider the K-theory of the group and subgroup C^* -algebras of the special or general linear groups over the ring of integers and of their canonical subgroups. We further consider the K-theory of the associated, crossed product C^* -algebras.

- 28 松本健吾 (上越教育大)* 双曲型力学系における漸近的連続軌道同型と Ruelle C^* -環について 15
 Kengo Matsumoto Asymptotic continuous orbit equivalence of hyperbolic dynamics and
 (Joetsu Univ. of Edu.) Ruelle C^* -algebras

概要 A Smale space is a hyperbolic dynamical system with local product structure. D. Ruelle constructed C^* -algebras from Smale spaces. The algebras are regarded as higher dimensional analogues of Cuntz–Krieger algebras. We introduce notions of asymptotic continuous orbit equivalence and asymptotic conjugacy in Smale spaces and characterize them in terms of their étale groupoids and their asymptotic Ruelle algebras with their dual actions, respectively.

14:15~15:30

- 29 鈴木悠平 (名大多元数理) 極小拡大の接合積 C^* -環の純無限性について 15
 Yuhei Suzuki (Nagoya Univ.) On pure infiniteness of crossed products of minimal extensions

概要 We show that pure infiniteness of reduced crossed product is inherited to minimal extensions.

- 30 縄田紀夫 (大阪教育大教育) Trace scaling automorphisms of $\mathcal{W} \otimes \mathbb{K}$ 15
 Norio Nawata (Osaka Kyoiku Univ.) Trace scaling automorphisms of $\mathcal{W} \otimes \mathbb{K}$

概要 In this talk, we classify trace scaling automorphisms of $\mathcal{W} \otimes \mathbb{K}$ up to outer conjugacy, where \mathcal{W} is a certain simple separable nuclear stably projectionless C^* -algebra having trivial K -groups.

- 31 佐藤康彦 (京大理) UHF 環を吸収する C^* -環における自己同型の近似的ユニタリ同値 15
 Yasuhiko Sato (Kyoto Univ.) Projections associated with quasidiagonality

概要 In the recent breakthrough by Tikuisis, White, and Winter, it is shown that the universal coefficient theorem and nuclearity imply quasidiagonality for separable C^* -algebras. Precisely, they showed a faithful tracial state is quasi-diagonal under the natural assumption required in the classification theory. In this talk, we explain a technical ingredient in their proof and several alternative approaches to obtaining quasidiagonality.

- 32 澤田友佐 (名大多元数理) W^* -双加群の成す双圏 15
 山上 滋 (名大多元数理)
 Yusuke Sawada (Nagoya Univ.) The bicategory of W^* -bimodules
 Shigeru Yamagami (Nagoya Univ.)

概要 When a von Neumann algebra acts on a Hilbert space, the relative tensor product of the Hilbert spaces is defined. The notion of the relative tensor product was introduced by Alain Connes. There are two ways which we define it by changing left or right Hilbert space into the operator space. We call them the left and right relative tensor products respectively. We will show that the two categories consisting of all bimodules (i.e. Hilbert spaces on which von Neumann algebras act from left and right) with left and right relative tensor products are equivalent. This is a joint work with Shigeru Yamagami.

15:45~16:45 特別講演

磯野 優介 (京大数理研) Deformation/rigidity 理論と III 型フォンノイマン環

Yusuke Isono (Kyoto Univ.) Deformation/rigidity theory and type III von Neumann algebras

概要 Deformation/rigidity theory is initiated by S. Popa in 2001 to study *non-amenable* von Neumann algebras. Amenable von Neumann algebras naturally appear in physics and they were extensively studied. The study of non-amenable algebras also attracted attention but very few had been known until 1990s. Deformation/rigidity theory is a great success in the study of non-amenable algebras. It brought much progress and in fact solved lots of open problems for non-amenable algebras.

Most of technologies used in this new theory require so-called a *trace* (a generalization of traces on matrices). Since this requirement is crucial, the above development is mostly restricted to algebras with traces. Von Neumann algebras without any traces are called *type III*. For example, von Neumann algebras in physics are always of type III. They also appear in many other context such as ergodic theory, quantum groups, free probability theory etc.

In this talk, I focus on type III algebras, particularly non-amenable type III algebras. To study them, it is important to find a way of applying technologies of deformation/rigidity theory to type III algebras. This problem has been mainly studied by C. Houdayer and myself. I will survey recent progress on this problem.

統計数学

9月11日(月) 第I会場

9:15~12:00

- 1 小山 民雄 (神戸大理) 超函数による反転公式の表示とその応用 15
 Tamio Koyama (Kobe Univ.) An inversion formula utilizing hyperfunctions and it's application

概要 We show that an inversion formula for probability measures on the real line holds in a sense of the theory of hyperfunctions. As an application of our inversion formula, we give a representation of probability density functions utilizing characteristic functions.

- 2 高橋 博樹 (慶大理工) Lyapunov optimization for non-generic one-dimensional expanding Markov
 篠田 万穂 (慶大理工) maps 15
 Hiroki Takahashi (Keio Univ.) Lyapunov optimization for non-generic one-dimensional expanding Markov
 Mao Shinoda (Keio Univ.) maps

概要 For a non-generic, yet dense subset of C^1 expanding Markov maps of the interval we prove the existence of uncountably many Lyapunov optimizing measures which are ergodic, fully supported and have positive entropy. We also prove the existence of another non-generic dense subset for which the optimizing measure is unique and supported on a single periodic orbit. A key ingredient is a new C^1 perturbation lemma which allows us to interpolate between expanding Markov maps and the shift map on a finite number of symbols.

- 3 イェーリッシュヨハネス Spectral gap property for random dynamics on the real line and the
 (島根大総合理工) Hölder regularity of generalised Takagi functions 15
 角 大輝 (京大人間環境)
 Johannes Jaerisch (Shimane Univ.) Spectral gap property for random dynamics on the real line and the
 Hiroki Sumi (Kyoto Univ.) Hölder regularity of generalised Takagi functions

概要 We consider the random iteration of two expanding diffeomorphisms on the real-line without a common fixed point. We derive the spectral gap property of an associated transition operator acting on spaces of Hölder continuous functions. We introduce generalised Takagi functions on the real-line and we investigate their regularity properties.

- 4 四丸 直人 (岡山理大理) Another proof of Weyl's lemma 10
 高嶋 恵三 (岡山理大理)
 Naoto Shimaru (Okayama Univ. of Sci.) Another proof of Weyl's lemma
 Keizo Takashima
 (Okayama Univ. of Sci.)

概要 We give an upper bound for the discrepancy of irrational rotations $\{n\alpha\}$ in terms of the continued fraction expansion of α and the related Ostrowski expansion. Our result improves earlier bounds in the literature and substantially simplifies their proofs.

- 5 植村 英明 (愛知教育大教育) 一般 CONS の確率フーリエ係数による乱関数の復元について 15
 小川 重義 (立命館大理工)
 Hideaki Uemura (Aichi Univ. of Edu.) On the reconstruction of random function from its SFCs defined by an
 Shigeyoshi Ogawa (Ritsumeikan Univ.) arbitrary CONS

概要 We consider the reconstruction problem of a random function from the system of its stochastic Fourier coefficients (SFC in abbr.). We employ arbitrary orthonormal basis and Ogawa integral to construct SFCs. We first discuss the representation of Ogawa integral of a random function from its SFCs and H^1 basis, and then solve the reconstruction problem.

- 6 星野 浄生 (阪府大理) 有界変動過程の SFC による同定 15
 Kiyoki Hoshino (Osaka Pref. Univ.) Identification of finite variation processes from the SFC

概要 We consider the question whether a random function (or a stochastic derivative as an extension) is identified from the stochastic Fourier coefficient (SFC). We give an answer for the stochastic derivatives driven by finite variation processes. Especially, any finite variation process is identified from the SFC of Ogawa type. Also, we reconstruct, independently of values of the Brownian motion, nonnegative absolutely continuous noncausal Wiener functionals from the SFC of Skorokhod type.

- 7 星野 壮登 (早大理工) Global well-posedness of complex Ginzburg–Landau equation with a
 space-time white noise 15
 Masato Hoshino (Waseda Univ.) Global well-posedness of complex Ginzburg–Landau equation with a
 space-time white noise

概要 We show the global-in-time well-posedness of the complex Ginzburg–Landau (CGL) equation with a space-time Gaussian white noise on the 3-dimensional torus. The local well-posedness was obtained by Hoshino, Inahama and Naganuma, as an application of the theory of paracontrolled calculus. For the global well-posedness, we use a similar argument to Mourrat and Weber’s work about the global well-posedness of the dynamical Φ_3^4 model. By improving their method, we show a priori L^{2p} estimate of the solution for $p > \frac{3}{2}$.

- 8 星野 壮登 (早大理工) A coupled KPZ equation, its two types of approximations and existence
 舟木 直久 (早大理工) of global solutions 15
 Masato Hoshino (Waseda Univ.) A coupled KPZ equation, its two types of approximations and existence
 Tadahisa Funaki (Waseda Univ.) of global solutions

概要 We discuss the multi-component coupled Kardar–Parisi–Zhang (KPZ) equation and its two types of approximations. By applying the paracontrolled calculus introduced by Gubinelli, Imkeller and Perkowski, we show that these approximations have a common limit under well adjusted choices of renormalization factors. Moreover, if the coupling constants satisfy the so-called “trilinear” condition, then the Wiener measure becomes stationary for the limit, so that this limit exists globally in time when the initial value is sampled under the stationary measure.

- 9 和田 正樹 (福島大人間発達文化) Large time asymptotics for fundamental solutions of critical Schrödinger
 operators 15
 Masaki Wada (Fukushima Univ.) Large time asymptotics for fundamental solutions of critical Schrödinger
 operators

概要 Let $\{X_t\}_{t \geq 0}$ be the symmetric α -stable process with generator $\mathcal{L} = -(-\Delta)^{\alpha/2}$ for $0 < \alpha \leq 2$ and μ be a positive Radon measure in a certain class. We define the Schrödinger operator $\mathcal{L}^\mu = \mathcal{L} + \mu$ and consider the fundamental solution of the equation $\partial u / \partial t = \mathcal{L}^\mu u$. If μ is critical, the behavior of the fundamental solution is different from that of the transition density function of $\{X_t\}_{t \geq 0}$. In this talk, we give large time asymptotics for fundamental solutions of critical Schrödinger operators.

14:15~15:00

- 10 吉川 和宏 (立命館大理工) Composed order statistics and multivariate compound Poisson processes
 青山 崇洋 (岡山大環境理工) 15
 Kazuhiro Yoshikawa Composed order statistics and multivariate compound Poisson processes
 (Ritsumeikan Univ.)
 Takahiro Aoyama (Okayama Univ.)

概要 In this talk, we give an order statistic with random vectors to construct multivariate compound Poisson processes. The method works well for some processes, especially generated by zeta distributions. For example, Aoyama and Nakamura introduced generalized Euler products attached to a subclass of multidimensional infinitely divisible distributions. We will construct the compound Poisson processes corresponding their infinitely divisible distributions generated by the Euler products.

- 11 平尾 将剛 (愛知県立大情報) Frame potentials of determinantal point processes on the d -sphere 15
 Masatake Hirao (Aichi Pref. Univ.) Frame potentials of determinantal point processes on the d -sphere

概要 In the recent years, finite frame theory has come to draw a lot of attention since there exist many applications, e.g, numerical analysis, algebraic design theory, directional statistics, compressed sensing and so on. In this talk we show that determinantal point processes on the sphere give almost tight finite frames. We give two expectations of frame potentials of spherical ensembles and harmonic ensembles, which are the typical types of DPPs on the sphere. We also discuss random matrices induced by determinantal point processes on the sphere.

- 12 竹居 正登 (横浜国大工) 完全2部グラフ上の vertex-reinforced random walks 10
 石川 智啓 (横浜国大工)
 Masato Takei (Yokohama Nat. Univ.) Vertex-reinforced random walks on complete bipartite graphs
 Tomohiro Ishikawa
 (Yokohama Nat. Univ.)

概要 We consider vertex-reinforced random walks on complete bipartite graphs, and study their limiting behavior.

15:15~16:15 特別講演

- 岡村 和樹 (京大数理研) 複雑なグラフ上のランダムウォークの諸性質について
 Kazuki Okamura (Kyoto Univ.) Several properties for random walks on graphs

概要 In this talk we will state several properties for random walks on graphs. We will mainly focus on simple random walks on infinite connected graphs such as fractal graphs and percolation clusters. This talk will consist of two large parts. First, we will state the range of random walk on graphs satisfying a uniform condition, which includes several fractal graphs. Second, we will state a level-2 quenched large deviation principle for simple random walk on a class of percolation clusters including long-range correlations. The second part is based on a joint work with Noam Berger and Chiranjib Mukherjee.

16:30~17:30 特別講演

坂井 哲 (北 大 理) 強磁性イジング模型の相転移・臨界現象に関する研究の最近の動向
 Akira Sakai (Hokkaido Univ.) Recent progress in researches on phase transitions and critical behavior
 for Ising ferromagnets

概要 In this talk, I will review a class of most important results, chosen based on my personal preference, on phase transitions and critical behavior for the Ising model, a model of ferromagnetism in classical equilibrium statistical mechanics. First, I will overview the results obtained by the end of the previous century and recall the problems left unsolved then. Next, I will review how some of those problems have been solved since 2001. The common tool to solve most of those problems is the random-current representation. I will explain the derivation of the representation and its implication by showing some examples, such as exponential decay of the subcritical two-point function, uniqueness of the critical point, the mean-field bound on the 1-arm exponent, and the lace expansion for the two-point function. Finally, I will summarize the remaining open problems for future researches.

9月12日(火) 第I会場

9:20~11:30

- 13 藤田 敏治 (九 工 大 工) ノンシリアル動的計画 —合流型推移— 15
 才川 尚輝 (九 工 大 工)
 Toshiharu Fujita On nonserial dynamic programming —Converging branch systems—
 (Kyushu Inst. of Tech.)
 Naoki Saikawa (Kyushu Inst. of Tech.)

概要 In this study, we consider a new decision process model with a converging branch system which is one of the nonserial transition systems. We give the formulation of the model and introduce a recursive method to solve it by using dynamic programming.

- 14 松原 和樹 (中央学院大商) The existence of two-pairwise additive cyclic BIB designs of block size
 景山 三平 two 15
 (東京理大理数センター)
 Kazuki Matsubara (ChuoGakuin Univ.) The existence of two-pairwise additive cyclic BIB designs of block size
 Sanpei Kageyama (Tokyo Univ. of Sci.) two

概要 The existence of pairwise additive cyclic balanced incomplete block (BIB) designs with $k = 2$ and $\lambda = 1$ has been discussed in the literature. In this talk, for an odd prime $p \geq 5$, 2-pairwise additive cyclic BIB designs with $(v, k, \lambda) = (3p, 2, 1), (2p, 2, 2)$ are mainly constructed through methods of block replacements. Finally, the existence of 2-pairwise additive cyclic BIB designs with $k = 2$ and $\lambda \geq 1$ is shown entirely.

- 15 弓場 弘 (国際自然研) A*-optimal balanced third-order designs of resolution $R^*({10,01})$ with
兵頭 義史 $N < \nu(m)$ for 3^m factorials 15
(岡山理大総合情報研・国際自然研)
栗田 正秀 (国際自然研)
Hiromu Yumiba (Int. Inst. for Nat. Sci.) A*-optimal balanced third-order designs of resolution $R^*({10,01})$ with
Yoshifumi Hyodo $N < \nu(m)$ for 3^m factorials
(Okayama Univ. of Sci./Int. Inst. for Nat. Sci.)
Masahide Kuwada
(Int. Inst. for Nat. Sci.)

概要 We consider the third-order linear model for 3^m factorials. In previous talks (MSJ Autumn Meeting 2016; MSJ Spring Meeting 2017), we have established a necessary and sufficient condition for a simple array (SA) to be a balanced third-order (3^m -BTO) design of resolution $R^*({10,01})$, where the number of assemblies ($= N$) is less than the number of non-negligible factorial effects ($= \nu(m)$) and $m \geq 6$.

Let T be a 3^m -BTO design of resolution $R^*({10,01})$ derived from an SA with N assemblies, and further let $\sigma^2 S_T$ be the trace of the variance-covariance matrix of the estimators concerning with all the main effects based on T . If $S_T \leq S_{T^*}$ for any T^* , then T is said to be A*-optimal, where T^* is a 3^m -BTO design of resolution $R^*({10,01})$ derived from an SA with N assemblies. In this talk, we give A*-optimal 3^m -BTO designs of resolution $R^*({10,01})$ derived from SA's for $m = 6, 7, 8$, where $N < \nu(m)$.

- 16 盧 曉南 (東京理大理工) Locating arrays with error-correcting ability 15
神保 雅一 (中部大現代教育)
Xiao-Nan Lu (Tokyo Univ. of Sci.) Locating arrays with error-correcting ability
Masakazu Jimbo (Chubu Univ.)

概要 Locating arrays are introduced for identifying interaction faults and their locations in component-based systems. This class of problems are closely related with covering arrays and group testing, but the constructions are less known. Under the assumption that the system contains (at most) d faults, each involving (at most) t interacting factors, the notion of a (\bar{d}, t) -locating array is proposed. In this talk, I will focus on $(\bar{1}, t)$ -locating arrays. Moreover, by taking the similar consideration to error-correcting codes, the notion of locating arrays with error-correcting ability will be introduced.

- 17 内藤 貫太 (島根大総合理工) Regression with stagewise minimization on the risk 15
吉田 拓真 (鹿児島大理工)
Kanta Naito (Shimane Univ.) Regression with stagewise minimization on the risk
Takuma Yoshida (Kagoshima Univ.)

概要 This talk is concerned with regression based on the empirical risk minimization. The estimator is composed as a convex combination of the word (learner) in dictionary. The word is selected in each step of the proposed stagewise algorithm, which minimizes a certain divergence measure. A non-asymptotic error bound of the estimator is developed, and it is seen that the error bound becomes sharp as the number of iteration of the algorithm increases.

- 18 前園 宜彦 (九大数理) On direct kernel estimator of density ratio 10
 本山 真誠 (JA 全農)
 Yoshihiko Maesono (Kyushu Univ.) On direct kernel estimator of density ratio
 Masanari Motoyama (ZEN-NOH)

概要 Let $f(x)$ and $g(x)$ denote probability density functions and $g(x_0) \neq 0$. In this paper we discuss the density ratio $f(x_0)/g(x_0)$. A naive estimator is constituted from separate estimators of $f(x_0)$ and $g(x_0)$, which we call an indirect estimator. The other estimator is proposed by Cwik and Mielniczuk (1989), which we call a direct estimator. Here we propose a new direct estimator, and derive asymptotic mean squared error. We also prove central limit theorem of the new estimator. We also compare mean squared errors of the proposed estimator and others by simulation

- 19 柿沢 佳秀 (北大経済) 対称分布をベースにした q-MIG カーネル密度推定 15
 Yoshihide Kakizawa (Hokkaido Univ.) Symmetrical-based q-MIG kernel density estimation

概要 We consider estimation of the probability density for nonnegative data. In that case, the standard kernel density estimator is, in general, inconsistent near the boundary, due to the so-called boundary bias. Many authors have suggested various remedies, e.g., renormalization, reflection, and generalized jackknifing (see Jones (1993) for a review). On the other hand, over the last decade, there has been growing interest in the use of asymmetric kernel (AK), whose support matches the support of the density to be estimated. We propose a new AK density estimator using a q-MIG kernel. Here, “MIG” is a mixture of symmetrical-based inverse Gaussian (IG) and its reciprocal (RIG), and “q-MIG” is its generalization via Yang’s (2006) dual transformation, including a subfamily of log-symmetrical densities as a special case.

11:30~12:00 統計数学分科会総会

13:15~14:15

- 20 生 亀 清 貴 (東京理大理工) 順序カテゴリ正方分割表におけるダイヤモンドモデルの分解 10
 Kiyotaka Iki (Tokyo Univ. of Sci.) Decomposition of diamond model for square contingency tables with ordered categories

概要 For square contingency table with the same row and column ordinal classifications, this paper shows that the diamond model holds if and only if the weighted covariance for the difference between the row and column classifications and the sum of them equals zero and the uniform association diamond model holds.

- 21 田 畑 耕 治 (東京理大理工) 正方分割表における非対称モデルと対称性の分解 10
 Kouji Tahata (Tokyo Univ. of Sci.) Asymmetry model and decomposition of symmetry for square contingency tables

概要 For the analysis of square contingency tables with ordered categories, the present paper proposes a model that indicates the structure of asymmetry for cell probabilities. The model is the closest to the symmetry model in terms of the f -divergence under certain conditions, and includes the asymmetry models, which have been proposed by many statisticians, in the special cases. Also, it is shown the theorem that the symmetry model can be separated into some models by using the proposed model. It may be useful to see the reason for the poor fit of the symmetry model.

- 22 八木文香 (東京理大理) Transformed T^2 -type statistics for testing equality of two mean vectors
with monotone samples 15
瀬尾隆 (東京理大理)
Ayaka Yagi (Tokyo Univ. of Sci.) Transformed T^2 -type statistics for testing equality of two mean vectors
with monotone samples
Takashi Seo (Tokyo Univ. of Sci.)

概要 We consider the null distribution of the Hotelling's T^2 -type statistic for testing equality of two mean vectors when the two data matrices are of the same monotone missing pattern. As with the one-sample problem, a simplified T^2 statistic and an asymptotic expansion of its null distribution using decomposition of the test statistic are derived. Decomposition of the test statistic proposed in the study allowed to calculate the asymptotic expansion more easily. Further, we present the transformed test statistics based on the Bartlett adjustment. Finally, by a Monte Carlo simulation, we investigate the accuracy and asymptotic behavior of the approximation for chi-squared distribution.

- 23 兵頭昌 (阪府大工) 多標本問題に対するユークリッド距離を利用した平均ベクトルと分散共
小川颯 (阪府大工) 分散行列の同時検定 15
西山貴弘 (専修大経営)
Masashi Hyodo (Osaka Pref. Univ.) Simultaneous test of mean vector and variance covariance matrix using
Hayate Ogawa (Osaka Pref. Univ.) Euclidean norm for multi-sample problem
Takahiro Nishiyama (Senshu Univ.)

概要 In this talk, we propose an L^2 -norm-based statistic and its asymptotic distribution for simultaneous test of the mean vector and covariance matrix for multi-sample problem. An asymptotic distribution of test statistic are derived under a high-dimensional framework to deal with high-dimensional problems. This result is used for asymptotic size adjustment and derivation of asymptotic power. Finally, we study the finite sample and dimension performance of this test via Monte Carlo simulations.

9月13日(水) 第I会場

9:40~12:10

- 24 佃康司 (東大総合文化) 二つの一般リッジ推定量を等しくする共分散構造 15
倉田博史 (東大総合文化)
Koji Tsukuda (Univ. of Tokyo) Covariance structure associated with equality between two general ridge
Hiroshi Kurata (Univ. of Tokyo) estimators

概要 The Rao structure about the dispersion matrix in the general linear model is a well-known necessary and sufficient condition which guarantees that the ordinary least square estimator becomes the best linear unbiased estimator. In this presentation, we discuss a general ridge estimator to derive an extension of the Rao structure, that is, a necessary and sufficient condition under which two general ridge estimators coincide with each other.

- 25 石井 晶 (東京理大理工) 高次元データにおける固有空間の構造に基づいた共分散行列の同等性検定
 矢田 和善 (筑波大数理物質) 15
 青嶋 誠 (筑波大数理物質)
 Aki Ishii (Tokyo Univ. of Sci.) Equality tests of high-dimensional covariance matrices based on eigen-
 Kazuyoshi Yata (Univ. of Tsukuba) structures
 Makoto Aoshima (Univ. of Tsukuba)

概要 In this talk, we consider the equality test of covariance matrices for high-dimensional data. Aoshima and Yata (2017, SS) proposed two eigenvalue models for high-dimensional data. One is called the strongly spiked eigenvalue (SSE) model and the other one is called the non-SSE (NSSE) model. Li and Chen (2012) proposed a test statistic under the NSSE model. We verify that the statistic is asymptotically distributed as a chi-squared distribution under the SSE model. With the help of the asymptotic distribution, we proposed an equality test under the SSE model.

- 26 矢田 和善 (筑波大数理物質) Asymptotic normality for inference on high-dimensional mean vectors
 青嶋 誠 (筑波大数理物質) under the SSE model 15
 Kazuyoshi Yata (Univ. of Tsukuba) Asymptotic normality for inference on high-dimensional mean vectors
 Makoto Aoshima (Univ. of Tsukuba) under the SSE model

概要 In this talk, we consider the asymptotic normality for inference on high-dimensional mean vectors under two disjoint models: the strongly spiked eigenvalue (SSE) model and the non-SSE (NSSE) model. We first consider a distance-based statistics. We verify that it is asymptotically distributed as a normal distribution under the NSSE model. We also show that the asymptotic normality does not hold under the SSE model. We propose a new statistics by the estimation of eigenstructures for the SSE model. We verify that the proposed statistics is asymptotically distributed as a normal distribution under the SSE model. With the help of the asymptotic normality, we consider inferences on multi-sample and mean vectors under the SSE model.

- 27 中山 優吾 (筑波大数理物質) Asymptotic properties of support vector machines in high-dimension,
 矢田 和善 (筑波大数理物質) low-sample-size settings 15
 青嶋 誠 (筑波大数理物質)
 Yugo Nakayama (Univ. of Tsukuba) Asymptotic properties of support vector machines in high-dimension,
 Kazuyoshi Yata (Univ. of Tsukuba) low-sample-size settings
 Makoto Aoshima (Univ. of Tsukuba)

概要 In this talk, we consider asymptotic properties of the support vector machine (SVM) in high-dimension, low-sample-size (HDLSS) settings. We show that the SVM holds a consistency property in which misclassification rates tend to zero as the dimension goes to infinity under certain severe conditions. We show that the SVM is very biased in HDLSS settings and its performance is affected by the bias directly. In order to overcome such difficulties, we propose a bias-corrected SVM (BC-SVM). We show that the BC-SVM gives preferable performances in HDLSS settings for typical kernel functions. Finally, we check the performance of the BC-SVM by numerical simulations.

- 28 Yujie Xue (早大理工) Modified LASSO estimators of the models with long-memory disturbances 10
 谷口正信 (早大理工)
Yujie Xue (Waseda Univ.) Modified LASSO estimators of the models with long-memory disturbances
 Taniguchi Masanobu (Waseda Univ.)

概要 When we deal with actual problems by model building, it is often commonly assumed that the response variable and covariates satisfy linear relationship. One of the usual assumptions is that the disturbances follow identically independent distribution. Nevertheless the correlation of them may occur when the data are collected sequentially in time, especially in the field of economics and geophysics. In this talk, we assume the errors are strongly dependent. Then the asymptotic theory for modified LASSO estimators is discussed.

- 29 藤森 洸 (早大理工) The Dantzig selector for high-dimensional linear models of diffusion processes 10
 Kou Fujimori (Waseda Univ.) The Dantzig selector for high-dimensional linear models of diffusion processes

概要 The Dantzig selector, which was proposed by Candés and Tao in 2007, is an estimation procedure for regression models in a high-dimensional and sparse setting. In this presentation, linear models of diffusion processes with unknown drift matrices and diagonal diffusion matrices are discussed. We will consider the estimation problems for drift and diffusion matrices based on the discrete time observation in high-dimensional and sparse settings for drift matrices. To estimate drift matrices, we will apply the Dantzig selector and prove the l_q consistency of the estimator for every $q \in [1, \infty]$ under some appropriate conditions.

- 30 長幡英明 (早大理工) Analysis of variance for high dimensional time series 15
 谷口正信 (早大理工)
Hideaki Nagahata (Waseda Univ.) Analysis of variance for high dimensional time series
 Masanobu Taniguchi (Waseda Univ.)

概要 For independent observations, analysis of variance (ANOVA) has been enough tailored. Recently there has been much demand for ANOVA of high dimensional and dependent observations in many fields. However ANOVA for high dimensional and dependent observations has been immature. In this paper, we study ANOVA for high dimensional and dependent observations. Specifically, we show asymptotics of classical tests proposed for independent observations and give a sufficient condition for them to be asymptotically normal. Some numerical examples for simulated and real financial data are given as applications of these results. The extension in this paper is not straightforward and contains a lot of novel aspects for the analysis of variance for high dimensional time series.

- 31 高野嘉寿彦 (信州大工学部) 1次自己回帰過程の空間の幾何学的性質について 10
 Kazuhiko Takano (Shinshu Univ.) Geometric properties of system spaces of autoregressive process of degree 1

概要 We study a system space of autoregressive process of degree 1. System spaces of time series in information geometry have a Fisher metric as a Riemannian metric, and admit an α -connection which is defined by the power spectrum. This space is a two dimensional α -flat statistical manifold. Moreover, for $\alpha = -1, 0, 1$ we discuss α -geodesics and almost complex structures which are parallel with respect to the α -connection.

- 32 劉 言 (早大理工) A test for stationarity by copula spectral density 15
 Yan Liu (Waseda Univ.) A test for stationarity by copula spectral density

概要 We consider a hypothesis testing problem on the stationarity in locally stationary processes. In the existing literature, several test statistics have been proposed in the framework of local periodogram generated from the local stationary processes. However, we need sufficient moments of the stochastic processes under that framework. Thus, we propose a new test statistic constructed from the local quantile periodogram, where the measure of stationarity is also redefined in the quantiles. In addition, we extend our test statistic to the empirical likelihood ratio statistic to test the hypothesis. The theoretical results and numerical results under the alternative hypotheses will be given in the talk.

14:15~15:15 特別講演

- 清 智也 (東大情報理工) 多次元確率分布のスケーリング問題
 Tomonari Sei (Univ. of Tokyo) A scaling problem of multi-dimensional probability distributions

概要 We discuss a scaling problem of continuous probability distributions on the Euclidean space, where scaling means a coordinate-wise transformation in order that some functional identity is satisfied. In linear algebra, it is known that any non-negative definite matrix has a unique diagonal scaling such that the transformed matrix is quasi-doubly stochastic, whenever it is strictly copositive. We generalize the result to the space of probability distributions, where the set of matrices is identified with the Gaussian family. It is shown that, under a strictly copositive condition for distributions, there exists a unique coordinate-wise transformation such that the transformed distribution satisfies a Stein-type identity. The result is interpreted as an alternative representation of copulas. The proof is based on an energy minimization problem over a subset of the Wasserstein space. Some open problems will be discussed.

15:30~16:30 特別講演

- 明石 郁哉 (早大理工) 経験尤度法・基準化法に基づく非正規時系列モデルの頑健な統計的推測法の構成
 Fumiya Akashi (Waseda Univ.) Robust statistical inference for non-standard time series models based on the empirical likelihood and normalization methods

概要 This talk introduces robust statistical inference for various time series models under non-standard settings. In these few decades, non-standard aspects of real data in some sense are frequently observed in practical situations. First, a prominent example of long-range dependence was found by Hurst (1951, Trans. Amer. Soc. Civil Eng.) via the analysis of records of water flows through the Nile and though other rivers. Second, Mandelbrot (1963, J. Polit. Econ.) and Fama (1965, J. Bus.) found heavy-tailed economic and financial data which were poorly captured by the Gaussian models. When a statistical model has long-range dependence and/or heavy-tails, the limit distributions of fundamental statistics (e.g., sample mean) are not expressed in a closed form, and the rate of convergence contains the Hurst-index of long-range dependence and the tail-index of the underlying innovation density. Such properties make the situation complicated, and it is unfeasible to use the classical maximum likelihood method or the method of moments directly. To overcome the hurdles, we make use of some statistical methodologies involving the empirical likelihood, self-weighting and self-normalization methods. The empirical likelihood method proposed by Owen (1988, Biometrika) is a modern important statistical framework without knowledge of the underlying distribution. In particular, we integrate the concepts of the empirical likelihood and the least absolute deviations-based self-weighting method proposed by Ling (2005, J. Roy. Stat. Soc.), and construct the robust empirical likelihood statistic which is not affected by the nuisance parameters of the model and has the standard chi-square limit distribution. On the other hand, we also overcome the difficulties brought by the long-range dependence by using the self-normalized subsampling method proposed by Bai et al. (2016, Ann. Stat.). Finally, a unified, feasible and robust framework for various time series models under the non-standard situation is established.

応 用 数 学

9月11日(月) 第IV会場

10:00~12:00

- 1 佐竹翔平 (神戸大システム情報) Constructions of Ramanujan graphs and related results 15
 Shohei Satake (Kobe Univ.) Constructions of Ramanujan graphs and related results

概要 Ramanujan graph were defined by Lubotzky–Phillips–Sarnak in 1988. It is well known that Ramanujan graphs have nice properties as networks. In combinatorics, to give explicit constructions of Ramanujan graphs is recognized as a very interesting problems. In this talk, we give some explicit constructions of Ramanujan graphs as Cayley graphs over finite fields and rings. Moreover, we will discuss some properties which our graphs have.

- 2 足立智子 (東邦大理) 完全二部グラフの順序付けと長方形 RAID への応用 10
 Tomoko Adachi (Toho Univ.) Ordering of the complete bipartite graph and its application to rectangular RAID system

概要 We replace the problem of the RAID system in computer science with the problem of cyclic orderings in graph theory. We pay attention to constructions of cyclic orderings called cluttered orderings. There have been several studies on cluttered orderings of the complete bipartite graph $K_{\ell,\ell}$. Mueller et al. presented cluttered orderings in the case of $\ell = 3t, 10t$. In this paper, we investigate cluttered orderings of the complete bipartite graph $K_{\ell,m}$.

- 3 藤村 丞 On the number of perfect matchings of line graphs 10
 (福岡大情報基盤センター)
 白石修二 (福岡大理)
 Sho Fujimura (Fukuoka Univ.) On the number of perfect matchings of line graphs
 Shuji Shiraiishi (Fukuoka Univ.)

概要 We give a method for counting perfect matchings in line graphs. Consequently, for graphs G of maximum degree at most 3, we give a closed formula for the number of perfect matchings in $L(G)$.

- 4 向江頼士 (木更津工高専) Impurity of projective planar graphs 15
 佐野照和 (木更津工高専)
 Raiji Mukae Impurity of projective planar graphs
 (Kisarazu Nat. Coll. of Tech.)
 Terukazu Sano
 (Kisarazu Nat. Coll. of Tech.)

概要 We shall determine the impurity of projective planar graphs. The concept of impurity is related to edge-maximal graphs.

- 5 田中康平 (信州大経済) 離散的オイラー積分と数え上げ問題への応用 15
 Kohei Tanaka (Shinshu Univ.) Discrete Euler calculus and its application to the counting problem

概要 This talk introduces the integration theory with respect to the Euler characteristics of posets (categories), as a discrete analog of Baryshnikov and Ghrist's work on topological Euler calculus. As its application, we consider the counting problem in sensor network theory. We enumerate targets which lie on an acyclic network graph with sensors detecting the targets, by using discrete Euler calculus.

- 6 本田 あおい (九工大情報工) 包除積分の単調性と interaction operator 15
岡崎 悦明 (フuzzyシステム研)
Aoi Honda (Kyushu Inst. of Tech.) Monotonicity of inclusion-exclusion integral and interaction operator
Yoshiaki Okazaki
 (Fuzzy Logic Systems Inst.)

概要 We have proposed the inclusion-exclusion integral which is an integral with respect to monotone measure and interaction operator. In this talk, we show a concrete way of construction of interaction operator based on t-norm and give a sufficient condition for monotonicity of the inclusion-exclusion integral. Moreover we give several examples of the operators which satisfy this sufficient condition.

- 7 佐々木 義卓 (大阪体育大) 制限付きロンサム行列の数え上げ 10
大野 泰生 (東北大理)
Yoshitaka Sasaki Counting restricted lonesum matrices
 (Osaka Univ. of Health and Sport Sci.)
Yasuo Ohno (Tohoku Univ.)

概要 A lonesum matrix is a $(0,1)$ -matrix which can be uniquely reconstructed from its row and column sums. We plan to talk about counting restricted lonesum matrices. We also discuss recurrence formulas for poly-Bernoulli numbers which derived from such counting formulas.

- 8 水澤 篤彦 Simple non-confusing travel groupoid の特徴づけと cycle 上の数え上げ 15
松本ディオゴけんじ
 (芝浦工大工)
Atsuhiko Mizusawa On characterization of simple non-confusing travel groupoids
Diogo Kendy Matsumoto
 (Shibaura Inst. of Tech.)

概要 A travel groupoid is an algebraic structure which has information of a graph and walks (paths) on the graph. We study a special travel groupoid called a simple non-confusing travel groupoid and characterize it by spanning trees on the graph associated to the simple non-confusing groupoid. We also count the number of simple non-confusing groupoids on cycle graphs.

14:15~15:45

- 9 松原 和樹 (中央学院大商) Continuous flattening of α -trapezoidal polyhedra 15
奈良 知恵 (明大MIMS)
Kazuki Matsubara (ChuoGakuin Univ.) Continuous flattening of α -trapezoidal polyhedra
Chie Nara (Meiji Univ.)

概要 It was proved that any orthogonal polyhedron is continuously flattened by using a property of a rhombus. We investigated the method precisely, and found that there are infinitely many ways to flatten such polyhedra. In this talk, we prove that the infimum of the area of moving creases is zero for α -trapezoidal polyhedra. As a by-product we provide a continuous flattening motion whose area of moving creases is arbitrarily small for more general types of polyhedra.

- 10 安藤 清 (国立情報学研・JST ERATO) A new forbidden subgraph for 5-contractible edges 15
 Kiyoshi Ando (Nat. Inst. of Information/JST ERATO) A new forbidden subgraph for 5-contractible edges

概要 An edge of a k -connected graph is said to be k -contractible if the contraction of the edge results in a k -connected graph. A k -connected graph with no k -contractible edge is said to be contraction-critically k -connected. Kawarabayashi showed that $K_1 + (P_3 \cup K_2)$ is a forbidden subgraph of contraction-critically 5-connected graphs. We present a new forbidden subgraph which has $K_1 + (P_3 \cup K_2)$. This is an extension of the previous result due to Kawarabayashi.

- 11 斎藤 明 (日大文理) Rainbow forbidden subgraphs in edge-colored graphs 15
 C. Magnant (Georgia Southern Univ.)
 Akira Saito (Nihon Univ.) Rainbow forbidden subgraphs in edge-colored graphs
 Colton Magnant (Georgia Southern Univ.)

概要 A pair (G, c) of a graph and an edge-coloring $c: E(G) \rightarrow \mathbf{N}$ is called an edge-colored graph. If c is an injection, we say that (G, c) is a rainbow graph. For a connected graph H , an edge-colored graph (G, c) is said to be rainbow H -free if G does not contain a rainbow subgraph which is isomorphic to H . By definition, if H' is a connected subgraph of H , every rainbow H' -free graph is rainbow H -free. In this talk, we report a reverse phenomenon. Let $K_{1,k}^+$ denote the graph of order $k+2$ which is obtained from $K_{1,k}$ by performing a simple subdivision to one edge. Then we show that every rainbow $K_{1,k}^+$ -free complete graph edge-colored in sufficiently many colors is rainbow $K_{1,k}$ -free. We also show that for a connected graph H and a connected proper subgraph H' , if every rainbow H -free complete graph edge-colored in sufficiently many colors is rainbow H -free, then $(H', H) = (K_{1,k}, K_{1,k}^+)$ for some k .

- 12 藤田 慎也 (横浜市国際総合) 辺着色グラフの次数条件と閉路の存在について 10
 Shinya Fujita (Yokohama City Univ.) Degree conditions for cycles in edge-colored graphs

概要 In this talk, some recent results on degree conditions for properly colored cycles and rainbow cycles in edge-colored graphs will be reviewed.

- 13 土屋守正 (東海大理) On series parallel orders and strict-double-bound graphs 10
 田代慎一郎 (東海大理)
 Morimasa Tsuchiya (Tokai Univ.) On series parallel orders and strict-double-bound graphs
 Shinichiro Tashiro (Tokai Univ.)

概要 For a poset P , the strict-double-bound graph (sDB-graph $\text{sDB}(P)$) is the graph on $V(P)$ for which vertices u and v of $\text{sDB}(P)$ are adjacent if and only if u is not v and there exist x and y in $V(P)$ distinct from u and v such that x is a lower bound of u, v and y is an upper bound of u, v . For a poset P , P is a series parallel order if P contains no induced subposet isomorphic to the N -poset. We obtain the following result. For a series parallel order P , if P_3 is an induced subgraph of a component with at least four vertices of $\text{sDB}(P)$, then P_3 is contained C_4 , $K_4 - e$, $K_{1,3}$ or 3-pan as an induced subgraph.

- 14 早水 桃子 Universal tree-based network の最小サイズについて 15
 (統計数理研・JST さきがけ)
鍛冶 静雄
 (山口大理・JST さきがけ)
藤重 悟 (京大数理研・京大*)
Momoko Hayamizu On the size of universal tree-based networks
 (Inst. of Stat. Math./JST PRESTO)
Shizuo Kaji
 (Yamaguchi Univ./JST PRESTO)
Satoru Fujishige
 (Kyoto Univ./Kyoto Univ.*)

概要 A tree-based network on a set X of leaves is said to be universal if any rooted binary phylogenetic tree on X can be its base tree. In my earlier work, the concept of universal tree-based networks was defined and it was shown that there exist infinitely many universal tree-based networks for any number $|X|$ of leaves. In this talk, I will discuss the minimum size of universal tree-based networks. This talk is based on joint work with Satoru Fujishige (RIMS, Kyoto University) and Shizuo Kaji (Yamaguchi University, JST PRESTO).

16:00~17:00 特別講演

- 千葉 周也 (熊本大先端科学) 指定した成分数の 2-因子と次数和条件
Shuya Chiba (Kumamoto Univ.) 2-factors with a specified number of components and degree sum conditions

概要 O. Ore (1960) proved that if the degree sum of every pair of non-adjacent vertices is at least the order of the graph, then the graph is hamiltonian. The study of sufficient conditions on degrees for the existence of a Hamilton cycle started from such a classical result. In this talk, as one of the generalizations of Ore's theorem, we focus on degree sum conditions for the existence of a 2-factor with a specified number of components in general simple graphs, bipartite graphs and directed graphs, and we survey results including recent progress on the research field. We also discuss the difference from the results on Hamilton cycles.

9月12日(火) 第IV会場

9:30~12:00

- 15 三橋 秀生 (法政大理工) 有限グラフ上の四元数 Szegedy ウォークと第 2 種重み付きゼータ関数 .. 15
今野 紀雄 (横浜国大理工)
松江 要
 (九大IMI・九大I2CNER)
佐藤 巖 (小山工高専)
Hideo Mitsuhashi (Hosei Univ.) Quaternionic quantum walks of Szegedy type on finite graphs and second weighted zeta functions
Norio Konno (Yokohama Nat. Univ.)
Kaname Matsue
 (Kyushu Univ./Kyushu Univ.)
Iwao Sato (Oyama Nat. Coll. of Tech.)

概要 We define a quaternionic analogue of the Szegedy walk on a graph and study its right spectral properties. The condition for the transition matrix of the quaternionic Szegedy walk on a graph to be quaternionic unitary is given. In order to derive the spectral mapping theorem for the quaternionic Szegedy walk, we show a quaternionic analogue of the determinant expression of the second weighted zeta function of a graph. Our main results determine explicitly all the right eigenvalues of the quaternionic Szegedy walk by using complex right eigenvalues of the corresponding doubly weighted matrix. We also show the manner of obtaining eigenvectors corresponding to right eigenvalues derived from those of doubly weighted matrix.

- 16 佐藤 巖 (小山工高専) A generalized Bartholdi zeta function of a graph 15
 三橋 秀生 (法政大理工)
 森田 英章 (室蘭工大工)
 Iwao Sato (Oyama Nat. Coll. of Tech.) A generalized Bartholdi zeta function of a graph
 Hideo Mitsuhashi (Hosei Univ.)
 Hideaki Morita
 (Muroran Inst. of Tech.)

概要 We define an $(n + 1)$ -variable Bartholdi zeta function and an $(n + 1)$ -variable Bartholdi L -function of a graph G , and give determinant expressions of them. We present a decomposition formula for the $(n + 1)$ -variable Bartholdi zeta function of a regular covering of G . Furthermore, we express the $(n + 1)$ -variable Bartholdi zeta function of a regular covering of G as a product of its $(n + 1)$ -variable Bartholdi L -functions.

- 17 今野 紀雄 (横浜国大工) Partition-based quantum walk 15
 R. Portugal (LNCC)
 佐藤 巖 (小山工高専)
 瀬川 悦生 (東北大情報)
 Norio Konno (Yokohama Nat. Univ.) Partition-based quantum walk
 Renato Portugal (LNCC)
 Iwao Sato (Oyama Nat. Coll. of Tech.)
 Etsuo Segawa (Tohoku Univ.)

概要 We introduce an extension model called partition-based quantum walk, which includes most quantum walk models driven by two local operators, such as the coined model, Szegedy's model, and the 2-tessellable staggered model. We show that all those families of quantum walk models using two local operators are unitary equivalent. The new framework is based on two equivalence-class partitions of the computational basis, which establishes the notion of local dynamics.

- 18 松江 要 单体的量子ウォーク version 2 —グラフ上 coined walk との対応— 15
 (九大IMI・九大I2CNER)
 小栗 栖修 (金沢大理工)
 瀬川 悦生 (東北大情報)
 Kaname Matsue Simplicial quantum walks version 2 —correspondence to coined walks
 (Kyushu Univ./Kyushu Univ.) on graphs—
 Osamu Ogurisu (Kanazawa Univ.)
 Etsuo Segawa (Tohoku Univ.)

概要 We propose a new version of quantum walks on simplicial complexes (named simplicial quantum walk), which is an alternative of preceding studies by authors. We show that it is unitary equivalent to a bipartite walk on associated bipartite graphs, coined quantum walk on a graph. Moreover, if simplicial complexes are orientable, the simplicial quantum walk is unitary equivalent to coined quantum walk on a graph with duplication structure.

- 19 松江 要 (九大IMI・九大I2CNER) 単体の量子探索 15
 小栗 栖 修 (金沢大理工)
 瀬川 悦生 (東北大情報)
Kaname Matsue (Kyushu Univ./Kyushu Univ.) Simplicial quantum search
 Osamu Ogurisu (Kanazawa Univ.)
 Etsuo Segawa (Tohoku Univ.)

概要 Here we show that the quantum search on the specific simplicial complex corresponding to the triangulation of n -dimensional unit square driven by this new simplicial quantum walk works well, namely, a marked simplex can be found with probability $1 + o(1)$ with in a time $O(\sqrt{N})$, where N is the number of simplices with the dimension of marked simplex.

- 20 吉江 佑介 (東北大情報) A periodicity of the Grover walk on Bethe trees 10
 瀬川 悦生 (東北大情報)
 谷口 哲至 (広島工大工)
 久保田 匠 (東北大情報)
Yusuke Yoshie (Tohoku Univ.) A periodicity of the Grover walk on Bethe trees
 Etsuo Segawa (Tohoku Univ.)
 Tetsuji Taniguchi (Hiroshima Inst. of Tech.)
 Sho Kubota (Tohoku Univ.)

概要 A Bethe tree is a rooted tree such that in each level the vertices have equal degree. In this paper we focus on the periodicity of the Grover walk on Bethe trees. The Grover walk is a kind of quantum walks on graphs, and the time evolution operator of the Grover walk is determined by the graph. A periodicity is a special feature of quantum walk, and we have found some graphs to induce a periodic Grover walk. We find the classes of Bethe trees which induce a periodic Grover walk under an assumption.

- 21 船川 大樹 (北大理) 2次元4状態量子ウォークの局在化について 15
 布田 徹 (北大理)
 笹山 智司 (北大理)
 鈴木 章斗 (信州大工)
Daiju Funakawa (Hokkaido Univ.) Localization of a 2-dimensional 4-state quantum walk
 Toru Fuda (Hokkaido Univ.)
 Satoshi Sasayama (Hokkaido Univ.)
 Akito Suzuki (Shinshu Univ.)

概要 We consider a discrete-time 2-dimensional 4-state quantum walk. The evolution of the quantum walk is described by a unitary operator U , which is the product of a space-dependent coin operator and a shift operator weighted with probabilities q_1 and q_2 . Supposing that the coin operator and the shift operator are self-adjoint and unitary,

Fuda, Funakawa, and Suzuki proved that localization occurs $|q| = \sqrt{|q_1|^2 + |q_2|^2}$ is sufficiently small. In this talk, we study about the localization without smallness of $|q|$.

- 22 布田 徹 (北 大 理) 1次元スプリット・ステップ量子ウォークの弱極限分布 15
 船川 大樹 (北 大 理)
 鈴木 章斗 (信州大工)
 Toru Fuda (Hokkaido Univ.) Weak limit distribution of a one-dimensional split-step quantum walk
 Daiju Funakawa (Hokkaido Univ.)
 Akito Suzuki (Shinshu Univ.)

概要 In this talk, we consider a discrete-time one-dimensional two-state quantum walk called a split-step quantum walk. The evolution operator of the split-step quantum walk is defined by the product of a shift operator and a space-dependent coin operator. Weak limit theorem of the split-step quantum walk and its explicit limit distribution are presented.

- 23 一階 智弘 (名大多元数理) 超群とグラフ上の酔歩 15
 澤田 友佑 (名大多元数理)
 Tomohiro Ikkai (Nagoya Univ.) Hypergroups and random walks on graphs
 Yusuke Sawada (Nagoya Univ.)

概要 Wildberger gave a way to construct a finite hypergroup from a random walk on a certain kind of finite graphs. His method is applicable to a random walk on a certain kind of infinite graphs. In this talk, we formulate his method and give some examples that produce hypergroups and that do not produce hypergroups.

9月13日(水) 第IV会場

9:30~11:40

- 24 堀口 俊二 拡張ニュートン法とフラクタル (実関数の場合) 15
 Shunzi Horiguchi Extended Newton method and fractals (in case of real variable function)

概要 The convergences of the extended Newton method to the roots of different initial values cause catastrophe, chaos, and fractals.

- 25 大林 一平 (東北大AIMR) パーシステントホモロジーと機械学習を用いたデータ解析 15
 Ipei Obayashi (Tohoku Univ.) Data analysis using persistence homology and machine learning

概要 In this talk, I will show you methods of data analysis using persistent homology and machine learning. Persistent homology enables us to describe the shape of data quantitatively from the viewpoint of homology and it is useful to study heterogeneous geometric structures. Machine learning enables us to detect characteristic patterns from data. By the combination of persistence homology and machine learning, we can quantitatively and statistically find characteristic geometric pattern hidden behind the data. Persistence Image and linear machine learning models are used for our methods. This combination gives us a very intuitive visualization of the learned result. "Inverse Problem" techniques for persistence diagrams are also effectively used to visualize the learned result.

- 26 坂口文則 (福井大工) 微分方程式整数型解法と差分方程式と数論との接点 15
 Fuminori Sakaguchi (Univ. of Fukui) Some possible relationships among an integer-type algorithm for solving ODEs, difference equations and number theory

概要 A integer-type algorithm for accurately solving linear ODEs by means only of four arithmetic operations among integers had been proposed by the author. Some direct ‘decipherments’ of numerical results by this algorithm enable us to see what is the essence of the accuracy of this algorithm, because this algorithm uses ‘exact’ Gaussian-integer-valued expansion coefficient sequences. By the decipherments, in this study, it turned out that this algorithm is closely related to number theory, in that a numerical expansion coefficient sequence by this algorithm is a rational linear combination of rational solution sequences of homogeneous linear difference equations with non-constant rational coefficients which accurately approximates their finite-norm irrational linear combination described in terms of algebraic extension of rational field. The relationship of this algorithm to continued fractions can be explained in this context.

- 27 三宅常時 (宇部工高専) 非線形自律系の定常解を追跡する一計算法 10
 勝田祐司 (宇部工高専)
 George Miyake (Ube Nat. Coll. of Tech.) A computational method for stationary solutions in nonlinear autonomous dynamical system
 Yuji Katsuta (Ube Nat. Coll. of Tech.)

概要 A computational method for equilibria of a nonlinear autonomous system is considered by using bisection method. The method led to analyze dynamical behavior and qualitative properties of the autonomous system.

- 28 穴田浩一 (早大高等学院) スケール不変性を利用した爆発レートの数値的推定について 15
 石渡哲哉 (芝浦工大システム理工)
 牛島健夫 (東京理大理工)
 Koichi Anada (Waseda Univ. Senior High School) A numerical method of estimating blow-up rates for nonlinear evolution equations by using rescaling algorithm
 Tetsuya Ishiwata (Shibaaura Inst. of Tech.)
 Takeo Ushijima (Tokyo Univ. of Sci.)

概要 In this talk, we proposed a numerical method for estimating blow-up rate of blow-up solutions for a class of nonlinear evolution equations which have a scaling invariance. To use this scaling invariance we adopt the rescaling algorithm to the problems and numerically estimate the blow-up rates. Applying the method to several examples, we examine the effectiveness of the method.

- 29 宇田智紀 (京大理) 形状微分を用いた周期定常渦斑の数値計算 15
 Tomoki Uda (Kyoto Univ.) Shape derivative and its application to vortex patch equilibria in a periodic array

概要 We propose a new shape derivative formula for contour integrals with logarithmic kernels which yields a numerical scheme to compute vortex patch equilibria. Owing to its simplicity, any steady configuration of point vortices can be extended to that of vortex patches. As a test problem, a periodic array of vortex patches is considered to show the efficiency of the new formula.

- 30 土屋 拓也 (早大理工) de Sitter 時空における半線形 Klein-Gordon 方程式の初期値問題における数値安定性について 15
中村 誠 (山形大理工)
Takuya Tsuchiya (Waseda Univ.) On the numerical analysis of the Cauchy problem for semi-linear Klein-Gordon equation in de Sitter spacetime
Makoto Nakamura (Yamagata Univ.)

概要 The numerical analysis of the Cauchy problem for semi-linear Klein-Gordon equation in the de Sitter spacetime is considered. The solution of the equation expresses the property of expansion or contraction depending on conditions. Some of the terms in the equation present the dissipative and antidissipative effects. Since it is difficult to study the property and the effects analytically, we investigate them with numerical simulations. In addition, we study the numerical stability of the solutions.

- 31 浦川 遼介 (早大理工) 測地線による Einstein 方程式の数値解の検証 15
土屋 拓也 (早大理工)
米田 元 (早大理工)
Ryosuke Urakawa (Waseda Univ.) Verification of Einstein's equation by geodesics
Tsuchiya Takuya (Waseda Univ.)
Yoneda Gen (Waseda Univ.)

概要 When we search for numerical solutions to the Einstein's equation, we typically monitor the conservation of the constraints as a sanity check against numerical errors in the solution. The conservation of constraints is a necessary but not sufficient condition for the solutions to the Einstein's equation. We propose a method by using the geodesic equation to improve the reliability of the solution.

14:15~16:15

- 32 木下 武彦 Hilbert 空間における線形作用素に対する逆作用素ノルム評価の改良 ... 15
渡部 善隆
 (九大情報基盤研究開発センター)
中尾 充宏 (早大理工)
Takehiko Kinoshita An improvement of norm bound computation for inverses of linear operators in Hilbert spaces
Yoshitaka Watanabe (Kyushu Univ.)
Mitsuhiro T. Nakao (Waseda Univ.)

概要 We propose a computer-assisted procedure to prove the invertibility of a linear operator in a Hilbert space and to compute a verified norm bound of its inverse. A number of the authors have previously proposed two verification approaches that are based on projection and constructive a priori error estimates. The approach of the present talk is expected to bridge the gap between the two previous procedures in actual numerical verifications. Several verification examples that confirm the actual effectiveness of the proposed procedure are reported.

- 33 剣持 智哉 (東大数理) 放物型方程式に対する有限要素法の時間について一様な誤差評価 15
Tomoya Kemmochi (Univ. of Tokyo) Time-global error estimates for finite element approximation of parabolic problems

概要 We consider the finite element approximation for the linear homogeneous parabolic problem with the homogeneous Neumann boundary condition. We assume that the elliptic operator in the equation does not have lower order terms, that is, the operator is not positive definite. In this case, both exact and approximate solutions are globally bounded; thus the error is also bounded uniformly in time. However, in many literature, they use the Gronwall inequality, which causes the exponentially increasing term with respect to the time variable. As far as we know, there are no literature on time-global error estimates for these problems. In this talk, we present the time-global L^∞ - L^p -error estimates for sufficiently smooth initial data.

- 34 千葉悠喜 (東大数理) N 次元球状領域上の Poisson 方程式に対する不連続 Galerkin 法 15
齊藤宣一 (東大数理)
Yuki Chiba (Univ. of Tokyo) Discontinuous Galerkin method for an N -dimensional spherically sym-
Norikazu Saito (Univ. of Tokyo) metric Poisson equation

概要 It is useful to make numerical method for nonlinear PDEs in higher dimension for researching critical phenomena of it. So, we consider a spherically symmetric Poisson equation in N -dimensional ball. The previous study proposed finite element method using weight function x^{N-1} and showed optimal weighted L^2 error estimate. However, there is a disadvantage of increasing error near origin. The another approach to use weight function x showed optimal L^∞ error estimate. In this paper, we see the PDE as singularly perturbed convection-diffusion equation through later approach, and apply discontinuous Galerkin method to it. We show some estimates and offer some numerical results.

- 35 中田行彦 (島根大総理工) Delay equations for reinfection dynamics 15
Yukihiko Nakata (Shimane Univ.) Delay equations for reinfection dynamics

概要 “Re”infection of recovered individuals, as a consequence of waning immunity and change of his/her susceptibility add further complexity in understanding disease transmission dynamics, forming a delayed feedback from infective population to susceptible population. In this talk we discuss dynamical aspects of a series of epidemic models, paying attention to reinfection dynamics, formulated by delay differential equations (DDE) and renewal equations (RE). We introduce a mathematical model by delay differential equations to provide a possible explanation of periodic outbreak of a childhood disease observed in Japan. Simple threshold dynamics is shown for a general SIS epidemic model formulated by a nonlinear renewal equation. On the other hand, we shall show that heterogeneous susceptibility can induce epidemic, after approaching to the trivial equilibrium.

- 36 鈴木宏昌 (滋賀大教育) 3変数反応拡散系における余次元2分岐の解析的および数値的考察 15
西浦廉政 (東北大AIMR)
Hiromasa Suzuki (Shiga Univ.) Analytical and numerical approaches to codimension-two bifurcation in
Yasumasa Nishiura (Tohoku Univ.) a three-component reaction-diffusion system

概要 We study the stability and the bifurcation structure of standing pulse solutions to a singularly perturbed three-component reaction-diffusion system. We can show the detailed information of dependence on the parameters about the stability properties and the bifurcation structure.

- 37 清水雄貴 (京大理) トーラス幾何による N 点渦環の安定化 15
坂上貴之 (京大理)
Yuuki Shimizu (Kyoto Univ.) Toroidal geometry stabilizing a ring of N point vortices on a torus
Takashi Sakajo (Kyoto Univ.)

概要 On a plane and a sphere, an N -ring is unstable for $N > 7$. We introduce on the inner side of a toroidal surface, however, an N -ring is stable when the aspect ratio of the torus is sufficiently large for any fixed N .

- 38 柴山允瑠 (京大情報)^b 制限 n 体問題の非可積分性 15
Mitsuru Shibayama (Kyoto Univ.) Non-integrability of the restricted n -body problem

概要 We prove the non-integrability of the spacial n -body problem. In order to prove it, we focus on the singularity of the extended differential equations and then apply the Morales–Ramis theory to it. We also discuss the non-integrability of the spacial restricted $n + 1$ -body problem.

16:30~17:30 特別講演

- 松江 要 サドルが紡ぐ縁 —えにし—: 精度保証付き数値計算と力学系
(九大IMI・九大I2CNER)
- Kaname Matsue Saddles create connections: Rigorous numerics and dynamical systems
(Kyushu Univ./Kyushu Univ.)

概要 Validated computations (rigorous numerics) have been applied to dynamical systems, such as the existence of equilibria, periodic orbits, connecting orbits and bifurcations and so on as well as their stability, for a couple of decades. I believe that one of true significance of rigorous numerics to dynamical systems is validation of objects which are very difficult to detect from both mathematical and numerical approach. The root will be either reduction of problems to fixed point problems for nonlinear maps or analysis based on saddle-type equilibria or invariant sets. In this talk, I develop an overview of rigorous numerics to dynamical systems derived from saddles in terms of topological tools such as isolating blocks, cones and Lyapunov functions, as well as various results by analytic approach.

9月14日(木) 第IV会場

9:30~10:30

- 39 大枝和浩(早大 GEC) Stationary problem of a diffusive prey-predator system with protection zones 15
Kazuhiro Oeda (Waseda Univ.) Stationary problem of a diffusive prey-predator system with protection zones

概要 This talk is concerned with the Neumann problem of a diffusive Lotka–Volterra prey-predator system with finitely many protection zones for the prey species. We discuss the stability of non-negative constant solutions. Moreover, we study the existence and non-existence of positive stationary solutions by applying the bifurcation theory.

- 40 寺本 敬(旭川医科大医) An action functional approach to localized patterns in a three-component FitzHugh–Nagumo model 10
P. van Heijster
(Queensland Univ. of Tech.)
Chao-Nien Chen
(Nat. Tsing-Hua Univ.)
西浦 廉政(東北大 AIMR)
Takashi Teramoto An action functional approach to localized patterns in a three-component FitzHugh–Nagumo model
(Asahikawa Medical Univ.)
Peter van Heijster
(Queensland Univ. of Tech.)
Chao-Nien Chen
(Nat. Tsing-Hua Univ.)
Yasumasa Nishiura (Tohoku Univ.)

概要 We analyze the stationary localized patterns in a singularly perturbed three-component FitzHugh–Nagumo model. We derive explicit conditions for the existence and stability of these type of localized solutions by combining geometric singular perturbation techniques and an action functional approach. The action functional replaces the Melnikov integral approach and Evans function computation to derive existence conditions and critical information on the stability of the localized patterns.

- 41 鈴木 貴 (阪大基礎工) 骨代謝モデリング—動的平衡の崩壊 5
Takashi Suzuki (Osaka Univ.) Bone metabolism modeling —break down of dynamical equilibrium

概要 We study break down of bone metabolism, using mathematical modeling. The principal part of this model is composed of two pathways of maturation, that is, from pre-osteoblast to osteoblast and from pre-osteoclast to osteoclast. There is also a pathway of acceleration to the formation of pre-osteoclast by pre-osteoblast. This pathway is evoked by a cytokine, called RANKL. Experimental data, on the other hand, suggest a differentiation annihilation factor, to the maturation pathways above. Here we formulate the above feedback loops as a system of ordinary differential equations, pick up dynamical equilibria, and study their break down.

- 42 鈴木 貴 (阪大基礎工) 細胞シグナルモデリング—減衰振動の再現性 5
Takashi Suzuki (Osaka Univ.) Cell signal modeling —reproducibility of damping oscillation

概要 The transcription factor NF-kB induces expression of multiple genes by shuttling between cytoplasm and nucleus. Previous studies have reported that the transcription of target gene is activated by phosphorylation of NF-kB, and decreased by dephosphorylation. In this study, we constructed a new mathematical model considering phosphorylation and mathematically analyzed how the phosphorylation of NF-kB effects on the oscillation phenomena. As a result, our new model, explained an appearance of a stable periodic orbit, which appeared in a transitional manner in response to the attenuation of an external stimulus, and also indicated that the NF-kB oscillation occurred by attracting to the periodic orbit.

- 43 西浦 廉政 (東北大AIMR) ナノ微粒子ミクロ相分離における縞模様からタマネギ型への形態遷移モ
E. Avalos (東北大AIMR) デリング 15
寺本 敬 (旭川医科大医)
藪 浩 (東北大AIMR)
Yasumasa Nishiura (Tohoku Univ.) Lamellae to onion transformation in block copolymer nanoparticles via
Edgar Avalos (Tohoku Univ.) coupled Cahn–Hilliard equations
Takashi Teramoto
(Asahikawa Medical Univ.)
Hiroshi Yabu (Tohoku Univ.)

概要 Annealing of copolymers has become a tool of great importance to reconfigure nanoparticles. We present experimental results of annealing copolymer nanoparticles and a mathematical model to describe the morphological transformation from lamellae to onion. A good correspondence between experimental findings and predictions of the model is observed. The model based on an appropriate free energy leads to a set of Cahn–Hilliard equations that correctly describes the dynamical transformation from lamellae particles to onion and reverse onion-like particles, regardless of the nature of the annealing process. This universality makes possible to describe a variety of experimental conditions involving nanoparticles underlying a heating process. A notable advantage of the proposed approach is that it makes possible to selectively control the interaction between the confined copolymer and the surrounding media.

10:45～11:45 特別講演

中野直人 (JSTさきがけ・北大理) Hilbert–Schmidt 積分作用素論による遅延座標埋め込み再考: 経験的モデリングのための力学再構成に向けて
Naoto Nakano (JST PRESTO/Hokkaido Univ.) Revisiting delay-embedding in terms of Hilbert–Schmidt integral operator theory: Towards dynamical reconstruction for empirical modelling

概要 Delay embedding is well-known for non-linear time-series analysis, and it is used in several research fields such as physics, informatics, neuroscience and so on. The celebrated theorem of Takens ensures validity of the delay embedding analysis: embedded data preserves topological properties, which the original dynamics possesses, if one embeds into some phase space with sufficiently high dimension. This means that, for example, an attractor can be reconstructed by the delay coordinate system topologically. However, configuration of an embedded dataset may easily vary with the delay width and the delay dimension, namely, “the way of embedding”. In a practical sense, this sensitivity may cause degradation of reliability of the method, therefore it is natural to require robustness of the result obtained by the embedding method in certain sense. In this study, we investigate the mathematical structure of the framework of delay-embedding analysis to provide Ansatz to choose the appropriate way of embedding, in order to utilise for time-series prediction. In short, mathematical theories of the Hilbert–Schmidt integral operator and the corresponding Sturm–Liouville eigenvalue problem underlie the framework. Using these mathematical theories, one can derive error estimates of mode decomposition obtained by the present method and can obtain the phase-space reconstruction by using the leading modes of the decomposition. In this talk, we will show some results for some numerical and experimental datasets to validate the present method.

トポロジー

9月11日(月) 第Ⅲ会場

10:00~12:00

- 1 門田直之(大阪電通大工) Signatures of surface bundles 15
 Naoyuki Monden Signatures of surface bundles
 (Osaka Electro-Comm. Univ.)

概要 The signature of a surface bundle over a surface has some restrictions, for examples, it is dividable by 4 and vanishes if the base genus is 0 or 1. Bryan and Donagi constructed examples over a genus-2 surface with non-zero signatures. However, the signatures and the genera of their examples are sporadic. In this talk, for any positive integer n , we give a surface bundle of fiber genus g over a surface of genus 2 with signature $4n$ and a section of self-intersection 0 if g is greater than or equal to $39n$.

- 2 門田直之(大阪電通大工) Stable commutator lengths of Dehn twists 15
 Naoyuki Monden Stable commutator lengths of Dehn twists
 (Osaka Electro-Comm. Univ.)

概要 In this talk, we give explicit factrizations of certain powers of Dehn twists as products of commutators. As a corollary, we improve upper bounds for stable commutator lengths of Dehn twists. Moreover, we show that the stable commutator length in the mapping class group is different from that in the hyperelliptic mapping class group for a surface of large genus.

- 3 直江央寛(東北大理) コルクとそのシャドウ複雑度 10
 Hironobu Naoe (Tohoku Univ.) Corks and their shadow complexities

概要 For a simply connected closed 4-manifold M , any 4-manifold exotic to M is obtained from X by twisting a contractible Stein domain called a cork. We study a cork from a viewpoint of the notion of shadows. A shadow of a 4-manifold is a simple polyhedron collapsed from the manifold. By using a shadow, Costantino defined a complexity of a 4-manifold, which is the minimum number of vertex of its shadow. We show that there are no corks with complexity zero and that there are infinitely many corks with complexity 1 and 2.

- 4 安井弘一(阪大情報) Nonexistence of twists and surgeries generating exotic 4-manifolds 15
 Kouichi Yasui (Osaka Univ.) Nonexistence of twists and surgeries generating exotic 4-manifolds

概要 It is well known that for any exotic pair of simply connected closed 4-manifolds, one is obtained by twisting the other along a contractible submanifold. In contrast, we show that for each positive integer n , there exists an infinite family of pairwise exotic simply connected closed 4-manifolds such that, for any 4-manifold X and any compact (not necessarily connected) codimension zero submanifold W with $b_1(\partial W) < n$, the family cannot be generated by twisting X along W and varying the gluing map. As a corollary, we show that there exists no ‘universal’ 4-manifold with boundary such that any exotic family is generated by twisting along an embedded copy of the 4-manifold. Moreover, we give similar results for surgeries.

- 5 大場 貴裕 (東工大理) Surfaces in D^4 with the same boundary and fundamental group 15
 Takahiro Oba (Tokyo Tech) Surfaces in D^4 with the same boundary and fundamental group

概要 This talk is concerned with symplectic surfaces in a symplectic 4-disk (D^4, ω) bounded by the same transverse link in the standard contact 3-sphere (S^3, ξ_{st}) . There are some examples of transverse links (or knots) bounding more than two distinct symplectic surfaces. All these surfaces can be distinguished by the fundamental groups of their complements. In this talk, I will present a family of pairs of two distinct symplectic surfaces whose boundaries are the same transverse knot and whose complements have isomorphic fundamental groups. To tell apart the two surfaces of each pair, I take double branched covers branched along them.

- 6 馬場 伸平 (Univ. Heidelberg) 複素射影構造のノード付き曲面への退化 15
 Shinpei Baba (Univ. Heidelberg) Neck-pinching of \mathbb{CP}^1 -structures on surfaces and convergence of holonomy representations

概要 A \mathbb{CP}^1 -structure on a surface is an atlas modeled on \mathbb{CP}^1 with transition maps in $\mathrm{PSL}(2, \mathbb{C})$, and a \mathbb{CP}^1 -structure corresponds to a pair of a Riemann surface and a holomorphic quadratic differential on it. In addition, each \mathbb{CP}^1 -structure has a holonomy representation from its fundamental group into $\mathrm{PSL}(2, \mathbb{C})$. In this talk, we consider a path of diverging \mathbb{CP}^1 -structures on a closed oriented surface such that, their holonomy representations converge. We discuss about its limit under the assumption that the Riemann surface structures are pinched along some loops.

- 7 高橋 典寿 (立命館大理工) 商空間がトーラスとなる対合と可換なリーマン面上の周期的写像の分類
 野沢 啓 (立命館大理工) 15
 Norihisa Takahashi (Ritsumeikan Univ.) On a classification of periodic maps on surfaces which commute with
 Hiraku Nozawa (Ritsumeikan Univ.) certain involutions

概要 Ishizaka classified up to conjugation orientation preserving periodic maps of a surface which commute with a hyperelliptic involution. Here, an involution on a surface is hyperelliptic if and only if the quotient space is a sphere. In this talk, we consider maps which commute with involutions whose quotient space is a torus. We will classify up to conjugation orientation preserving irreducible periodic maps which commute with such involutions to give a complete list. It turned out that there are only finite conjugacy classes. We present a representative of each conjugacy class with certain decomposition of surfaces into fundamental domains.

- 8 岡田 真央 (東大数理) Local rigidity of certain actions of nilpotent-by-cyclic groups on the
 sphere 10
 Mao Okada (Univ. of Tokyo) Local rigidity of certain actions of nilpotent-by-cyclic groups on the
 sphere

概要 Let $G = \mathrm{SU}(n, 1)$ be the isometry group of complex hyperbolic space X and $G = KAN$ an Iwasawa decomposition. We proved local rigidity of actions of certain finitely generated subgroups Γ of AN on the imaginary boundary of X .

14:15~15:15 特別講演

- 河澄 響 矢 (東大数理) 写像類群, Goldman–Turaev Lie 双代数, 柏原 Vergne 問題
 Nariya Kawazumi (Univ. of Tokyo) Mapping class groups, the Goldman–Turaev Lie bialgebra and the Kashiwara–Vergne problem

概要 Let Σ be a compact connected oriented surface with non-empty boundary and a framing f . Then a subset of the mapping class group of Σ , which includes the Torelli group, is naturally embedded into the (completed) Goldman–Turaev Lie bialgebra of Σ . A framed version of the Turaev cobracket vanishes on the image of the embedding. So we need a formal description of the Goldman–Turaev Lie bialgebra. In the genus 0 case, the set of expansions inducing a formal description of the bialgebra is naturally bijective to the set of solutions of the Kashiwara–Vergne problem in the formulation of Alekseev–Torossian. In view of this bijection, we can formulate a Kashiwara–Vergne problem associated with (Σ, f) . The set of its solutions is non-empty except some of the genus 1 case. This talk is based on a joint work with Anton Alekseev, Yusuke Kuno and Florian Naef.

15:30~17:30

- 9 野崎 雄太 (東大数理) Every lens space contains a genus one homologically fibered knot 15
 Yuta Nozaki (Univ. of Tokyo) Every lens space contains a genus one homologically fibered knot

概要 We prove that every lens space contains a genus one homologically fibered knot, which is contrast to the fact that some lens spaces contain no genus one fibered knot. In the proof, the discriminant of a binary quadratic form and the Chebotarev density theorem in number theory play central roles.

- 10 藤井 道彦 (琉球大理) The spherical growth series for certain Seifert fiber spaces 10
 Michihiko Fujii (Univ. of Ryukyus) The spherical growth series for certain Seifert fiber spaces

概要 We consider groups presented as $G(p, q) = \langle x, y \mid x^p = y^q \rangle$, with integers p and q satisfying $2 \leq p \leq q$. The groups are geometrically realized as the fundamental groups of Seifert fiber spaces over 2-dimensional disks with two cone points. We present rational function expressions for the spherical growth series of such groups with respect to the generating set $\{x, y, x^{-1}, y^{-1}\}$.

- 11 北野 晃朗 (創価大理工) 結び目のスライス和で与えられるホモロジー 3 球面の $SL(2; \mathbb{C})$ -既約表現の Reidemeister torsion の値の有限性について 10
 Teruaki Kitano (Soka Univ.) Finiteness of the values of Reidemeister torsion for a spliced sum along knots with $SL(2; \mathbb{C})$ -irreducible representations

概要 Let M be a homology 3-sphere. Reidemeister torsion is a topological invariant of M with a representation $\rho : \pi_1(M) \rightarrow SL(2; \mathbb{C})$. It gives a complex valued function on the space of conjugacy classes of $SL(2; \mathbb{C})$ -irreducible representations. We show the image is a finite set for some splicing manifolds along torus knots or the figure-eight knot.

- 12 山田 裕一 (電通大) Exceptional Dehn surgeries along the Mazur link 10
 Yuichi Yamada (Univ. of Electro-Comm.) Exceptional Dehn surgeries along the Mazur link

概要 A hyperbolic 3-manifold that has torus boundaries can change to a non-hyperbolic 3-manifold by filling the boundary by a solid torus. Such surgeries are called exceptional surgeries. We study the exceptional Dehn surgeries along the Mazur link and Akbulut–Yasui links.

- 13 斎藤 敏夫 (上越教育大) An approach to defining Hempel distance of generalized Heegaard splittings 10
Toshio Saito (Joetsu Univ. of Edu.) An approach to defining Hempel distance of generalized Heegaard splittings

概要 It is known that Hempel distance of Heegaard splittings has a good relationship with topology and geometry of 3-manifolds. We discuss an approach to defining Hempel distance of generalized Heegaard splittings.

- 14 境 圭一 (信州大理) Generic 閉曲線の一般化された連結和に対する Arnold 不変量の公式 ... 15
杉山 龍太郎
Keiichi Sakai (Shinshu Univ.) Generalized connected sum formula for the Arnold invariants of generic
Ryutaro Sugiyama plane curves

概要 We define the generalized connected sum for generic closed plane curves, generalizing the strange sum defined by Arnold, and completely describe how the Arnold invariants J^\pm and St behave under the generalized connected sums.

- 15 石上 茜 (東海大理) Type 1 の lens を含む (2,4,2) 型の minimal 5-chart について 15
Akane Ishigami (Tokai Univ.) Minimal 5-charts of type (2,4,2) with a lens of type 1

概要 Let Γ be a chart. For each label k , we denote by Γ_k the union of edges of label k and their vertices. Let Γ be a chart. If Γ has exactly 8 white vertices and $\Gamma_m \cap \Gamma_{m+1}$ has 2 white vertices, $\Gamma_{m+1} \cap \Gamma_{m+2}$ has 4 white vertices, $\Gamma_{m+2} \cap \Gamma_{m+3}$ has 2 white vertices, then Γ is called a chart of type (2,4,2). In this talk we study for a minimal 5-chart of type (2,4,2) with a lens of type 1 which has less than or equal to 14 black vertices.

- 16 中村伊南沙 (東大数理) 1-ハンドル付加による被覆曲面結び目の単純化 10
Inasa Nakamura (Univ. of Tokyo) Simplifying covering surface-knots by an addition of 1-handles with
chart loops

概要 We consider a surface-knot in the form of a simple branched covering over an oriented surface-knot F , which is called a covering surface-knot over F . Such a surface-knot is presented by a certain graph called a chart on a surface diagram of F . For a covering surface-knot, an addition of 1-handles with chart loops is a simplifying operation which deforms the chart to a union of free edges and 1-handles with chart loops. Here, we obtain properties of such simplifications.

- 17 佐藤 進 (神戸大理) リボン曲面タンゲルと曲面絡み目のダブル 10
Shin Satoh (Kobe Univ.) Ribbon surface-tangles and doubles of surface-links

概要 A ribbon surface-tangle is a compact surface in upper four-space with no minimal points such that the boundary presents a trivial link. Any surface-link is the closure of some ribbon surface-tangle. We introduce an equivalence relation among ribbon surface-tangles, called a root-equivalence, and prove that the closures of two ribbon surface-tangles present the same surface-link if and only if they are root-equivalent. As an application, we see that the double of a surface-link is well-defined up to stable equivalence. We also study several properties of the double.

9月12日(火) 第Ⅲ会場

10:10~10:25 2017年度日本数学会幾何学賞授賞式

10:30~11:30 2017年度日本数学会幾何学賞受賞特別講演 (幾何学学科会と合同)

小林 治

Weyl のゲージ理論, Schwarz 微分, そしてある球面定理

Kobayashi Osamu

Weyl's gauge theory, the Schwarzian derivative and a sphere theorem

概要 For many years I have been interested in conformal differential geometry and projective differential geometry. In this talk I would like to explain what these geometries mean at the present day and what can be expected in the future. I am now, as of June 2017, planning the talk and it will be concerned chiefly with a new conformal invariant which is similar to Yamabe's conformal invariant in many respects. In addition a conjecture, a sphere theorem, will be presented. If time permits I touch upon projective differential geometry and discuss some complements to Weyl's setting. This talk as a whole is a derivation from H. Weyl's "Reine Infinitesimalgeometrie" and subsequent developments by K. Yano, H. Yamabe and M. Obata.

13:00~14:00 2017年度日本数学会幾何学賞受賞特別講演 (幾何学学科会と合同)

作間 誠 (広島大理) 3次元多様体のファイバー曲面とヘガード曲面

Makoto Sakuma (Hiroshima Univ.) Fiber surfaces vs Heegaard surfaces of 3-manifolds

概要 Though fiber surfaces and Heegaard surfaces have completely different natures, we can find various analogies between them. We describe the analogies from the view points of (1) the branched fibration theorem, (2) monodromy groups, (3) McShane's identity and (4) geometric structures.

9月13日(水) 第Ⅲ会場

10:00~12:00

18 宮澤 治子

交差の多重化から得られるウェルデッド絡み目の不変量 10

(津田塾大数学・計算機科学研)

和田 康載 (早大教育)

安原 晃 (津田塾大学芸)

Miyazawa Haruko (Tsuda Coll.)

Invariants of welded links derived from multiplexing of crossings

Kodai Wada (Waseda Univ.)

Akira Yasuhara (Tsuda Coll.)

概要 For an ordered n -component link diagram D , we construct a virtual link diagram $D(m_1, m_2, \dots, m_n)$ which is obtained from D by multiplexing of the crossings of D , where m_i is an integer. If two link diagrams D and D' are equivalent, then $D(m_1, m_2, \dots, m_n)$ and $D'(m_1, m_2, \dots, m_n)$ are equivalent as welded links. Since an invariant of $D(m_1, m_2, \dots, m_n)$ is that of D , we try to find new invariants of D via $D(m_1, m_2, \dots, m_n)$.

19 中西 康剛 (神戸大理)

A set of local moves generating the writhe polynomial 10

佐藤 進 (神戸大理)

Yasutaka Nakanishi (Kobe Univ.)

A set of local moves generating the writhe polynomial

Shin Satoh (Kobe Univ.)

概要 In the talk, we will consider a relationship between an invariant and local moves for virtual knots. We show that two virtual knots have the same writhe polynomial if and only if they are related by a finite sequence of certain local moves.

- 20 嘉藤桂樹 (東工大) Interior polynomial for signed bipartite graphs and the HOMFLY polynomial 15
 Keiju Kato (Tokyo Tech) Interior polynomial for signed bipartite graphs and the HOMFLY polynomial

概要 There are many relations between graph theory and knot theory. In particular, certain knot invariants have been expressed in terms of graph invariants. As an example, the interior polynomial is an invariant of bipartite graphs, and a part of the HOMFLY polynomial of a special alternating link coincides with the interior polynomial of the Seifert graph of the link. We extend the interior polynomial to signed bipartite graphs, and we show that, in the planar case, it is equal to a part of the HOMFLY polynomial of a naturally associated link. We also establish some other, more basic properties of this new notion. This leads to new identities involving the original interior polynomial.

- 21 中村拓司 (大阪電通大工) 仮想結び目のある半順序について 10
 中西康剛 (神戸大理)
 佐藤進 (神戸大理)
 Takuji Nakamura On a partial order of virtual knots
 (Osaka Electro-Comm. Univ.)
 Yasutaka Nakanishi (Kobe Univ.)
 Shin Satoh (Kobe Univ.)

概要 For the set of virtual knots \mathcal{V} , we define a relation \leq as follows. Let K, K' be virtual knots. We write $K' \leq K$ if for any virtual knot diagram D of K , we obtain a virtual knot diagram of K' by replacing several real crossings of D with virtual crossings.

In this talk, we show that (\mathcal{V}, \leq) is a partially ordered set. We also show that any finite subset of \mathcal{V} has an upper bound with respect to this order.

- 22 野坂武史 (東工大) べき単的マグナス展開を用いた Milnor 不変量 10
 小谷久寿
 Takefumi Nosaka (Tokyo Tech) Milnor invariants via unipotent Magnus embeddings
 Hisatoshi Kodani

概要 We develop a diagrammatic computation of the Milnor invariant of links, in terms of central group extensions and unipotent Magnus embeddings. As a corollary, we compute the invariants of the Milnor link and of some links. We also powerfully extend the higher invariant, by improving indeterminacy therein.

- 23 野坂武史 (東工大) 自由群のべき零商の群コサイクル 10
 Takefumi Nosaka (Tokyo Tech) Cocycles of nilpotent quotients of free groups

概要 We focus on the cohomology of the k -th nilpotent quotient of the free group, F/F_k . This paper describes all the group 2-, 3-cocycles in terms of Massey products, and gives expressions for some of the 3-cocycles. We also give simple proofs of some of the results on Milnor invariants and the Johnson–Morita homomorphisms.

- 24 A. T. Tran (Univ. of Texas, Dallas) Higher-dimensional twisted Alexander invariants for metabelian representations 10
 山口祥司 (秋田大教育文化)
 Anh T. Tran (Univ. of Texas, Dallas) Higher-dimensional twisted Alexander invariants for metabelian representations
 Yoshikazu Yamaguchi (Akita Univ.)

概要 We will discuss the asymptotic behavior of the twisted Alexander invariants for higher-dimensional representations of a knot group and the relation to the asymptotic behavior of the Reidemeister torsion. We will focus metabelian representations of a knot group into $SL(2; \mathbb{C})$. It is known that metabelian representations are related to exceptional surgeries along a knot. We will see the relation to our previous result on the asymptotic behavior of the Reidemeister torsion for the resulting manifold obtained by an exceptional surgery along a twist knot.

- 25 伊藤昇 (東大数理) On n -trivialities of classical and virtual knots for some unknotting operations 10
 櫻井みぎ和 (茨城工高専)
 Noboru Ito (Univ. of Tokyo) On n -trivialities of classical and virtual knots for some unknotting operations
 Migiwa Sakurai (Ibaraki Nat. Coll. of Tech.)

概要 In 1990, Vassiliev introduced a filtered space of knot invariants via a standard unknotting operation, called crossing change. In 2000, Goussarov, Polyak, and Viro introduced another degree and filtration via another unknotting operation, called virtualization, for classical and virtual knots. In these theories, a notion of n -trivialities has played a significant role. However, for an integer $n (> 2)$, any example of n -trivial classical and virtual knot by virtualizations is still missing. In this talk, we obtain an example of n -trivial knots by virtualizations. We also introduce a new filtration of Vassiliev-type invariants by using an unknotting operation, called Forbidden moves. We obtain n -trivial knots of this new degree.

- 26 滝岡英雄 (阪市大数学研) 自明な $(2, 1)$ ケーブル Γ 多項式をもつ結び目の無限族について 10
 Hideo Takioka (Osaka City Univ.) Infinitely many knots with the trivial $(2, 1)$ -cable Γ -polynomial

概要 It is known that there exist many polynomial invariants for knots. For example, Alexander–Conway, Jones, Γ , Q , HOMFLYPT, Kauffman polynomials are well known. These polynomials of the trivial knot are one. The problem is whether there exists a non-trivial knot such that these polynomials are one. It is known that there exists such a knot for the Alexander–Conway, Γ , Q polynomials. However, it is still an open problem for the other polynomial invariants. Moreover, we consider the $(p, 1)$ -cable versions of these polynomial invariants for an integer $p (\geq 2)$. These $(p, 1)$ -cable versions of the trivial knot are one. The problem is whether there exists a non-trivial knot such that these $(p, 1)$ -cable versions are one. It is known that there exists such a knot for the Alexander–Conway polynomial. However, it is still an open problem for the other polynomial invariants. In this talk, we show that there exist infinitely many knots such that the $(2, 1)$ -cable version of the Γ -polynomial for the knots is one.

14:15~15:15 特別講演

- 内藤 貴仁 (東大数理) Sullivan's coproduct on the reduced loop homology
 Takahito Naito (Univ. of Tokyo) Sullivan's coproduct on the reduced loop homology

概要 In the theory of string topology initiated by Chas and Sullivan, the homology of free loop spaces of manifolds (called the loop homology) has very rich algebraic structures. The loop product is a multiplication on the loop homology and it is the most basic operation in string topology. Cohen and Godin discovered a 2-dimensional TQFT structure on the homology. Moreover, Godin showed that the loop homology is a homological conformal field theory. In this talk, I will discuss a coproduct on the reduced loop homology which is introduced by Sullivan. The coproduct and the loop product give the loop homology an infinitesimal bialgebra structure. I will explain how to construct Sullivan's coproduct and give some computational examples.

15:30~17:30

- 27 西村 悠矢 (京都産大理) Leinster の弱高次元圏におけるモデル圏構造について 15
 Yuya Nishimura (Kyoto Sangyo Univ.) On model structures of Leinster's weak higher categories.

概要 Model structures are known to be important to explore various higher categories. The topic of this talk is canonical model structures on weak higher categories.

- 28 奥山 真吾 (香川高専)* 部分的モノイドにラベルをもつ区間の配置空間 15
 島川 和久 (岡山大理)
 Shingo Okuyama Configuration space of intervals with partially summable labels
 (Kagawa Nat. Coll. of Tech.)
 Kazuhisa Shimakawa (Okayama Univ.)

概要 A configuration space of intervals in 1-dimensional Euclidean space with partially summable labels is constructed. It is a kind of an extension of the configuration space with partially summable labels constructed by the second author and at the same time a generalization of the configuration space of intervals with labels in a based space constructed by the first author. An approximation theorem of the preceding configuration space is generalized to our case. More precisely, we construct a configuration space of intervals in \mathbb{R} with labels in a partial abelian monoid M , and show that it is weakly homotopy equivalent to the space of based loops on the classifying space of M .

- 29 一木 俊助 (横浜国大環境情報) Generic linear perturbations 15
 Shunsuke Ichiki (Yokohama Nat. Univ.) Generic linear perturbations

概要 In his celebrated paper "Generic projections", John Mather has shown that almost all linear projections from a submanifold of a vector space into a subspace yield a stable mapping in the nice dimensions. In this talk, an improvement of the Mather result is given. Namely, almost all linear perturbations of a smooth mapping from a submanifold of \mathbb{R}^m into \mathbb{R}^ℓ yield a stable mapping in the nice dimensions.

- 30 E. B. Batista (Federal Univ. of Cariri) Stability of C^∞ convex integrands 15
 韓 呼 和 (横浜国大環境情報)
 西 村 尚 史 (横浜国大環境情報)
 Erica Boizan Batista (Federal Univ. of Cariri) Stability of C^∞ convex integrands
 Huhe Han (Yokohama Nat. Univ.)
 Takashi Nishimura (Yokohama Nat. Univ.)

概要 In this talk, it is shown that the set consisting of stable convex integrands $S^n \rightarrow \mathbb{R}_+$ is open and dense in the set consisting of C^∞ convex integrands with respect to Whitney C^∞ topology. Moreover, it is given examples representing well why stable convex integrands are preferred.

- 31 清 水 達 郎 (京大数理研) 写像の特異点集合の自己交差と不足符号数 10
 Tatsuro Shimizu (Kyoto Univ.) Self-intersection of singular sets of maps and signature defect

概要 We give a geometric proof of that the k -times self-intersection of singular set of a generic smooth map from n -dimensional manifold X to R^p coincides with the corank (of Jacobian) = k singular set of any generic map from X to R^{p+k-1} as homology classes with $Z/2$ coefficient ($(n-p+2)k > n+1$). As an application we give a description of the signature defect of framed 3-manifold from the point of view of singular sets of maps.

- 32 那 須 正 和 * Essentially weakly onesided resolving endomorphisms of the shift 15
 Masakazu Nasu Essentially weakly onesided resolving endomorphisms of the shift

概要 Characterizations are given for essentially weakly onesided resolving endomorphisms of subshifts.

- 33 那 須 正 和 * On the limits of onesided resolving directions of endomorphisms of sub-
 shifts of finite type 15
 Masakazu Nasu On the limits of onesided resolving directions of endomorphisms of sub-
 shifts of finite type

概要 We present the results that there exists an automorphism of a full shift having a limit of onesided resolving directions of type II or III and that no automorphism of a transitive subshift of finite type has an irrational unique non-expansive direction.

9月14日(木) 第III会場

10:00~11:30

- 34 加 藤 久 男 (筑波大数理物質) トポロジカルエントロピーと1次元連続体の幾何学的構造 15
 Hisao Kato (Univ. of Tsukuba) Topological entropy and topological structures of 1-dimensional continua

概要 In this talk, we define a new notion of “free tracing property by free chains” on G -like continua and we prove that a positive topological entropy homeomorphism on a G -like continuum admits a Cantor set Z such that every tuple of finite points in Z is an IE -tuple of f and Z has the free tracing property by free chains. Also, by use of this notion, we prove the following theorem: If G is any graph and a homeomorphism f on a G -like continuum X has positive topological entropy, then there is a Cantor set Z which is related to both the chaotic behaviors of Kerr and Li in dynamical systems and composants of indecomposable continua in topology. This theorem implies that chaotic dynamics induce complicated topology.

- 35 下村尚司 (名経大経済) Topological rank does not increase by natural extension of Cantor minimals 10
 Takashi Shimomura Topological rank does not increase by natural extension of Cantor minimals
 (Nagoya Univ. of Economics)

概要 Downarowicz and Maass (2008) defined the topological rank for all homeomorphic Cantor minimal dynamical systems. This definition can be extended to all continuous surjective Cantor minimal systems. We have made it clear that taking natural extension does not increase the topological rank.

- 36 下村尚司 (名経大経済) Proximal Cantor systems with topological rank 2 are residually scrambled 10
 Takashi Shimomura Proximal Cantor systems with topological rank 2 are residually scrambled
 (Nagoya Univ. of Economics)

概要 Downarowicz and Maass (2008) introduced the topological rank on all homeomorphic Cantor minimal dynamical systems. This definition can be easily extended to homeomorphic Cantor proximal dynamical systems. We consider the homeomorphic proximal Cantor dynamical systems with topological rank 2. We show that they are all residually scrambled. Evidently, such systems have at most two ergodic measures. We have obtained a necessary and sufficient condition for the unique ergodicity of these systems. In addition, we show that the number of ergodic measures of systems that are topologically mixing can be 1 and 2. Moreover, there exist examples that are topologically weakly mixing, not topologically mixing, and uniquely ergodic. Finally, we show that the number of ergodic measures of systems that are not weakly mixing can be 1 and 2.

- 37 越野克久 (神奈川大工) Hyperspaces homeomorphic to Hilbert spaces 15
 Katsuhisa Koshino (Kanagawa Univ.) Hyperspaces homeomorphic to Hilbert spaces

概要 Let $\text{Comp}(X)$ be the hyperspace consisting of non-empty compact subsets of a space X with the Vietoris topology, and $C(X)$ be the hyperspace of compact and connected sets in X , that is considered as a subspace of $\text{Comp}(X)$. In this talk, we characterize a metrizable space X whose hyperspaces $\text{Comp}(X)$ and $C(X)$ are homeomorphic to a non-separable Hilbert space.

- 38 矢ヶ崎達彦 (京都工繊大工芸) Local and end deformation theorems for uniform embeddings 15
 Tatsuhiko Yagasaki (Kyoto Inst. Tech.) Local and end deformation theorems for uniform embeddings

概要 This talk is concerned with local and end deformation properties of spaces of uniform embeddings in metric manifolds. Using the local deformation theorem by Cernavskii and Edwards–Kirby, we show that any metric manifold with a geometric group action has the local deformation property for uniform embeddings (LD). As an example, the κ -cone end ($\kappa \leq 0$) over any compact Lipschitz metric manifold is shown to have the property (LD). We also introduce a notion of end deformation of uniform embeddings (ED) and show that the 0-cone end over any compact Lipschitz metric manifold has the property (ED). A role of uniform isotopies in uniform topology is also clarified.

- 39 山崎薫里 (高崎経大経済) Insertion of poset-valued maps with the way-below and -above relations 10
 Kaori Yamazaki Insertion of poset-valued maps with the way-below and -above relations
 (Takasaki City Univ. of Econ.)

概要 We give insertion theorems for maps with values in bi-bounded complete and bicontinuous posets by using the way-below relation and the way-above relation.

無限可積分系

9月11日(月) 第V会場

9:50~12:00

- 1 筧 三郎 (立教大理) 擬不変量を持つ可解カオスと虚数乗法 15
 久保 涼平 (立教大理)
 Saburo Kakei (Rikkyo Univ.) Solvable chaos with pseudo-invariants and complex multiplication
 Ryohei Kubo (Rikkyo Univ.)

概要 We propose several solvable chaotic systems that have pseudo-invariants. The solution of the systems can be represented by elliptic functions associated to elliptic curves with complex multiplication. We also discuss invariant measures of the systems.

- 2 永野 中行 (東大数理) 保型形式の視点から見た可換な微分作用素 15
 Atsuhira Nagano (Univ. of Tokyo) Differential operators from the viewpoint of automorphic forms

概要 The speaker will present a result of linear ordinary differential equations which are parametrized on Hermitian symmetric space and invariant under the action of symmetric groups. They are generalization of the classical Lamé equation. We will see a relation between such differential equations and automorphic forms for symplectic groups.

- 3 中屋 敷 厚 (津田塾大学芸) KP 階層のtau関数の展開係数について 15
 岡田 聡 一 (名大多元数理)
 執行 洋子 (津田塾大学芸)
 Atsushi Nakayashiki (Tsuda Coll.) On the expansion coefficients of KP Tau function
 Soichi Okada (Nagoya Univ.)
 Yoko Shigyo (Tsuda Coll.)

概要 If the tau function does not vanish at the origin, it is known that the coefficients are given by Giambelli formula and that it characterizes solutions of the KP hierarchy. In this talk, we deal with a generalization of Giambelli formula to the case when the tau function vanishes at the origin.

- 4 小川原 弘士 (熊本大先端科学) 代数的差分方程式の解の代数的独立性について 15
 Hiroshi Ogawara (Kumamoto Univ.) On algebraic independence of solutions to certain algebraic difference equations

概要 In the talk, we give criteria for algebraic independence of solutions to a certain system of first order algebraic difference equations. First, as a background of our results, Ostrowski gave a criterion for algebraic independence of integrals of given functions. Inspired by Ostrowski's result, Kolchin gave a criterion for algebraic independence of exponentials of integrals of given functions. Hardouin gave difference analogues of Ostrowski's result and Kolchin's result, that is, criteria for algebraic independence of functions satisfying first order linear difference equations. We generalize the analogues to more general systems of algebraic difference equations. As an application of our results, we show algebraic independence of multiple gamma functions and derivatives of the gamma function. As another application, we show algebraic independence of the logarithmic function, q -polylogarithm functions and q -exponential functions.

- 5 伊藤公毅 (豊橋技科大) q 差分加群の解層とそのコホモロジー 15
 Kôki Itô (Toyohashi Univ. of Tech.) solution sheaf of q -difference module and its cohomology

概要 Aomoto introduces the global q -difference de Rham complex and the Čech complex. These two complex should be isomorphic to each other in general case. To prove this, we introduce a sheaf-theoretic version of these complexes. We introduce the sheaf $\mathcal{D}_{\mathbb{P}^1}^\bullet$ of q -difference operators, which is a q -analogue of the sheaf of differential operators on \mathbb{P}^1 . We define the sheaf $\mathcal{D}_{\mathbb{P}^1}^\bullet$ on some simplicial space \mathbb{P}^1_\bullet (which we also define). For some $\mathcal{D}_{\mathbb{P}^1}^\bullet$ -module \mathcal{M}^\bullet we consider $\mathbb{R}\mathcal{H}om_{\mathcal{D}_{\mathbb{P}^1}^\bullet}(\mathcal{O}^\bullet, \mathcal{M}^\bullet)$, where \mathcal{O}^\bullet is the structure sheaf on \mathbb{P}^1_\bullet . Both the (global) q -difference de Rham complex and the Čech complex are quasi-isomorphic to $\mathbb{R}\Gamma\mathbb{R}\mathcal{H}om_{\mathcal{D}_{\mathbb{P}^1}^\bullet}(\mathcal{O}^\bullet, \mathcal{M}^\bullet)$.

- 6 大川 領 (早大基幹理工) Functional equations of Nekrasov functions proposed by Ito–Maruyoshi–Okuda 15
 Ryo Okawa (Waseda Univ.) Functional equations of Nekrasov functions proposed by Ito–Maruyoshi–Okuda

概要 We give a proof of functional equations of Nekrasov partition functions for A_1 -singularity, suggested by Ito–Maruyoshi–Okuda. We follow method by Nakajima–Yoshioka based on the theory of wall-crossing formula developed by Mochizuki.

- 7 小西由紀子 (京大理)* 齋藤構造の概双対性と複素鏡映群 15
 三鍋聡司 (東京電機大工)
 白石勇貴 (京大理)
 Yukiko Konishi (Kyoto Univ.) Almost duality for Saito structure and complex reflection groups
 Satoshi Minabe (Tokyo Denki Univ.)
 Yuuki Shiraishi (Kyoto Univ.)

概要 We extend Dubrovin’s almost duality of Frobenius structures to Saito structures without metric. Then we formulate and study the existence and uniqueness problem of the natural Saito structure on the orbit spaces of finite complex reflection groups from the viewpoint of the almost duality. We give a complete answer to the problem for the irreducible groups.

14:20~16:30

- 8 岡田 聡一 (名大多元数理) C 型ルート系に付随した Schur Q 関数 15
 Soichi Okada (Nagoya Univ.) Schur Q -functions associated to the root system of type C

概要 Schur Q -functions are a family of symmetric functions introduced by Schur in his study of projective representations of symmetric groups. They are obtained by putting $t = -1$ in the Hall–Littlewood functions associated to the root system of type A . (Schur functions are the $t = 0$ specialization.) This talk concerns symplectic Q -functions, which are obtained by putting $t = -1$ in the Hall–Littlewood functions associated to the root system of type C . We discuss several Pfaffian identities as well as a combinatorial description for them. Also we present some positivity conjectures.

- 9 伊藤雅彦 (琉球大理) Selberg 型 BC_n 楕円超幾何積分の行列式公式 15
 野海正俊 (神戸大理)
 Masahiko Ito (Univ. of Ryukyus) A determinant formula for the BC_n elliptic hypergeometric integrals of Selberg type
 Masatoshi Noumi (Kobe Univ.)

概要 We will present a determinant formula for the BC_n elliptic hypergeometric integrals of Selberg type. The formula means that the determinant whose entries are the BC_n type elliptic Selberg integrals is expressed explicitly as a product of elliptic gamma functions. For this purpose we make use of the elliptic Lagrange interpolation functions, which play important roles in both definition of the determinant and proof of the product formula.

- 10 竹村 剛一 (中大理工) Degenerations of Ruijsenaars–van Diejen operator and q -Painlevé equations 15

Kouichi Takemura (Chuo Univ.) Degenerations of Ruijsenaars–van Diejen operator and q -Painlevé equations

概要 It is known that the Painlevé VI is obtained by connection preserving deformation of some linear differential equations, and the Heun equation is obtained by a specialization of the linear differential equations. We investigate degenerations of the Ruijsenaars–van Diejen difference operators and show difference analogues of the Painlevé–Heun correspondence.

- 11 増田 哲 (青学大理工) 4階笹野系の q -類似たち 15

Tetsu Masuda (Aoyama Gakuin Univ.) q -analogues of the Sasano system of the fourth order

概要 The Sasano system is a higher order generalization of the sixth Painlevé equations. The Sasano system of the $2N$ -th order admits the affine Weyl group symmetry of type $D_{2N+2}^{(1)}$ as the group of Bäcklund transformations. We propose, in this talk, some q -analogues of the Sasano system of the fourth order. Each of our systems also admits the affine Weyl group symmetry.

- 12 鈴木 貴雄 (近畿大理工) $W(A_{2n+1}^{(1)})$ -対称性を持つ一般化 q -パンルヴェ VI 方程式に付随する双線型関係式 15

Takao Suzuki (Kinki Univ.) A bilinear form of the generalized q -Painlevé VI system with $W(A_{2n+1}^{(1)})$ symmetry

概要 The q -Painlevé VI equation (q - P_{VI}) was introduced by Jimbo and Sakai. Recently, from a viewpoint of the Heine’s basic hypergeometric function ${}_{n+1}\phi_n$, we proposed a higher order generalization of q - P_{VI} , which admits an extended affine Weyl group symmetry of type $A_{2n+1}^{(1)}$. In this talk, we formulate τ -functions for the generalized q - P_{VI} on the root lattice $Q(A_{2n+1})$ and show that they satisfy Hirota–Miwa type bilinear relations.

- 13 大久保直人 (東大数理) q -パンルヴェ方程式の退化とクラスター代数 15
鈴木 貴雄 (近畿大理工)

Naoto Okubo (Univ. of Tokyo) Degeneration of q -discrete Painlevé equations and cluster algebras
Takao Suzuki (Kinki Univ.)

概要 We construct cluster algebras the coefficients of which satisfy several q -discrete Painlevé equations. These cluster algebras are obtained from quivers with the mutation-period property. We will also show that degeneration of q -discrete Painlevé equations corresponds to confluence of quivers.

- 14 長尾 秀人 (明石工高専) $E_7^{(1)}$ 型 q 差分パンルヴェ方程式の種々の特異点配置 15
山田 泰彦 (神戸大理工)

Hidehito Nagao (Akashi Coll. of Tech.) Various singularity configurations for q -Painlevé equations of type $E_7^{(1)}$
Yasuhiko Yamada (Kobe Univ.)

概要 When we geometrically construct the Painlevé systems by certain rational surfaces called the spaces of initial values, we can variously select deformation directions and singularity configurations in $\mathbb{P}^1 \times \mathbb{P}^1$. However, in order to obtain a simple evolution equation, it is necessary to select the deformation direction and coordinates (singularity configuration) suitably. In this talk, we give some examples of simple evolution equations for the case of q - $E_7^{(1)}$ type, and we discuss a relation among them and their special solutions.

- 15 竹 縄 知 之 (東京海洋大海洋工) 2次元可積分写像のファイバー依存非自励化と離散 Painlevé 方程式 15
 Tomoyuki Takenawa Fiber-dependent deautonomization of integrable 2D mappings and discrete Painlevé equations
 (Tokyo Univ. of Marine Sci. and Tech.)

概要 In this talk we establish a way of performing the deautonomization for a pair of an autonomous mapping and an anti-canonical divisor. Starting from a single autonomous mapping but varying the type of a chosen fiber, we obtain different types of discrete Painlevé equations using this deautonomization procedure. We also introduce a technique which allows us to obtain factorized expressions of discrete Painlevé equations, including the elliptic case. Further, by imposing certain restrictions on such non-autonomous mappings we obtain new and simple elliptic difference Painlevé equations, including examples whose symmetry groups do not appear explicitly in Sakai's classification. (Based on joint work with A. S. Carstea and A. Dzhamay.)

16:40~17:40 特別講演

- 上 岡 修 平 (京 大 情 報) Partition functions for reverse plane partitions derived from the two-dimensional Toda molecule
 Shuhei Kamioka (Kyoto Univ.) Partition functions for reverse plane partitions derived from the two-dimensional Toda molecule

概要 Reverse plane partitions are combinatorial objects of tableau-type. One of the most important topics in reverse (and ordinary) plane partitions is the exploration of generating functions and partition functions which can be nicely factored. In this talk we show a close connection between reverse plane partitions and an integrable dynamical system called the discrete two-dimensional (2D) Toda molecule. We consider reverse plane partitions of arbitrary shape and of bounded size of parts and find that each non-vanishing solution to the discrete 2D Toda molecule gives a partition function which can be nicely factored. For the result a combinatorial interpretation of the discrete 2D Toda molecule in terms of lattice paths is crucial. As an instance from a specific solution we obtain a new partition function which generalizes both MacMahon's generating function for boxed plane partitions and the trace generating function for reverse plane partitions of unbounded size of parts.

9月12日(火) 第V会場

9:50~12:00

- 16 茂 木 康 平 (東京海洋大海洋工) 6頂点模型の Izergin–Korepin 解析 10
 Kohei Motegi Izergin–Korepin analysis on the wavefunctions of the six-vertex models
 (Tokyo Univ. of Marine Sci. and Tech.)

概要 We analyze the wavefunctions of the six-vertex models by extending the celebrated Izergin–Korepin analysis which was originally invented by Korepin and Izergin to study the domain wall boundary partition functions. We show that the method can give a systematic way to study the relation between the wavefunctions and symmetric functions. We mainly illustrate the method by taking the basic wavefunctions of the XXZ-type six-vertex models as an example. Next, we show the result for the case of triangular boundary conditions, Felderhof models and their generalizations such as the elliptic analogue.

- 17 桑野泰宏 (鈴鹿医療科学大医工) イジング模型: 2次元磁場付きの系と3次元系 15
 Yas-Hiro Quano (Suzuka Univ. of Med. Sci.) Ising model: 2dim system in a magnetic field and 3 dim system

概要 We consider the two dimensional Ising model by using the CTM (corner transfer matrix) method. We derive the corner Hamiltonian for the Ising model in a magnetic field by perturbatively inserting the magnetic field. This is just a perturbative approximation because the existence of a real magnetic field violates some assumptions for the validity of the CTM method. Thus we find that the CTM method goes well at a particular pure imaginary magnetic field, and obtain the explicit expression of the spontaneous magnetization.

- 18 小島武夫 (山形大理工) 量子アフィン超代数 $U_q(\widehat{sl}(M|N))$ の自由場表現 15
 Takeo Kojima (Yamagata Univ.) A bosonization of the quantum affine superalgebra $U_q(\widehat{sl}(M|N))$

概要 A bosonization of the quantum affine superalgebra $U_q(\widehat{sl}(M|N))$ is presented for an arbitrary level $k \in \mathbf{C}$. Screening operators that commute with $U_q(\widehat{sl}(M|N))$ are presented for the level $k \neq -M + N$. This talk is based on arXiv.1701.03645 to appear in *Commun. Math. Phys.*

- 19 齋藤洋介 (阪市大数学研) フェルミオンと量子トーラス代数の楕円化 15
 Yosuke Saito (Osaka City Univ.) Elliptic analog of an algebra of fermions and the quantum torus algebra

概要 Nakatsu–Takasaki have studied partition functions of five-dimensional gauge theories and topological string theories by using an algebra of fermions. Recently, it has been realized that partition functions of six-dimensional gauge theories are represented by elliptic functions. As an attempt to apply the method due to Nakatsu–Takasaki to the six-dimensional cases, the author has introduced an elliptic analog of an algebra of fermions. An elliptic analog of the quantum torus algebra is also constructed.

- 20 大久保勇輔 (ロシア HSE) Ding–Iohara–Miki 代数のレベル N 表現の特異ベクトル 15
 Yusuke Ohkubo (Nat. Res. Univ. Higher School of Econ.) Singular vectors of the level N representation of Ding–Iohara–Miki algebra

概要 In this talk, I explain that singular vectors of a certain algebra arising from the level N representation of a Hopf algebra called Ding–Iohara–Miki algebra correspond to generalized Macdonald functions (also called AFLT basis or fixed point basis). This correspondence is a generalization of coincidence between singular vectors of the deformed W algebra and ordinary Macdonald functions. Moreover, we also obtain a formula for Kac determinant of the level N representation of DIM algebra.

- 21 Ivan Chi Ho Ip (京大理) On tensor product decomposition of positive representations 15
 Ivan Chi Ho Ip (Kyoto Univ.) On tensor product decomposition of positive representations

概要 Recently a cluster realization of Drinfeld–Jimbo quantum groups $U_q(\mathfrak{g})$ has been found via the positive representations, where an embedding into a quantum torus algebra $\mathcal{A}_{\mathfrak{g}}$ is described by certain quiver diagram. Using this new realization, we discuss its application towards the proof of the tensor product decomposition of the positive representations of split real quantum groups restricted to the Borel part, as well as the proof by Schrader–Shapiro on the full decomposition in type A_n , thus solving part of the long-standing conjecture of the closure of positive representations under taking the tensor product.

22 今野 均 (東京海洋大海洋工) 楕円量子群と楕円重み関数 15

Hitoshi Konno Elliptic quantum group and elliptic weight function

(Tokyo Univ. of Marine Sci. and Tech.)

概要 By using representation theory of the elliptic quantum group $U_{q,p}(\widehat{\mathfrak{sl}}_N)$, we present a systematic method of deriving the weight functions. The resultant \mathfrak{sl}_N type elliptic weight functions are new, and give elliptic and dynamical analogue of those obtained by Mimachi and Tarasov–Varchenko in the trigonometric case. We then discuss some basic properties of the elliptic weight functions. We also present an explicit formula for formal elliptic hypergeometric integral solution to the elliptic q -KZ equation.

23 今野 均 (東京海洋大海洋工) 楕円量子群の Gelfand–Tsetlin 基底上の有限次元表現 15

Hitoshi Konno Finite-dimensional representations of elliptic quantum group on the

(Tokyo Univ. of Marine Sci. and Tech.) Gelfand–Tsetlin basis

概要 We construct a finite-dimensional representation of the elliptic quantum group $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ on the Gelfand–Tsetlin basis. The resultant representation is described in terms of the partitions of $[1, n]$ in a combinatorial way, and gives an elliptic and dynamical analogue of the geometric representation of $U_q(\widehat{\mathfrak{sl}}_N)$ on the equivariant K -theory constructed by Vasserot and Nakajima. By comparing the elliptic weight functions presented in the previous talk and the elliptic stable envelopes proposed by Aganagic–Okounkov, we conjecture that our finite-dimensional representation provides a geometric representation of $U_{q,p}(\widehat{\mathfrak{sl}}_N)$ on the equivariant elliptic cohomology.

13:00~14:00 特別講演

粟田 英資 (名大多元数理) Ding–庵原–三木代数の表現論

Hidetoshi Awata (Nagoya Univ.) The representation theory of the Ding–Iohara–Miki algebra

概要 We will discuss on the recent development of the representation theory of the quantum toroidal algebra of type $\mathfrak{gl}(1)$ i.e. the Ding–Iohara–Miki algebra.