

Two-weighted estimates for positive operators and Doob maximal operators on filtered measure spaces

By Wei CHEN, Chunxiang ZHU, Yahui ZUO and Yong JIAO

(Received Mar. 6, 2018)
(Revised Jan. 14, 2019)

Abstract. We characterize strong type and weak type inequalities with two weights for positive operators on filtered measure spaces. These estimates are probabilistic analogues of two-weight inequalities for positive operators associated to the dyadic cubes in \mathbb{R}^n due to Lacey, Sawyer and Uriarte-Tuero [30]. Several mixed bounds for the Doob maximal operator on filtered measure spaces are also obtained. In fact, Hytönen-Pérez type and Lerner-Moen type norm estimates for Doob maximal operator are established. Our approaches are mainly based on the construction of principal sets.

1. Introduction

The theory of weighted inequalities in harmonic analysis is an old subject, which can probably be traced back to the beginning of integration. The A_p condition first appeared in a paper of Rosenblum [42], but systematic investigation was initiated by [36], [9] and [37] etc. The A_p condition is geometric, meaning to only involve the weights and not the operators. Later, Sawyer [43] introduced the test condition S_p and characterized the two-weight estimates for the classical Hardy-Littlewood maximal operator. The testing condition essentially amounts to testing the uniform estimates on characteristic functions of dyadic cubes. In addition, Sawyer [44] proved that for operators such as fractional integrals, Poisson kernels, and other nonnegative kernels, the two-weight estimate still holds if one assumes the testing condition not only on the operator itself, but also on its formal adjoint (see [14] and [15] for more information).

Dyadic Harmonic Analysis can be traced back to the early years of the 20th century, and Haar's basis of orthogonal functions has profound and still useful connections to combinatorial and probabilistic reasoning. This subject has recently acquired a renewed attention by Stefanie Petermichl [41], that a notion of Haar shifts can be used to recover deep results about the Hilbert transform (see [38] and [27] for more information). As is well known, to get sharp one-weight estimates of usual operators in classical harmonic analysis, a standard way is a dyadic discretization technique. Using it, Hytönen [16] gave the solution of the A_2 conjecture, which states that any Calderón-Zygmund operator satisfies the following bound on weighted Lebesgue spaces:

$$(1.1) \quad \|T\|_{L^p(w)} \lesssim [w]_{A_p}^{\max(1, \frac{1}{p-1})}.$$

2010 Mathematics Subject Classification. Primary 60G46; Secondary 60G42.

Key Words and Phrases. Martingale, Positive operator, Doob maximal operator, Weighted inequality.

Its simpler proofs were found by several authors (see [19, 32]) and inequality (1.1) has seen several improvements (see [33, 18, 21, 28]). These improvements come in the form of the so-called mixed estimates. The idea behind the mixed estimates is that one only needs the full strength of the A_p constant for part of the estimates, while the other part only requires something weaker. The smaller quantities come in the form of A_r constants for large r or A_∞ constants. The dyadic discretization technique is also valid for (linear) positive operators (see [29, 30, 24, 25, 50]) and the (fractional) maximal operator (see [4, 43, 31, 29, 17, 21]).

With the development of weighted theory in harmonic analysis, its probabilistic counterpart was also studied. This is weighted theory on martingale spaces. The history of martingale theory goes back to the early 1950s when Doob [13] pointed out the connection between martingales and analytic functions. Standard introductions to martingale theory can be found in Dellacherie and Meyer [11], Doob [12], Kazamaki [26], Long [34], Neveu [39], Weisz [52] and Williams [53]. Recently, Schilling [45] and Stroock [46] developed martingale theory for σ -finite measure spaces rather than just for probability spaces, so that they are immediately applicable to analysis on the Euclidean space \mathbb{R}^n without the need of auxiliary truncations or decompositions into probability spaces. Doob's maximal operator, which is a generalization of the dyadic Hardy-Littlewood maximal operator, and a martingale transform, which is an analogue of a singular integral in classical harmonic analysis, are important tools in stochastic analysis. For Doob's maximal operator, assuming some regularity condition on A_p weights, one-weight inequality was studied first by Izumisawa and Kazamaki [22]. The added property is superfluous (see Jawerth [23] or Long [34]). Two-weight weak inequalities were studied by Uchiyama [51] and Long [34], and two-weight strong inequalities were studied by Long and Peng [35] and Chang [6]. Weighted inequalities involving Carleson measure for generalized Doob's maximal operator were obtained by Chen and Liu [8].

In martingale theory, as we see above, weighted inequalities first appeared in 1970s, but they have been developing slowly. One reason is that some decomposition theorems and covering theorems which depend on algebraic structure and topological structure are invalid on probability space. Recently, there are two new approaches to weighted theory in martingale spaces. One is very closely related to Burkholder's method (see [5]). This is the so-called Bellman's method, which also rests on the construction of an appropriate special function. The technique has been used very intensively mostly in analysis, in the study of Carleson embedding theorems, BMO estimates, square function inequalities, bounds for maximal operators, estimates for weights and many other related results. For more complete references, we refer to the bibliographies of [49]. In martingale spaces, this theory was further developed in a series of papers by Bañuelos and Osękowski (see, e.g., [1, 2, 3]) and a monograph [40] by Osękowski. The other is the construction of principal sets on filtered measure spaces which is a quadruplet $(\Omega, \mathcal{F}, \mu; (\mathcal{F}_i)_{i \in \mathbb{Z}})$. The germ of principal sets appeared as the sparse family on \mathbb{R}^n (see [21, 10] for more information) and the principal sets were successfully constructed on filtered measure spaces in [47, p.942-943]. Using the construction, Tanaka and Terasawa [48] obtained a characterization for the boundedness of positive operators on filtered measure spaces. In addition, the construction was reinvestigated by Chen and Jiao [7] and a new property of the construction was found (see Section 3, P.3).

The purpose of this paper is to develop a theory of weights for positive operators

and Doob maximal operators on filtered measure spaces. To better explain our aim, we first recall the main results of [30]. Let $\nu = \{\nu_Q : Q \in \mathcal{Q}\}$ be non-negative constants associated to dyadic cubes, and define a positive linear operator by

$$T_\nu f = \sum_{Q \in \mathcal{Q}} \nu_Q \mathbb{E}_Q f \cdot \chi_Q,$$

where $\mathbb{E}_Q f := |Q|^{-1} \int_Q f dx$. Let σ, w be non-negative locally integral weights on \mathbb{R}^n . Lacey, Sawyer and Uriarte-Tuero [30, Theorem 1.11] characterize the two-weight strong type inequalities

$$(1.2) \quad \|T_\nu(f\sigma)\|_{L^q(w)} \lesssim \|f\|_{L^p(\sigma)}, \quad 1 < p \leq q < \infty,$$

in term of Sawyer-type testing conditions. In the present paper, we consider the positive operator $T_\alpha(\cdot, \sigma)$ (see Subsection 2.1 for the definition) on filtered measure spaces which is the generalization of positive dyadic operator $T_\nu(\cdot, \sigma)$

The following theorem is our first main result, which characterizes two-weight strong type inequality for positive operators on filtered measure spaces. Let p' be the conjugate exponent number of $1 < p < \infty$. All other unexplained notations can be found in Section 2 and Section 3.

THEOREM 1.1. *Let $1 < p \leq q < \infty$. Let $\omega \in A_1$ and $\sigma \in A_1$. Then the following statements are equivalent:*

(1) *There exists a positive constant C such that*

$$(1.3) \quad \|T_\alpha(f\sigma, g\omega)\|_{L^1(d\mu)} \leq C \|f\|_{L^p(\sigma)} \|g\|_{L^{q'}(\omega)};$$

(2) *There exist positive constants C_1 and C_2 such that for any $E \in \mathcal{F}_i^0, i \in \mathbb{Z}$,*

$$(1.4) \quad \left(\int_E \left(\sum_{j \geq i} \mathbb{E}_j(\sigma) \alpha_j \right)^q \omega d\mu \right)^{\frac{1}{q}} \leq C_1 \sigma(E)^{\frac{1}{p}},$$

$$(1.5) \quad \left(\int_E \left(\sum_{j \geq i} \mathbb{E}_j(\omega) \alpha_j \right)^{p'} \sigma d\mu \right)^{\frac{1}{p'}} \leq C_2 \omega(E)^{\frac{1}{q'}}.$$

Moreover, we denote the smallest constants C, C_1 and C_2 in (1.3), (1.4) and (1.5) by $\|T_\alpha(\cdot, \sigma)\|, [\omega, \sigma]_{\alpha, q', p'}$ and $[\sigma, \omega]_{\alpha, p, q}$, respectively. Then it follows that $[\omega, \sigma]_{\alpha, q', p'} \leq \|T_\alpha(\cdot, \sigma)\|, [\sigma, \omega]_{\alpha, p, q} \leq \|T_\alpha(\cdot, \sigma)\|$, and

$$\|T_\alpha(\cdot, \sigma)\| \lesssim [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} + [\sigma, \omega]_{\alpha, p, q} [\sigma]_{A_1}.$$

REMARK 1.2. It is clear that $\|T_\alpha(f\sigma, g\omega)\|_{L^1(d\mu)} = \int_\Omega \sum_{i \in \mathbb{Z}} \alpha_i \mathbb{E}_i(f\sigma) \mathbb{E}_i(g\omega) d\mu$. Then

$$\int_\Omega \sum_{i \in \mathbb{Z}} \alpha_i \mathbb{E}_i(f\sigma) \mathbb{E}_i(g\omega) d\mu = \sum_{i \in \mathbb{Z}} \int_\Omega \alpha_i \mathbb{E}_i(f\sigma) \mathbb{E}_i(g\omega) d\mu = \sum_{i \in \mathbb{Z}} \int_\Omega \alpha_i \mathbb{E}_i(f\sigma)(g\omega) d\mu.$$

It follows that

$$\sum_{i \in \mathbb{Z}} \int_{\Omega} \alpha_i \mathbb{E}_i(f\sigma)(g\omega) d\mu = \int_{\Omega} \sum_{i \in \mathbb{Z}} \alpha_i \mathbb{E}_i(f\sigma)(g\omega) d\mu.$$

Thus $\|T_{\alpha}(f\sigma, g\omega)\|_{L^1(d\mu)} = \int_{\Omega} T_{\alpha}(f\sigma)(g\omega) d\mu$.

Since Remark 1.2 and $L^q(\omega) - L^{q'}(\omega)$ duality, the first statement of Theorem 1.1 is equivalent to the fact that the positive operator $T_{\alpha}(\cdot\sigma)$ is bounded from $L^p(\sigma)$ to $L^q(\omega)$, which extends the inequality (1.2). Moreover, in the very special case that $\sigma = 1$, Theorem 1.1 partially improves Tanaka and Terasawa [47, Theorem 1.1]. Indeed, as pointed out in [47, p. 923], the expected conditions are (1.4) and (1.5). However, for some technical reasons, instead of the condition (1.4), they postulate a strong condition (see [47, (1.5)] or Remark 1.3 below).

Recall that Lacey, Sawyer and Uriarte-Tuero [30, Theorem 1.11] studied two-weight inequalities for positive operator associated to the dyadic cubes in \mathbb{R}^n . As is well known, they obtained two characterizations for the boundedness of the positive operator, which were the local one and global one. Treil [50] reinvestigated strong type inequality and obtained a short proof for the part involving the local one. For more information and references, see Tanaka and Terasawa [48]. The arguments in [30] and [50] are related to dyadic technique extensively, so they are invalid in filtered measure spaces. Instead of dyadic technique, our method is mainly based on the construction of principal sets (see Section 3).

REMARK 1.3. Let $\alpha_i, i \in \mathbb{Z}$, be a nonnegative bounded \mathcal{F}_i -measurable function and $\bar{\alpha}_i \in \mathcal{L}^+$, where $\bar{\alpha}_i := \sum_{j \geq i} \alpha_j$. Assuming that

$$(1.6) \quad \mathbb{E}_i \bar{\alpha}_i \approx \bar{\alpha}_i,$$

holds, [47, Theorem 1.1] showed that (1.5) implies (1.3) in the special case $\sigma = 1$.

As a corollary of Theorem 1.1, we have the following one-weight estimate.

COROLLARY 1.4. *Let $1 < p \leq q < \infty$. Then the following statements are equivalent:*

(1) *There exists a positive constant C such that*

$$(1.7) \quad \|T_{\alpha}(f\omega, g\omega)\|_{L^1(d\mu)} \leq C \|f\|_{L^p(\omega)} \|g\|_{L^{q'}(\omega)};$$

(2) *There exist positive constants C_1 and C_2 such that for any $E \in \mathcal{F}_i^0, i \in \mathbb{Z}$,*

$$(1.8) \quad \left(\int_E \left(\sum_{j \geq i} \mathbb{E}_j(\omega) \alpha_j \right)^q \omega d\mu \right)^{\frac{1}{q}} \leq C_1 \omega(E)^{\frac{1}{p}},$$

$$(1.9) \quad \left(\int_E \left(\sum_{j \geq i} \mathbb{E}_j(\omega) \alpha_j \right)^{p'} \omega d\mu \right)^{\frac{1}{p'}} \leq C_2 \omega(E)^{\frac{1}{q'}}.$$

Moreover, we denote the smallest constants C, C_1 and C_2 in (1.7), (1.8) and (1.9) by $\|T_{\alpha}(\cdot\sigma)\|, [\omega, \omega]_{\alpha, q', p'}$ and $[\omega, \omega]_{\alpha, p, q}$, respectively. Then it follows that $[\omega, \omega]_{\alpha, q', p'} \leq$

$\|T_\alpha(\cdot\sigma)\|, [\omega, \omega]_{\alpha, p, q} \leq \|T_\alpha(\cdot\sigma)\|$, and

$$\|T_\alpha(\cdot\sigma)\| \lesssim [\omega, \omega]_{\alpha, q', p'} + [\omega, \omega]_{\alpha, p, q}.$$

If $\omega = 1$, then Corollary 1.4 reduces to the following, which is the main result of [48, Theorem 1.2].

COROLLARY 1.5. *Let $1 < p \leq q < \infty$. Then the following statements are equivalent:*

(1) *There exists a positive constant C such that*

$$\|T_\alpha(f, g)\|_{L^1(d\mu)} \leq C \|f\|_{L^p(d\mu)} \|g\|_{L^{q'}(d\mu)};$$

(2) *There exists a positive constant C such that for any $E \in \mathcal{F}_i^0, i \in \mathbb{Z}$,*

$$\begin{cases} \left(\int_E \left(\sum_{j \geq i} \alpha_j \right)^q d\mu \right)^{\frac{1}{q}} \leq C \mu(E)^{\frac{1}{p}}, \\ \left(\int_E \left(\sum_{j \geq i} \alpha_j \right)^{p'} d\mu \right)^{\frac{1}{p'}} \leq C \mu(E)^{\frac{1}{q'}}. \end{cases}$$

Our second main result is two-weight weak type inequalities for positive operators in a filtered measure space, which is corresponding to [30, Theorem 1.8].

THEOREM 1.6. *Let $1 < p \leq q < \infty$. Then the following statements are equivalent:*

(1) *There exists a positive constant C such that*

$$(1.10) \quad \|T_\alpha(f\sigma)\|_{L^{q, \infty}(\omega)} \leq C \|f\|_{L^p(\sigma)};$$

(2) *There exists a positive constant C such that for any $E \in \mathcal{F}_i^0, i \in \mathbb{Z}$,*

$$(1.11) \quad \left(\int_E \left(\sum_{j \geq i} \mathbb{E}_j(\omega) \alpha_j \right)^{p'} \sigma d\mu \right)^{\frac{1}{p'}} \leq C \omega(E)^{\frac{1}{q'}}.$$

Moreover, we denote the smallest constants C in (1.10) and (1.11) by $\|T_\alpha(\cdot\sigma)\|$ and $[\sigma, \omega]_{\alpha, p, q}$, respectively. Then it follows that $[\sigma, \omega]_{\alpha, p, q} \leq \|T_\alpha(\cdot\sigma)\| \lesssim [\sigma, \omega]_{\alpha, p, q}$.

We now turn to the Doob maximal operator. We prove several mixed A_p - A_∞ bounds on filtered measure spaces. They are Hytönen-Pérez type and Lerner-Moen type norm estimates; see [21] and [33].

THEOREM 1.7. *Let $1 < p < \infty$.*

(1) *If $(v, \omega) \in B_p$, then $\|M\|_{L^p(v) \rightarrow L^p(\omega)} \lesssim [v, \omega]_{B_p}^{\frac{1}{p}}$;*

(2) *If $(v, \omega) \in A_p$ and $\sigma := \omega^{-\frac{1}{p-1}} \in A_\infty^*$, then $\|M\|_{L^p(v) \rightarrow L^p(\omega)} \lesssim [v, \omega]_{A_p}^{\frac{1}{p}} [\sigma]_{A_\infty^*}^{\frac{1}{p}}$;*

(3) *If $(\omega) \in A_p$ and $\sigma = \omega^{-\frac{1}{p-1}}$, then $\|M\|_{L^p(\omega) \rightarrow L^p(\omega)} \lesssim [\sigma]_{(A_{p'})^{\frac{1}{p'}} (A_\infty^*)^{\frac{1}{p}}} \left(1 + \log_2 [\omega]_{A_p} \right)^{\frac{1}{p}}$.*

Theorem 1.7 (1) and Theorem 1.7 (2) are probabilistic versions of [21, Theorem 4.3]; Theorem 1.7 (3) is closely corresponding to [33, Theorem 1.1]. We mention that the probabilistic analogue of Hytönen-Pérez type estimate [21, Theorem 4.3] first appeared in Tanaka and Terasawa [47, Theorem 5.1]. They gave one-weight norm estimates which is similar to Theorem 1.7 (1). Their estimate has two suprema. In particular, if $\omega = v$ in Theorem 1.7 (1), we obtain a better constant than [47, Theorem 5.1].

The article is organized as follows. In Section 2, we state some preliminaries. We construct principal sets in Section 3. In Section 4, we provide the proofs of the above theorems.

Throughout the paper, the letters C , C_1 and C_2 will be used for constants that may change from one occurrence to another. We use the notation $A \lesssim B$ to indicate that there is a constant C , independent of the weight constant, such that $A \leq CB$. We write $A \approx B$ when $A \lesssim B$ and $B \lesssim A$.

2. Preliminaries

This section consists of the preliminaries for this paper.

2.1. Filtered Measure Space

In this subsection we introduce the filtered measure space, which is standard [47, 20] (see also references therein). Let a triplet $(\Omega, \mathcal{F}, \mu)$ be a measure space. Denote by \mathcal{F}^0 the collection of sets in \mathcal{F} with finite measure. The measure space $(\Omega, \mathcal{F}, \mu)$ is called σ -finite if there exist sets $E_i \in \mathcal{F}^0$ such that $\Omega = \bigcup_{i=0}^{\infty} E_i$. In this paper all measure spaces are assumed to be σ -finite. Let $\mathcal{A} \subset \mathcal{F}^0$ be an arbitrary subset of \mathcal{F}^0 . An \mathcal{F} -measurable function $f : \Omega \rightarrow \mathbb{R}$ is called \mathcal{A} -integrable if it is integrable on all sets of \mathcal{A} , i.e., $\chi_E f \in L^1(\mathcal{F}, \mu)$ for all $E \in \mathcal{A}$. Denote the collection of all such functions by $L^1_{\mathcal{A}}(\mathcal{F}, \mu)$.

If $\mathcal{G} \subset \mathcal{F}$ is another σ -algebra, it is called a sub- σ -algebra of \mathcal{F} . A function $g \in L^1_{\mathcal{G}^0}(\mathcal{G}, \mu)$ is called the conditional expectation of $f \in L^1_{\mathcal{G}^0}(\mathcal{F}, \mu)$ with respect to \mathcal{G} if there holds

$$\int_G f d\mu = \int_G g d\mu, \quad \forall G \in \mathcal{G}^0.$$

The conditional expectation of f with respect to \mathcal{G} will be denoted by $\mathbb{E}(f|\mathcal{G})$, which exists uniquely in $L^1_{\mathcal{G}^0}(\mathcal{G}, \mu)$ due to σ -finiteness of $(\Omega, \mathcal{G}, \mu)$.

A family of sub- σ -algebras $(\mathcal{F}_i)_{i \in \mathbb{Z}}$ is called a filtration of \mathcal{F} if $\mathcal{F}_i \subset \mathcal{F}_j \subset \mathcal{F}$ whenever $i, j \in \mathbb{Z}$ and $i < j$. We call a quadruplet $(\Omega, \mathcal{F}, \mu; (\mathcal{F}_i)_{i \in \mathbb{Z}})$ a σ -finite filtered measure space. It contains a filtered probability space with a filtration indexed by \mathbb{N} , a Euclidean space with a dyadic filtration and doubling metric space with dyadic lattice.

We write

$$\mathcal{L} := \bigcap_{i \in \mathbb{Z}} L^1_{\mathcal{F}_i^0}(\mathcal{F}, \mu).$$

Notice that

$$L^1_{\mathcal{F}_i^0}(\mathcal{F}, \mu) \supset L^1_{\mathcal{F}_j^0}(\mathcal{F}, \mu)$$

whenever $i < j$. For a function $f \in \mathcal{L}$ we will denote $\mathbb{E}(f|\mathcal{F}_i)$ by $\mathbb{E}_i(f)$. By the tower rule of conditional expectations, a family of functions $\mathbb{E}_i(f) \in L^1_{\mathcal{F}_i}(\mathcal{F}, \mu)$ becomes a martingale.

Let $(\Omega, \mathcal{F}, \mu; (\mathcal{F}_i)_{i \in \mathbb{Z}})$ be a σ -finite filtered measure space. Then a function $\tau : \Omega \rightarrow \{-\infty\} \cup \mathbb{Z} \cup \{+\infty\}$ is called a stopping time if for any $i \in \mathbb{Z}$, we have $\{\tau = i\} \in \mathcal{F}_i$. The family of all stopping times is denoted by \mathcal{T} . Fixing $i \in \mathbb{Z}$, we denote $\mathcal{T}_i = \{\tau \in \mathcal{T} : \tau \geq i\}$.

Suppose that function $f \in \mathcal{L}$, the Doob maximal operator is defined by

$$Mf = \sup_{i \in \mathbb{Z}} |\mathbb{E}_i(f)|.$$

Fix $i \in \mathbb{Z}$, we define the tailed Doob maximal operator by

$$*M_i f = \sup_{j \geq i} |\mathbb{E}_j(f)|.$$

Let $\alpha_i, i \in \mathbb{Z}$, be a nonnegative bounded \mathcal{F}_i -measurable function and set $\alpha = (\alpha_i)$. Let $f, g \in \mathcal{L}$. We define the positive operator $T_\alpha(f)$ and bilinear positive operator $T_\alpha(f, g)$ by

$$T_\alpha f := \sum_{i \in \mathbb{Z}} \alpha_i \mathbb{E}_i(f) \text{ and } T_\alpha(f, g) := \sum_{i \in \mathbb{Z}} \alpha_i \mathbb{E}_i(f) \mathbb{E}_i(g),$$

respectively.

2.2. Definitions of Weights

By a weight we mean a nonnegative function which belongs to \mathcal{L} and, by a convention, we will denote the set of all weights by \mathcal{L}^+ . Let $B \in \mathcal{F}$, $\omega \in \mathcal{L}^+$, we always denote $\int_\Omega \chi_B d\mu$ and $\int_\Omega \chi_B \omega d\mu$ by $|B|$ and $|B|_\omega$, respectively. Then we define several kinds of weights.

DEFINITION 2.1. Let v be a weight. We say that the weight v satisfies the condition A_1 , if there exists a positive constant C such that

$$(2.1) \quad \sup_{j \in \mathbb{Z}} \mathbb{E}_j(v) \leq Cv.$$

We denote by $[v]_{A_1}$ the smallest constant C in (2.1).

DEFINITION 2.2. Let v and ω be weights and $1 < p < \infty$. We say that the couple of weights (v, ω) satisfies the condition A_p , if there exists a positive constant C such that

$$(2.2) \quad \sup_{j \in \mathbb{Z}} \mathbb{E}_j(v) \mathbb{E}_j(\omega^{1-p'})^{\frac{p}{p'}} \leq C,$$

where $\frac{1}{p} + \frac{1}{p'} = 1$. We denote by $[v, \omega]_{A_p}$ the smallest constant C in (2.2).

DEFINITION 2.3. Let ω be a weight and $1 < p < \infty$. We say that the weight ω satisfies the condition A_p , if there exists a positive constant C such that

$$(2.3) \quad \sup_{j \in \mathbb{Z}} \mathbb{E}_j(\omega) \mathbb{E}_j(\omega^{1-p'})^{\frac{p}{p'}} \leq C,$$

where $\frac{1}{p} + \frac{1}{p'} = 1$. We denote by $[\omega]_{A_p}$ the smallest constant C in (2.3).

DEFINITION 2.4. Let ω be a weight. We say that the weight ω satisfies the condition A_∞^{exp} , if there exists a positive constant C such that

$$(2.4) \quad \sup_{j \in \mathbb{Z}} \mathbb{E}_j(\omega) \exp \mathbb{E}_j(\log \omega^{-1}) \leq C.$$

We denote by $[\omega]_{A_\infty^{\text{exp}}}$ the smallest constant C in (2.4).

DEFINITION 2.5. Let v and ω be weights and $1 < p < \infty$. Denote $\sigma = \omega^{-\frac{1}{p-1}} \in \mathcal{L}^+$. We say that the couple of weights (v, ω) satisfies the condition S_p^* , if

$$(2.5) \quad [v, \omega]_{S_p^*} := \sup_{i \in \mathbb{Z}, E \in \mathcal{F}_i^0} \left(\frac{\int_E {}^*M_i(\sigma \chi_E)^p v d\mu}{\sigma(E)} \right)^{\frac{1}{p}} < \infty.$$

DEFINITION 2.6. Let v and ω be weights and $1 < p < \infty$. Denote that $\sigma = \omega^{-\frac{1}{p-1}} \in \mathcal{L}^+$. We say that the couple of weights (v, ω) satisfies the condition B_p , if there exists a positive constant C such that for all $i \in \mathbb{Z}$ we have

$$(2.6) \quad \mathbb{E}_i(v) \mathbb{E}_i(\sigma)^p \leq C \exp \left(\mathbb{E}_i(\log(\sigma)) \right).$$

We denote by $[v, \omega]_{B_p}$ the smallest constant C in (2.6).

DEFINITION 2.7. Let ω be a weight. We say that the weight ω satisfies the condition A_∞^* , if there exists a positive constant C such that for all $i \in \mathbb{Z}$ and $E \in \mathcal{F}_i^0$ we have

$$(2.7) \quad \int_E {}^*M_i(\omega \chi_E) d\mu \leq C \omega(E).$$

We denote by $[\omega]_{A_\infty^*}$ the smallest constant C in (2.7).

REMARK 2.8. We summarize basic properties about the conditions. Let $\omega \in A_p$ and $\sigma = \omega^{1-p'}$. Then

1. $\sigma \in A_{p'}$ and $[\sigma]_{A_{p'}}^{\frac{1}{p'}} = [\omega]_{A_p}^{\frac{1}{p}}$;
2. $\omega \in A_\infty^{\text{exp}}$ and $[\omega]_{A_\infty^{\text{exp}}} \leq [\omega]_{A_p}$;
3. $\omega \in A_\infty^*$ and $[\omega]_{A_\infty^*} \lesssim [\omega]_{A_\infty^{\text{exp}}}$.

Following from Remark 2.8, we give the mixed condition $(A_{p'})^{\frac{1}{p'}} (A_\infty^*)^{\frac{1}{p}}$ by

$$(2.8) \quad [\sigma]_{(A_{p'})^{\frac{1}{p'}} (A_\infty^*)^{\frac{1}{p}}} := \sup_{i \in \mathbb{Z}, Q \in \mathcal{F}_i^0} \left(\text{esssup}_Q (\mathbb{E}(\omega | \mathcal{F}_i) \mathbb{E}(\sigma | \mathcal{F}_i)^{p-1}) \frac{\int_Q {}^*M_i(\sigma \chi_Q) d\mu}{|Q|} \right)^{\frac{1}{p}}.$$

3. Construction of principal sets

We mention that “the construction of principal sets” here first appeared in Tanaka and Terasawa [47], and we find a new property P. 3 of the construction. We repeat the

construction of principal sets here for the convenience of our checking the new property P. 3. We call this property P. 3 conditional sparsity. Our results are mainly based on the construction of principal sets and the conditional sparsity.

Let $i \in \mathbb{Z}$, $h \in \mathcal{L}^+$. Fixing $k \in \mathbb{Z}$, we define a stopping time

$$\tau := \inf\{j \geq i : \mathbb{E}(h|\mathcal{F}_j) > 2^{k+1}\}.$$

For $\Omega_0 \in \mathcal{F}_i^0$, we denote that

$$(3.1) \quad P_0 := \{2^{k-1} < \mathbb{E}(h|\mathcal{F}_i) \leq 2^k\} \cap \Omega_0,$$

and assume $\mu(P_0) > 0$. It follows that $P_0 \in \mathcal{F}_i^0$. We write $\mathcal{K}_1(P_0) := i$ and $\mathcal{K}_2(P_0) := k$. We let $\mathcal{P}_1 := \{P_0\}$ which we call the first generation of principal sets. To get the second generation of principal sets we define a stopping time

$$\tau_{P_0} := \tau\chi_{P_0} + \infty\chi_{P_0^c},$$

where $P_0^c = \Omega \setminus P_0$. We say that a set $P \subset P_0$ is a principal set with respect to P_0 if it satisfies $\mu(P) > 0$ and there exist $j > i$ and $l > k + 1$ such that

$$\begin{aligned} P &= \{2^{l-1} < \mathbb{E}(h|\mathcal{F}_j) \leq 2^l\} \cap \{\tau_{P_0} = j\} \cap P_0 \\ &= \{2^{l-1} < \mathbb{E}(h|\mathcal{F}_j) \leq 2^l\} \cap \{\tau = j\} \cap P_0. \end{aligned}$$

Noticing that such j and l are unique, we write $\mathcal{K}_1(P) := j$ and $\mathcal{K}_2(P) := l$. We let $\mathcal{P}(P_0)$ be the set of all principal sets with respect to P_0 and let $\mathcal{P}_2 := \mathcal{P}(P_0)$ which we call the second generalization of principal sets.

We now need to verify that

$$\mu(P_0) \leq 2\mu(E(P_0))$$

where

$$E(P_0) := P_0 \cap \{\tau_{P_0} = \infty\} = P_0 \cap \{\tau = \infty\} = P_0 \setminus \bigcup_{P \in \mathcal{P}(P_0)} P.$$

Indeed, we have

$$\begin{aligned} \mu(P_0 \cap \{\tau_{P_0} < \infty\}) &\leq 2^{-k-1} \int_{P_0 \cap \{\tau_{P_0} < \infty\}} \mathbb{E}(h|\mathcal{F}_{\tau_{P_0}}) d\mu \\ &= 2^{-k-1} \int_{P_0} \mathbb{E}(h|\mathcal{F}_{\tau_{P_0}}) \chi_{\{\tau_{P_0} < \infty\}} d\mu \\ &= 2^{-k-1} \int_{P_0} \sum_{j \geq i} \mathbb{E}(h|\mathcal{F}_{\tau_{P_0}}) \chi_{\{\tau_{P_0} = j\}} d\mu \\ &= 2^{-k-1} \int_{P_0} \sum_{j \geq i} \mathbb{E}(h|\mathcal{F}_j) \chi_{\{\tau_{P_0} = j\}} d\mu. \end{aligned}$$

It follows that

$$\begin{aligned}
\mu(P_0 \cap \{\tau_{P_0} < \infty\}) &\leq 2^{-k-1} \int_{P_0} \mathbb{E}_i \left(\sum_{j \geq i} \mathbb{E}(h \chi_{\{\tau_{P_0}=j\}} | \mathcal{F}_j) \right) d\mu \\
&= 2^{-k-1} \int_{P_0} \sum_{j \geq i} \mathbb{E}_i(h \chi_{\{\tau_{P_0}=j\}}) d\mu \\
&= 2^{-k-1} \int_{P_0} \mathbb{E}_i(h \chi_{\{\tau_{P_0} < \infty\}}) d\mu \\
&\leq 2^{-k-1} \int_{P_0} \mathbb{E}_i(h) d\mu \leq \frac{1}{2} \mu(P_0).
\end{aligned}$$

This clearly implies

$$\mu(P_0) \leq 2\mu(E(P_0)).$$

For any $P'_0 \in (P_0 \cap \mathcal{F}_i^0)$, there exists a set $\Omega''_0 \in \mathcal{F}_i^0$ such that

$$P'_0 = P_0 \cap \Omega''_0 = \{2^{k-1} < \mathbb{E}(h | \mathcal{F}_i) \leq 2^k\} \cap \Omega_0 \cap \Omega''_0.$$

Taking $\Omega'_0 = \Omega_0 \cap \Omega''_0$, we have $P'_0 = \{2^{k-1} < \mathbb{E}(h | \mathcal{F}_i) \leq 2^k\} \cap \Omega'_0$. Using Ω'_0 instead of Ω_0 in (3.1), we deduce that

$$\mu(P'_0) \leq 2\mu(E(P'_0)).$$

Moreover, we obtain that

$$\begin{aligned}
\int_{P'_0} \chi_{P_0} d\mu &= \mu(P'_0 \cap P_0) = \mu(P'_0) \leq 2\mu(E(P'_0)) = 2\mu(P'_0 \cap \{\tau = \infty\}) \\
&= 2\mu(P'_0 \cap P_0 \cap \{\tau = \infty\}) = 2 \int_{P'_0} \chi_{E(P_0)} d\mu \\
&= 2 \int_{P'_0} \mathbb{E}_i(\chi_{E(P_0)}) d\mu.
\end{aligned}$$

Since P'_0 is arbitrary, we have $\chi_{P_0} \leq 2\mathbb{E}_i(\chi_{E(P_0)})\chi_{P_0}$.

The next generalizations are defined inductively,

$$\mathcal{P}_{n+1} := \bigcup_{P \in \mathcal{P}_n} \mathcal{P}(P),$$

and we define the collection of principal sets \mathcal{P} by

$$\mathcal{P} := \bigcup_{n=1}^{\infty} \mathcal{P}_n.$$

It is easy to see that the collection of principal sets \mathcal{P} satisfies the following properties:

P.1 The sets $E(P)$ where $P \in \mathcal{P}$, are disjoint and $P_0 = \bigcup_{P \in \mathcal{P}} E(P)$;

P.2 $P \in \mathcal{F}_{\mathcal{K}_1(P)}$;

$$\text{P.3 } \chi_P \leq 2\mathbb{E}(\chi_{E(P)}|\mathcal{F}_{\mathcal{K}_1(P)})\chi_P;$$

$$\text{P.4 } 2^{\mathcal{K}_2(P)-1} < \mathbb{E}(h|\mathcal{F}_{\mathcal{K}_1(P)}) \leq 2^{\mathcal{K}_2(P)} \text{ on } P;$$

$$\text{P.5 } \sup_{j \geq i} \mathbb{E}_j(h\chi_P) \leq 2^{\mathcal{K}_2(P)+1} \text{ on } E(P);$$

$$\text{P.6 } \chi_{\{\mathcal{K}_1(P) \leq j < \tau(P)\}} \mathbb{E}_j(h) \leq 2^{\mathcal{K}_2(P)+1}.$$

We use the principal sets to represent the tailed Doob maximal operator and obtain the following lemma.

LEMMA 3.1. *Let $i \in \mathbb{Z}$ and $h \in \mathcal{L}^+$. Fixing $k \in \mathbb{Z}$ and $\Omega_0 \in \mathcal{F}_i^0$, we denote*

$$P_0 := \{2^{k-1} < \mathbb{E}(h|\mathcal{F}_i) \leq 2^k\} \cap \Omega_0.$$

If $\mu(P_0) > 0$, then

$$\begin{aligned} {}^*M_i(h)\chi_{P_0} &= {}^*M_i(h\chi_{P_0})\chi_{P_0} \\ &= \sum_{P \in \mathcal{P}} {}^*M_i(h\chi_{P_0})\chi_{E(P)} \\ &\leq 4 \sum_{P \in \mathcal{P}} 2^{(\mathcal{K}_2(P)-1)}\chi_{E(P)}. \end{aligned}$$

The following lemma is a Carleson embedding theorem associated with the collection of principal sets \mathcal{P} , which is essentially [48, Lemma 2.2]. We provide a different proof.

LEMMA 3.2. *We have*

$$\sum_{P \in \mathcal{P}} \mu(P)2^{p(\mathcal{K}_2(P)-1)} \leq 2(p')^p \|h\chi_{P_0}\|_{L^p(d\mu)}^p.$$

Proof of Lemma 3.2.

$$\begin{aligned} \sum_{P \in \mathcal{P}} \mu(P)2^{p(\mathcal{K}_2(P)-1)} &\leq \sum_{P \in \mathcal{P}} \int_P \mathbb{E}(h\chi_{P_0}|\mathcal{F}_{\mathcal{K}_1(P)})^p d\mu \\ &= \sum_{P \in \mathcal{P}} \int_P \mathbb{E}(h\chi_{P_0}|\mathcal{F}_{\mathcal{K}_1(P)})^p \chi_P d\mu. \end{aligned}$$

Combining it with P.3 of the construction of principal sets, we have

$$\begin{aligned} \sum_{P \in \mathcal{P}} \mu(P)2^{p(\mathcal{K}_2(P)-1)} &\leq 2 \sum_{P \in \mathcal{P}} \int_P \mathbb{E}(h\chi_{P_0}|\mathcal{F}_{\mathcal{K}_1(P)})^p \mathbb{E}(\chi_{E(P)}|\mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &\leq 2 \sum_{P \in \mathcal{P}} \int_P \mathbb{E}(h\chi_{P_0}|\mathcal{F}_{\mathcal{K}_1(P)})^p \chi_{E(P)} d\mu \\ &= 2 \sum_{P \in \mathcal{P}} \int_{E(P)} \mathbb{E}(h\chi_{P_0}|\mathcal{F}_{\mathcal{K}_1(P)})^p d\mu \end{aligned}$$

In the view of the definition of Doob's maximal operator, we have

$$\sum_{P \in \mathcal{P}} \mu(P) 2^{p(\mathcal{K}_2(P)-1)} \leq 2 \sum_{P \in \mathcal{P}} \int_{E(P)} (M(h\chi_{P_0}))^p d\mu \leq \int_{\Omega} (M(h\chi_{P_0}))^p d\mu.$$

It follows from boundedness of Doob's maximal operator that

$$\sum_{P \in \mathcal{P}} \mu(P) 2^{p(\mathcal{K}_2(P)-1)} \leq 2(p')^p \|h\chi_{P_0}\|_{L^p(d\mu)}^p.$$

□

The following lemma can be found in [47, Theorem 4.1] or [8, Theorem 3.2].

LEMMA 3.3. *Let v, ω be weights, $1 < p < \infty$ and $\sigma = \omega^{-\frac{1}{p-1}}$. Then the following statements are equivalent:*

1. *There exists a positive constant C_1 such that*

$$(3.2) \quad \|M(f)\|_{L^p(v)} \leq C_1 \|f\|_{L^p(\omega)},$$

where $f \in L^p(\omega)$;

2. *The couple of weights (v, ω) satisfies the condition S_p^* .*

Moreover, we denote the smallest constant C_1 in (3.2) by $\|M\|$. Then $\|M\| \sim [v, \omega]_{S_p^*}$

4. Proofs of main results

We provide the proofs of our main results in this section. For simplicity we denote operator T_α by T in the proofs of Theorem 1.1 and Theorem 1.6.

Before we give the proof of Theorem 1.1, we mention that our method is similar to that of the proof of the main result in Tanaka and Terasawa [48]. Our new ingredient is the definition of $F_j := \{\mathbb{E}_j^\omega(g)^{q'} \omega \leq \mathbb{E}_j^\sigma(f)^p \sigma\}$, which appears in (4.1). In general F_j is not a \mathcal{F}_i -measurable set. This creates a difficulty in (4.4). To overcome the difficulty, we assume that $\omega \in A_1$ and $\sigma \in A_1$.

When we compare Theorem 1.1 to the local characterization of Lacey, Sawyer and Uriarte-Tuero [30, Theorem 1.11], we do not know whether our assumptions $\omega \in A_1$ and $\sigma \in A_1$ are superfluous on filtered measure spaces. We recall that the proof of [30, Theorem 1.11] depends very much on the dyadic structure. It is clear that our testing condition (1.4) and (1.5) are the generalization of the local characterization of Lacey, Sawyer and Uriarte-Tuero [30, Theorem 1.11]. For the global characterization of [30, Theorem 1.11], we still have no idea to generalize it on filtered measure spaces.

Proof of Theorem 1.1. (1) \Rightarrow (2) is trivial and we omit it. Note that we do not use $\omega \in A_1$ and $\sigma \in A_1$ in this part.

(2) \Rightarrow (1) Let $i \in \mathbb{Z}$ be arbitrarily taken and be fixed. By a standard limiting argument, it suffices to prove that the inequality

$$\sum_{j \geq i} \int_{\Omega} \alpha_j \mathbb{E}_j(f\sigma) \mathbb{E}_j(g\omega) d\mu$$

$$\lesssim [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \|f\|_{L^p(\sigma)}^{p\theta} + [\sigma, \omega]_{\alpha, p, q} [\sigma]_{A_1} \|g\|_{L^{q'}(\omega)}^{q'\theta}, \quad \theta := \frac{1}{p} + \frac{1}{q'},$$

holds (the rest follows from the homogeneity).

We set

$$(4.1) \quad F_j := \{\mathbb{E}_j^\omega(g)^{q'} \omega \leq \mathbb{E}_j^\sigma(f)^p \sigma\} \text{ and } G_j := \Omega \setminus F_j.$$

We shall prove that

$$(4.2) \quad \sum_{j \geq i} \int_{\Omega} \chi_{F_j} \alpha_j \mathbb{E}_j(f \sigma) \mathbb{E}_j(g \omega) d\mu \lesssim [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \|f\|_{L^p(\sigma)}^{p\theta}$$

and

$$(4.3) \quad \sum_{j \geq i} \int_{\Omega} \chi_{G_j} \alpha_j \mathbb{E}_j(f \sigma) \mathbb{E}_j(g \omega) d\mu \lesssim [\sigma, \omega]_{\alpha, p, q} [\sigma]_{A_1} \|g\|_{L^{q'}(\omega)}^{q'\theta}.$$

Since the proofs of (4.2) and (4.3) can be done in a completely symmetric way, we only prove (4.2) in the following.

We estimate $\sum_{j \geq i} \int_E \chi_{F_j} \alpha_j \mathbb{E}_j(f \sigma) \mathbb{E}_j(g \omega) d\mu$ for $E = P_0 \in \mathcal{F}_i^0$, where $\sigma(P_0) > 0$ and, for some $k \in \mathbb{Z}$, $P_0 := \{2^{k-1} < \mathbb{E}_i^\sigma(f) \leq 2^k\}$. For the above i , P_0 , $\sigma d\mu$ and f , we apply the construction of principal sets. Using the principal sets \mathcal{P} , we can decompose the left-hand side of (4.2) as follows:

$$\begin{aligned} \sum_{j \geq i} \int_E \chi_{F_j} \alpha_j \mathbb{E}_j(f \sigma) \mathbb{E}_j(g \omega) d\mu &= \sum_{j \geq i} \int_E \chi_{F_j} \alpha_j \mathbb{E}_j^\sigma(f) \mathbb{E}_j^\omega(g) \mathbb{E}_j(\sigma) \mathbb{E}_j(\omega) d\mu \\ &= \sum_{P \in \mathcal{P}} \sum_{j \geq \mathcal{K}_1(P)} \int_{P \cap \{j < \tau_P\}} \chi_{F_j} \alpha_j \mathbb{E}_j^\sigma(f) \mathbb{E}_j^\omega(g) \mathbb{E}_j(\sigma) \mathbb{E}_j(\omega) d\mu. \end{aligned}$$

Because of $\omega \in A_1$, we have

$$\begin{aligned} &\sum_{j \geq \mathcal{K}_1(P)} \int_{P \cap \{j < \tau_P\}} \chi_{F_j} \alpha_j \mathbb{E}_j^\sigma(f) \mathbb{E}_j^\omega(g) \mathbb{E}_j(\sigma) \mathbb{E}_j(\omega) d\mu \\ (4.4) \quad &\leq 2^{\mathcal{K}_2(P)+1} [\omega]_{A_1} \sum_{j \geq \mathcal{K}_1(P)} \int_{P \cap \{j < \tau_P\}} \alpha_j \mathbb{E}_j(\sigma) \chi_{F_j} \mathbb{E}_j^\omega(g) \omega d\mu \\ &\leq 2^{\mathcal{K}_2(P)+1} [\omega]_{A_1} \sum_{j \geq \mathcal{K}_1(P)} \int_P \alpha_j \mathbb{E}_j(\sigma) \sup_{\mathcal{K}_1(P) \leq j < \tau(P)} (\chi_{F_j} \mathbb{E}_j^\omega(g)) \omega d\mu \\ &= 2^{\mathcal{K}_2(P)+1} [\omega]_{A_1} \int_P \sum_{j \geq \mathcal{K}_1(P)} \alpha_j \mathbb{E}_j(\sigma) \sup_{\mathcal{K}_1(P) \leq j < \tau(P)} (\chi_{F_j} \mathbb{E}_j^\omega(g)) \omega d\mu. \end{aligned}$$

Combining it with Hölder's inequality, we have

$$\sum_{j \geq \mathcal{K}_1(P)} \int_{P \cap \{j < \tau_P\}} \chi_{F_j} \alpha_j \mathbb{E}_j^\sigma(f) \mathbb{E}_j^\omega(g) \mathbb{E}_j(\sigma) \mathbb{E}_j(\omega) d\mu$$

$$\leq 2^{\mathcal{K}_2(P)+1} [\omega]_{A_1} \left(\int_P \left(\sum_{j \geq \mathcal{K}_1(P)} \alpha_j \mathbb{E}_j(\sigma) \right)^q \omega d\mu \right)^{\frac{1}{q}} \left(\int_P \left(\sup_{\mathcal{K}_1(P) \leq j < \tau(P)} (\chi_{F_j} \mathbb{E}_j^\omega(g)) \right)^{q'} \omega d\mu \right)^{\frac{1}{q'}}.$$

In view of the definition of F_j , we obtain

$$\begin{aligned} & \sum_{j \geq \mathcal{K}_1(P)} \int_{P \cap \{j < \tau_P\}} \chi_{F_j} \alpha_j \mathbb{E}_j^\sigma(f) \mathbb{E}_j^\omega(g) \mathbb{E}_j(\sigma) \mathbb{E}_j(\omega) d\mu \\ & \leq [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \left(2^{p(\mathcal{K}_2(P)+1)} \sigma(P) \right)^{\frac{1}{p}} \left(\int_P \left(\sup_{\mathcal{K}_1(P) \leq j < \tau(P)} (\mathbb{E}_j^\sigma(f)) \right)^p \sigma d\mu \right)^{\frac{1}{q'}} \\ & \leq [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \left(2^{p(\mathcal{K}_2(P)+1)} \sigma(P) \right)^{\frac{1}{p}} \left(2^{p(\mathcal{K}_2(P)+1)} \sigma(P) \right)^{\frac{1}{q'}}. \end{aligned}$$

It follows from $\theta = \frac{1}{p} + \frac{1}{q'} \geq 1$ that

$$\begin{aligned} \sum_{j \geq i} \int_E \chi_{F_j} \alpha_j \mathbb{E}_j(f\sigma) \mathbb{E}_j(g\omega) d\mu & \leq [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \sum_{P \in \mathcal{P}} \left(2^{p(\mathcal{K}_2(P)+1)} \sigma(P) \right)^\theta \\ & \leq [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \left(\sum_{P \in \mathcal{P}} 2^{p(\mathcal{K}_2(P)+1)} \sigma(P) \right)^\theta \\ & \lesssim [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \left(\sum_{P \in \mathcal{P}} 2^{p(\mathcal{K}_2(P)-1)} \sigma(P) \right)^\theta. \end{aligned}$$

Using Lemma 3.2, we have

$$(4.5) \quad \sum_{j \geq i} \int_E \chi_{F_j} \alpha_j \mathbb{E}_j(f\sigma) \mathbb{E}_j(g\omega) d\mu \lesssim [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \|f\chi_{P_0}\|_{L^p(\sigma)}^{p\theta}.$$

Note that

$$\begin{aligned} \sum_{j \geq i} \int_\Omega \chi_{F_j} \alpha_j \mathbb{E}_j(f\sigma) \mathbb{E}_j(g\omega) d\mu & = \sum_{j \geq i} \sum_{k \in \mathbb{Z}} \int_{\{2^{k-1} < \mathbb{E}_i^\sigma(f) \leq 2^k\}} \chi_{F_j} \alpha_j \mathbb{E}_j(f\sigma) \mathbb{E}_j(g\omega) d\mu \\ & = \sum_{k \in \mathbb{Z}} \sum_{j \geq i} \int_{\{2^{k-1} < \mathbb{E}_i^\sigma(f) \leq 2^k\}} \chi_{F_j} \alpha_j \mathbb{E}_j(f\sigma) \mathbb{E}_j(g\omega) d\mu. \end{aligned}$$

Combining this with (4.5), we have

$$\begin{aligned} & \sum_{j \geq i} \int_\Omega \chi_{F_j} \alpha_j \mathbb{E}_j(f\sigma) \mathbb{E}_j(g\omega) d\mu \\ & \lesssim [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \sum_{k \in \mathbb{Z}} \left(\int_{\{2^{k-1} < \mathbb{E}_i^\sigma(f) \leq 2^k\}} f^p \sigma d\mu \right)^\theta \\ & \leq [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \left(\sum_{k \in \mathbb{Z}} \int_{\{2^{k-1} < \mathbb{E}_i^\sigma(f) \leq 2^k\}} f^p \sigma d\mu \right)^\theta \\ & = [\omega, \sigma]_{\alpha, q', p'} [\omega]_{A_1} \|f\|_{L^p(\sigma)}^{p\theta}. \end{aligned}$$

Similarly, we obtain

$$\sum_{j \geq i} \int_{\Omega} \chi_{G_j} \alpha_j \mathbb{E}_j(f\sigma) \mathbb{E}_j(g\omega) d\mu \lesssim [\sigma, \omega]_{\alpha, p, q} [\sigma]_{A_1} \|g\|_{L^{q'}(\omega)}^{q'\theta}.$$

This completes the proof of Theorem 1.1. \square

Proof of Corollary 1.4. We change (4.1) to

$$(4.6) \quad F_j := \{\mathbb{E}_j^\omega(g)^{q'} \leq \mathbb{E}_j^\omega(f)^p\} \text{ and } G_j := \Omega \setminus F_j.$$

The proof of Corollary 1.4 is similar to that of Theorem 1.1, and we omit the details. \square

Now we intend to prove two-weight weak type inequality.

Proof of Theorem 1.6. (1) \Rightarrow (2) Note that $\omega \in L^1_{\mathcal{F}^0}$. It follows from duality for Lorentz spaces that

$$\|T(f\omega)\|_{L^{p'}(\sigma)} \leq \|T\| \|f\|_{L^{q',1}(\omega)}.$$

Fix $E \in \mathcal{F}_i^0, i \in \mathbb{Z}$. For $f = \chi_E$, we have

$$\left(\int_E \left(\sum_{j \geq i} \alpha_j \mathbb{E}_j(\omega) \right)^{p'} \sigma d\mu \right)^{\frac{1}{p'}} \leq \|T(f\omega)\|_{L^{p'}(\sigma)} \leq \|T\| \|f\|_{L^{q',1}(\omega)} = \|T\| \omega(E)^{\frac{1}{q'}}.$$

Thus $[\sigma, \omega]_{\alpha, p, q} \leq \|T\|$.

(2) \Rightarrow (1) Fix $f \in L^p(\sigma)$ and $\lambda > 0$. We bound the set $\{T(f\sigma) > 2\lambda\}$. For $n \in \mathbb{Z}$, we denote $T_n(f\sigma) = \sum_{j=-\infty}^{j=n} \alpha_j \mathbb{E}_j(f\sigma)$ and $T^n(f\sigma) = \sum_{j=n}^{\infty} \alpha_j \mathbb{E}_j(f\sigma)$. Let

$$\tau = \inf\{n : T_n(f\sigma) > \lambda\}$$

and $\mathcal{Q}_\lambda = \{\{\tau = n\} : n \in \mathbb{Z}\}$. For $n \in \mathbb{Z}$, we have

$$\lambda \chi_{\{\tau=n\}} \geq T_{n-1}(f\sigma) \chi_{\{\tau=n\}}.$$

Then,

$$\lambda \chi_{\{\tau=n\} \cap \{T(f\sigma) > 2\lambda\}} \leq T^n(f\sigma) \chi_{\{\tau=n\} \cap \{T(f\sigma) > 2\lambda\}}.$$

For $\eta \in (0, 1)$ to be determined later, we denote

$$\mathcal{E} = \left\{ \{\tau = n\} : \omega(\{\tau = n\} \cap \{T(f\sigma) > 2\lambda\}) < \eta \omega(\{\tau = n\}) \right\}$$

and $\mathcal{F} = \mathcal{Q}_\lambda \setminus \mathcal{E}$. It follows that

$$\begin{aligned} & (2\lambda)^q \omega(\{T(f\sigma) > 2\lambda\}) \\ & \leq \eta (2\lambda)^q \sum_{\mathcal{E}} \omega(\{\tau = n\}) + 2^q \lambda^q \eta^{-q} \sum_{\mathcal{F}} \omega(\{\tau = n\}) \left(\frac{\omega(\{\tau = n\} \cap \{T(f\sigma) > 2\lambda\})}{\omega(\{\tau = n\})} \right)^q \end{aligned}$$

$$\leq \eta(2\lambda)^q \sum_{\mathcal{E}} \omega(\{\tau = n\}) + 2^q \eta^{-q} \sum_{\mathcal{F}} \omega(\{\tau = n\}) \left(\frac{\int_{\{\tau=n\}} T^n(f\sigma)\omega d\mu}{\omega(\{\tau = n\})} \right)^q.$$

Note that

$$\begin{aligned} & \sum_{n \in \mathbb{Z}} \omega(\{\tau = n\}) \left(\frac{\int_{\{\tau=n\}} T^n(f\sigma)\omega d\mu}{\omega(\{\tau = n\})} \right)^q \\ &= \sum_{n \in \mathbb{Z}} \left(\int_{\{\tau=n\}} T^n(f\sigma)\omega d\mu \right)^q \omega(\{\tau = n\})^{1-q} \\ &= \sum_{n \in \mathbb{Z}} \left(\int_{\{\tau=n\}} T^n(\omega\chi_{\{\tau=n\}})f\sigma d\mu \right)^q \omega(\{\tau = n\})^{1-q}. \end{aligned}$$

Using Hölder's inequality, we obtain

$$\begin{aligned} & \sum_{n \in \mathbb{Z}} \omega(\{\tau = n\}) \left(\frac{\int_{\{\tau=n\}} T^n(f\sigma)\omega d\mu}{\omega(\{\tau = n\})} \right)^q \\ &\leq \sum_{n \in \mathbb{Z}} \left(\int_{\{\tau=n\}} T^n(\omega\chi_{\{\tau=n\}})^{p'} \sigma d\mu \right)^{\frac{q}{p'}} \left(\int_{\{\tau=n\}} |f|^p \sigma d\mu \right)^{\frac{q}{p}} \omega(\{\tau = n\})^{1-q} \\ &= \sum_{n \in \mathbb{Z}} \left(\left(\int_{\{\tau=n\}} T^n(\omega\chi_{\{\tau=n\}})^{p'} \sigma d\mu \right)^{\frac{1}{p'}} (\omega\{\tau = n\})^{-\frac{1}{q'}} \right)^q \left(\int_{\{\tau=n\}} |f|^p \sigma d\mu \right)^{\frac{q}{p}}. \end{aligned}$$

In view of the condition (1.11), we have

$$\begin{aligned} \sum_{n \in \mathbb{Z}} \omega(\{\tau = n\}) \left(\frac{\int_{\{\tau=n\}} T^n(f\sigma)\omega d\mu}{\omega(\{\tau = n\})} \right)^q &\leq [\sigma, \omega]_{\alpha, p, q}^q \left(\sum_{n \in \mathbb{Z}} \int_{\{\tau=n\}} |f|^p \sigma d\mu \right)^{\frac{q}{p}} \\ &= [\sigma, \omega]_{\alpha, p, q}^q \left(\int_{\Omega} |f|^p \sigma d\mu \right)^{\frac{q}{p}}. \end{aligned}$$

Thus

$$\|T_\alpha(f\sigma)\|_{L^{q, \infty}(\omega)} \leq C(\eta)[\sigma, \omega]_{\alpha, p, q} \|f\|_{L^p(\sigma)},$$

where $C(\eta) = \frac{2}{(1-2^q\eta)^{\frac{1}{q}}\eta}$. The function $C(\eta)$ attains its minimum for $\eta = \frac{q}{1+q} \frac{1}{2^q}$ and the minimum is equal to $2^{q+1} \frac{1+q}{q} (1+q)^{\frac{1}{q}}$. It follows that $\|T\| \lesssim [\sigma, \omega]_{\alpha, p, q}$. \square

Proof of Theorem 1.7. Let $i \in \mathbb{Z}$ be arbitrarily chosen and fixed. By Lemma 3.3, we estimate $\int_E {}^*M_i(\sigma\chi_E)^p v d\mu$ for any $E \in \mathcal{F}_i^0$.

Since

$$\int_E {}^*M_i(\sigma)^p v d\mu = \int_E {}^*M_i(\sigma\chi_E)^p v d\mu,$$

it suffices to estimate $\int_E {}^*M_i(\sigma\chi_E)^p v d\mu$ for $E = P_0 \in \mathcal{F}_i^0$, where $\mu(P_0) > 0$ and, for some $k \in \mathbb{Z}$, $P_0 := \{2^{k-1} < \mathbb{E}(\sigma|\mathcal{F}_i) \leq 2^k\}$.

For the above i , P_0 and σ , we apply the construction of principal sets. We have

$$\begin{aligned} \int_{P_0} {}^*M_i(\sigma)^p v d\mu &\leq \sum_{P \in \mathcal{P}} \int_{E(P)} {}^*M_i(\sigma)^p v d\mu \\ &\lesssim \sum_{P \in \mathcal{P}} \int_{E(P)} 2^{p(\mathcal{K}_2(P)-1)} v d\mu \\ &\leq \sum_{P \in \mathcal{P}} \int_P 2^{p(\mathcal{K}_2(P)-1)} v d\mu. \end{aligned}$$

Proof of (1). It follows from the definition of B_p that

$$\begin{aligned} \int_P 2^{p(\mathcal{K}_2(P)-1)} v d\mu &= \int_P 2^{p(\mathcal{K}_2(P)-1)} \mathbb{E}(v | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &\leq \int_P \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)})^p \mathbb{E}(v | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &\leq [v, \omega]_{B_p} \int_P \exp(\mathbb{E}(\log \sigma | \mathcal{F}_{\mathcal{K}_1(P)})) d\mu. \end{aligned}$$

Note that

$$\begin{aligned} \int_P \exp(\mathbb{E}(\log \sigma | \mathcal{F}_{\mathcal{K}_1(P)})) d\mu &= \int_P \exp(\mathbb{E}(\log(\sigma \chi_{P_0}) | \mathcal{F}_{\mathcal{K}_1(P)})) d\mu \\ &= \int_P \exp(\mathbb{E}(\log(\sigma \chi_{P_0}) | \mathcal{F}_{\mathcal{K}_1(P)})) \chi_P d\mu. \end{aligned}$$

In view of P.3 of the construction of principal sets, it follows that

$$\begin{aligned} \int_P 2^{p(\mathcal{K}_2(P)-1)} v d\mu &\leq 2[v, \omega]_{B_p} \int_P \exp(\mathbb{E}(\log(\sigma \chi_{P_0}) | \mathcal{F}_{\mathcal{K}_1(P)})) \mathbb{E}(\chi_{E(P)} | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &= 2[v, \omega]_{B_p} \int_P \exp(\mathbb{E}(\log(\sigma \chi_{P_0}) | \mathcal{F}_{\mathcal{K}_1(P)})) \chi_{E(P)} d\mu \\ &= 2[v, \omega]_{B_p} \int_{E(P)} \exp(\mathbb{E}(\log(\sigma \chi_{P_0}) | \mathcal{F}_{\mathcal{K}_1(P)})) d\mu. \end{aligned}$$

Using Jensen's inequality for conditional expectation, for any $q > 1$, we have

$$\exp\left(\mathbb{E}(\log(\sigma \chi_{P_0}) | \mathcal{F}_{\mathcal{K}_1(P)})\right) \leq \mathbb{E}\left((\sigma \chi_{P_0})^{\frac{1}{q}} | \mathcal{F}_{\mathcal{K}_1(P)}\right)^q \leq M((\sigma \chi_{P_0})^{\frac{1}{q}})^q.$$

Then

$$\begin{aligned} \int_{P_0} {}^*M_i(\sigma)^p v d\mu &\lesssim [v, \omega]_{B_p} \sum_{P \in \mathcal{P}} \int_{E(P)} M((\sigma \chi_{P_0})^{\frac{1}{q}})^q d\mu \\ &\leq [v, \omega]_{B_p} \int_{P_0} M((\sigma \chi_{P_0})^{\frac{1}{q}})^q d\mu. \end{aligned}$$

Combining it with the boundedness of Doob's maximal operator, we deduce that

$$\int_{P_0} {}^*M_i(\sigma)^p v d\mu \lesssim [v, \omega]_{B_p} (q')^q \int_{P_0} \sigma d\mu.$$

Letting $q \rightarrow \infty$, we obtain $(q')^q \rightarrow e$. Thus

$$\int_{P_0} {}^*M_i(\sigma)^p v d\mu \lesssim [v, \omega]_{B_p} \sigma(P_0).$$

Proof of (2). It follows from the definition of A_p that

$$\begin{aligned} \int_P 2^{p(\mathcal{K}_2(P)-1)} v d\mu &= \int_P 2^{p(\mathcal{K}_2(P)-1)} \mathbb{E}(v | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &\leq \int_P \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)})^p \mathbb{E}(v | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &\leq [v, \omega]_{A_p} \int_P \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu. \end{aligned}$$

Note that $\int_P \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu = \int_P \mathbb{E}(\sigma \chi_{P_0} | \mathcal{F}_{\mathcal{K}_1(P)}) \chi_P d\mu$. In view of P.3 of the construction of principal sets, it follows that

$$\begin{aligned} \int_P \mathbb{E}(\sigma \chi_{P_0} | \mathcal{F}_{\mathcal{K}_1(P)}) \chi_P d\mu &\leq 2 \int_P \mathbb{E}(\sigma \chi_{P_0} | \mathcal{F}_{\mathcal{K}_1(P)}) \mathbb{E}(\chi_{E(P)} | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &= 2 \int_{E(P)} \mathbb{E}(\sigma \chi_{P_0} | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &\leq 2 \int_{E(P)} {}^*M_{\mathcal{K}_1(P_0)}(\sigma \chi_{P_0}) d\mu. \end{aligned}$$

Then

$$\begin{aligned} \int_{P_0} {}^*M_i(\sigma)^p v d\mu &\lesssim [v, \omega]_{A_p} \sum_{P \in \mathcal{P}} \int_{E(P)} {}^*M_{\mathcal{K}_1(P_0)}(\sigma \chi_{P_0}) d\mu \\ &\leq [v, \omega]_{A_p} \int_{P_0} {}^*M_{\mathcal{K}_1(P_0)}(\sigma \chi_{P_0}) d\mu. \end{aligned}$$

Because of $\sigma \in A_\infty^*$, we have

$$\int_{P_0} {}^*M_i(\sigma)^p v d\mu \lesssim [v, \omega]_{A_p} [\sigma]_{A_\infty^*} \sigma(P_0).$$

Proof of (3). For $a \in \mathbb{Z}$, define

$$\mathcal{Q}^a = \{P \in \mathcal{P} : 2^{a-1} < \text{esssup}_P(\mathbb{E}(\omega | \mathcal{F}_{\mathcal{K}_1(P)}) \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)})^{p-1}) \leq 2^a\}.$$

It follows from Hölder's inequality that $1 = \mathbb{E}_j(\omega^{\frac{1}{p}} \omega^{-\frac{1}{p}})^p \leq \mathbb{E}_j(\omega) \mathbb{E}_j(\sigma)^{p-1} \leq [\omega]_{A_p}$, for any $j \in \mathbb{Z}$. Set $K = \lceil \log_2[\omega]_{A_p} \rceil + 1$, we have

$$\mathcal{P} = \bigcup_{a=0}^K \mathcal{Q}^a.$$

Then

$$\begin{aligned} \sum_{P \in \mathcal{P}} \int_P 2^{p(\mathcal{K}_2(P)-1)} \omega d\mu &= \sum_{P \in \mathcal{P}} \int_P 2^{p(\mathcal{K}_2(P)-1)} \mathbb{E}(\omega | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &\leq \sum_{P \in \mathcal{P}} \int_P \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)})^p \mathbb{E}(\omega | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu. \end{aligned}$$

Note that

$$\begin{aligned} &\left(\mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)})^p \mathbb{E}(\omega | \mathcal{F}_{\mathcal{K}_1(P)}) \right) \chi_P \\ &\leq \operatorname{esssup}_P (\mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)})^{p-1} \mathbb{E}(\omega | \mathcal{F}_{\mathcal{K}_1(P)}) \chi_P) \operatorname{esssup}_P \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)}) \chi_P. \end{aligned}$$

It follows that

$$\begin{aligned} \sum_{P \in \mathcal{P}} \int_P 2^{p(\mathcal{K}_2(P)-1)} \omega d\mu &\leq \sum_{a=0}^K 2^a \sum_{P \in \mathcal{Q}^a} \int_P \operatorname{esssup}_P \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &\leq 2 \sum_{a=0}^K 2^a \sum_{P \in \mathcal{Q}^a} \int_P \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &= 2 \sum_{a=0}^K 2^a \sum_{P \in \mathcal{Q}^a} \int_P \sigma d\mu. \end{aligned}$$

Let \mathcal{Q}_{\max}^a be the collection of maximal sets¹ in \mathcal{Q}^a , we obtain

$$\sum_{P \in \mathcal{Q}^a} \int_P \sigma d\mu = \sum_{Q \in \mathcal{Q}_{\max}^a} \sum_{P \in \mathcal{Q}^a, P \subseteq Q} \int_P \sigma d\mu = \sum_{Q \in \mathcal{Q}_{\max}^a} \sum_{P \in \mathcal{Q}^a, P \subseteq Q} \int_P \sigma \chi_P d\mu.$$

In view of P.3 of the construction of principal sets, it follows that

$$\begin{aligned} \sum_{P \in \mathcal{Q}^a} \int_P \sigma d\mu &\leq 2 \sum_{Q \in \mathcal{Q}_{\max}^a} \sum_{P \in \mathcal{Q}^a, P \subseteq Q} \int_P \sigma \mathbb{E}(\chi_{E(P)} | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu \\ &= 2 \sum_{Q \in \mathcal{Q}_{\max}^a} \sum_{P \in \mathcal{Q}^a, P \subseteq Q} \int_P \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)}) \chi_{E(P)} d\mu. \end{aligned}$$

Because of $\int_P \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(P)}) \chi_{E(P)} d\mu = \int_{E(P)} \mathbb{E}(\sigma \chi_Q | \mathcal{F}_{\mathcal{K}_1(P)}) d\mu$, we have

$$\begin{aligned} \sum_{P \in \mathcal{Q}^a} \int_P \sigma d\mu &\leq 2 \sum_{Q \in \mathcal{Q}_{\max}^a} \sum_{P \in \mathcal{Q}^a, P \subseteq Q} \int_{E(P)} {}^*M_{\mathcal{K}_1(Q)}(\sigma \chi_Q) d\mu \\ &\leq 2 \sum_{Q \in \mathcal{Q}_{\max}^a} \int_Q {}^*M_{\mathcal{K}_1(Q)}(\sigma \chi_Q) d\mu. \end{aligned}$$

¹Let $\mathcal{Q} \subset \mathcal{P}$. In view of Zorn's Lemma, for \mathcal{Q} ordered by containment, we have that \mathcal{Q} contains at least one maximal element. Then, we denote the collection of maximal elements in \mathcal{Q} by \mathcal{Q}_{\max} .

Then

$$\begin{aligned} \int_{P_0} {}^*M_i(\sigma)^p \omega d\mu &\lesssim \sum_{a=0}^K 2^a \sum_{Q \in \mathcal{Q}_{\max}^a} \int_Q {}^*M_{\mathcal{K}_1(Q)}(\sigma \chi_Q) d\mu \\ &\lesssim \sum_{a=0}^K \sum_{Q \in \mathcal{Q}_{\max}^a} \operatorname{esssup}_Q \left(\mathbb{E}(\omega | \mathcal{F}_{\mathcal{K}_1(Q)}) \mathbb{E}(\sigma | \mathcal{F}_{\mathcal{K}_1(Q)})^{p-1} \right) \int_Q {}^*M_{\mathcal{K}_1(Q)}(\sigma \chi_Q) d\mu. \end{aligned}$$

By (2.8) the definition of $(A_{p'})^{\frac{1}{p'}} (A_\infty^*)^{\frac{1}{p}}$, we have

$$\begin{aligned} \int_{P_0} {}^*M_i(\sigma)^p \omega d\mu &\lesssim [\sigma]_{(A_{p'})^{\frac{1}{p'}} (A_\infty^*)^{\frac{1}{p}}}^p \sum_{a=0}^K \sum_{Q \in \mathcal{Q}_{\max}^a} \int_Q \sigma d\mu \\ &\leq [\sigma]_{(A_{p'})^{\frac{1}{p'}} (A_\infty^*)^{\frac{1}{p}}}^p \sum_{a=0}^K \int_{P_0} \sigma d\mu \\ &= [\sigma]_{(A_{p'})^{\frac{1}{p'}} (A_\infty^*)^{\frac{1}{p}}}^p (K+1) \int_{P_0} \sigma d\mu. \end{aligned}$$

Thus

$$\begin{aligned} \int_{P_0} {}^*M_i(\sigma)^p \omega d\mu &\lesssim [\sigma]_{(A_{p'})^{\frac{1}{p'}} (A_\infty^*)^{\frac{1}{p}}}^p (3 + \log_2[\omega]_{A_p}) \int_{P_0} \sigma d\mu \\ &\lesssim [\sigma]_{(A_{p'})^{\frac{1}{p'}} (A_\infty^*)^{\frac{1}{p}}}^p (1 + \log_2[\omega]_{A_p}) \int_{P_0} \sigma d\mu. \end{aligned}$$

The proof of Theorem 1.7 is complete. \square

5. Acknowledgement

We thank the referee for many valuable comments and suggestions. These greatly improve the presentation of our results. The article was revised during W. Chen's visit to the Georgia Tech Mathematics Department, whose hospitality is gratefully acknowledged. The research of W. Chen is supported by the National Natural Science Foundation of China(11771379), the Natural Science Foundation of Jiangsu Province(BK20161326), and the Jiangsu Government Scholarship for Overseas Studies(JS-2017-228). The research of Y. Jiao is supported by the National Natural Science Foundation of China (Grant No. 11471337).

References

- [1] Bañuelos, R. and Osękowski A., *Burkholder inequalities for submartingales, Bessel processes and conformal martingales*, Amer. J. Math., **135**(2013), no. 6, 1675–1698.
- [2] Bañuelos, R. and Osękowski A., *Sharp maximal L_p -estimates for martingales*, Illinois J. Math., **58**(2014), no. 1, 149–165.
- [3] Bañuelos, R. and Osękowski A., *Sharp martingale inequalities and applications to Riesz transforms on manifolds, Lie groups and Gauss space*, J. Funct. Anal., **269**(2015), no. 6, 1652–1713.
- [4] Buckley S., *Estimates of operator norms on weighted spaces and reverse Jensen inequalities*, Trans. Amer. Math. Soc., **340**(1993), 253–272.
- [5] Burkholder D. L., *A sharp inequality for martingale transforms*, Ann. Probab., **7**(1979), 858–863.
- [6] Chang X. Q., *Some Sawyer type inequalities for martingales*, Studia Math., **111**(1994), 187–194.

- [7] Chen W. and Jiao Y., *Weighted estimates for the multilinear maximal function on filtered measure spaces*, <https://arxiv.org/abs/1705.11048>(2017).
- [8] Chen W. and Liu P. D., *Weighted inequalities for the generalized maximal operator in martingale spaces*, Chin. Ann. Math. Ser. B, **32**(2011), 781–792.
- [9] Coifman R. R., Fefferman C., *Weighted norm inequalities for maximal functions and singular integrals* Studia Math., **51**(1974), 241–250.
- [10] Damián W., Lerner A. K. and Pérez C., *Sharp weighted bounds for multilinear maximal functions and Calderon-Zygmund operators*, J. Fourier Anal. Appl., **21**(2015), 161–181.
- [11] Dellacherie C. and Meyer P. A., *Probabilities and potential*, North-Holland Publishing Co., Amsterdam, 1988.
- [12] Doob J. L., *Stochastic Processes*, John Wiley & Sons, Inc., New York; Chapman & Hall, Limited, London, 1953.
- [13] Doob J. L., *Semimartingales and subharmonic functions*, Trans. Amer. Math. Soc., **77**(1954), 86–121.
- [14] García-Cuerva J. and Rubio de Francia J. *Weighted norm inequalities and related topics* Volume 116 of North-Holland Mathematics Studies, North Holland Publishing Co., Amsterdam, 1985.
- [15] Grafakos, L., *Classical Fourier Analysis*, Third edition. Graduate Texts in Mathematics, 249. Springer, New York, 2014.
- [16] Hytönen T., *The sharp weighted bound for general Calderón-Zygmund Operators*, Ann. of Math., **175**(2012), 1473–1506.
- [17] Hytönen T. and Kairema A., *Systems of dyadic cubes in a doubling metric space*, Colloq. Math. **126**(2012), 1–33.
- [18] Hytönen T. and Lacey M. T., *The A_p - A_∞ inequality for general Calderón-Zygmund operators*, Indiana U. Math. J., **61**(2012), 2041–2092.
- [19] Hytönen T., Lacey M. T. and Pérez C., *Sharp weighted bounds for the q -variation of singular integrals*, Bull. London Math. Soc., **45**(2013), 529–540.
- [20] Hytönen T., van Neerven J., Veraar M. C. and Weis L. W., *Analysis in Banach spaces Volume I: Martingales and Littlewood-Paley theory*. Ergebnisse der Math. 3. Folge, vol. 63. Springer, Berlin (2016).
- [21] Hytönen T. and Pérez C., *Sharp weighted bounds involving A_∞* , Anal. and PDE., **6**(2013), 777–818.
- [22] Izumisawa M. and Kazamaki N., *Weighted norm inequalities for martingales*, Tôhoku Math. J., **29**(1977), 115–124.
- [23] Jawerth B., *Weighted inequalities for maximal operators: linearization, localization and factorization*, Amer. J. Math., **108**(1986), 361–414.
- [24] Kairema A., *Two-weight norm inequalities for potential type and maximal operators in a metric space*, Publ. Mat., **57**(2013), 3–56.
- [25] Kairema A., *Sharp weighted bounds for fractional integral operators in a space of homogeneous type*, Math. Scand., **114**(2014), 226–253.
- [26] Kazamaki N., *Continuous exponential martingales and BMO*, Lecture Notes in Mathematics, **1579** Springer-Verlag, Berlin, 1994.
- [27] Lacey M. T., *The linear bound in A_2 for Calderón-Zygmund operators: a survey*, Marcinkiewicz centenary volume, 97–114, Banach Center Publ., 95, Polish Acad. Sci. Inst. Math., Warsaw, 2011.
- [28] Lacey M. T., *An A_p - A_∞ inequality for the Hilbert transform*, Houston J. Math., **38**(2012), 799–814.
- [29] Lacey M. T., Moen K., Pérez C. and Torres R. H., *Sharp weighted bounds for fractional integral operators*, J. Funct. Anal. **259**(2010), 1073–1097.
- [30] Lacey M. T., Sawyer E. and Uriarte-Tuero I., *Two weight inequalities for discrete positive operators*, <https://arxiv.org/abs/0911.3437> (2009).
- [31] Lerner A., *An elementary approach to several results on the Hardy-Littlewood maximal operator*, Proc. Amer. Math. Soc., **136**(2008), 2829–2833.
- [32] Lerner A., *A simple proof of the A_2 conjecture*, Int. Math. Res. Not. IMRN, **14**(2013), 3159–3170..
- [33] Lerner A. and Moen K., *Mixed A_p – A_∞ estimates with one supremum*, Studia Math. **219**(2013), 247–267.
- [34] Long R. L., *Martingale spaces and inequalities*, Peking University Press, Beijing; Friedr. Vieweg & Sohn, Braunschweig, 1993.
- [35] Long R.L. and Peng L.Z., *(p, q) maximal inequalities with two weights in martingale theory*, (Chinese) Acta Math. Sinica, **29**(1986), 253–258.
- [36] Muckenhoupt B., *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc., **165**(1972), 207–226.

- [37] Muckenhoupt B. and Wheeden R., *Weighted norm inequalities for fractional integrals*, Trans. Amer. Math. Soc., **192**(1974), 261–274.
- [38] Nazarov F., Treil S. and Volberg A., *The Tb-theorem on non-homogeneous spaces*, Acta Math., **190**(2003), 151–239.
- [39] Neveu J., *Discrete-Parameter Martingales*. North-Holland Publishing Co., Amsterdam, revised edition, 1975.
- [40] Osekowski A., *Sharp Martingale and Semimartingale Inequalities*. Monogr. Mat., vol.72, Birkhäuser, Basel, 2012.
- [41] Petermichl S., *Dyadic shifts and a logarithmic estimate for Hankel operators with matrix symbol*, C. R. Acad. Sci. Paris Sér. I Math., **330**(2000), 455–460.
- [42] Rosenblum M., *Summability of Fourier series in $L_p(d\mu)$* , Trans. Amer. Math. Soc. **105**(1962), 32–42.
- [43] Sawyer E., *A characterization of a two-weight norm inequality for maximal operators*, Studia Math., **75**(1982), 1–11.
- [44] Sawyer E., *A characterization of two weight norm inequalities for fractional and Poisson integrals*, Trans. Amer. Math. Soc., **308**(1988), 533–545.
- [45] Schilling R., *Measures, integrals and martingales*, Cambridge University Press, New York, 2005.
- [46] Stroock D. W., *Probability theory. An analytic view.*, Second edition, Cambridge University Press, Cambridge, 2011.
- [47] Tanaka H. and Terasawa Y., *Positive operators and maximal operators in a filtered measure space*, J. Funct. Anal., **264**(2013), 920–946.
- [48] Tanaka H. and Terasawa Y., *A characterization for the boundedness of positive operators in a filtered measure space*, Arch. Math. (Basel), **101**(2013), 559–568.
- [49] Treil S., *Sharp A_2 estimates of Haar shifts via Bellman function*. Recent trends in analysis, 187–208, Theta Ser. Adv. Math., Theta, Bucharest, 2013.
- [50] Treil S., *A remark on two weight estimates for positive dyadic operators*, in: K. Gröchenig, Yu. Lyubarskii, K. Seip (Eds.), *Operator-Related Function Theory and Time-Frequency Analysis: The Abel Symposium 2012*, in: Abel Symposia, vol. 9, Springer, 2014, pp. 185–195.
- [51] Uchiyama A., *Weight functions on probability spaces*, Tôhoku Math. J., **30**(1978), 463–470.
- [52] Weisz F., *Martingale Hardy spaces and their applications in Fourier analysis*, Lecture Notes in Mathematics, vol. 1568, Springer-Verlag, Berlin, 1994.
- [53] Williams D., *Probability with martingales*. Cambridge Mathematical Textbooks. Cambridge University Press, Cambridge, 1991.

Wei CHEN

School of Mathematical Sciences, Yangzhou University, 225002
Yangzhou, China

E-mail: weichen@yzu.edu.cn

Chunxiang ZHU

School of Mathematical Sciences, Yangzhou University, 225002
Yangzhou, China

E-mail: cxzhu.yzu@163.com

Yahui ZUO

School of Mathematics and Statistics, Central South University,
Changsha 410075, China

E-mail: zuoyahui@csu.edu.cn

Yong JIAO

School of Mathematics and Statistics, Central South University,
Changsha 410075, China

E-mail: jiaoyong@csu.edu.cn